

Supplementary Table 1. Distribution of participants by presence of symptoms and test reactivity


	Non-reactive N=1,838	Reactive N=166	Total N=2,004
Symptomatic			
Fever	20 (1.1)	7 (4.2)	27 (1.4)
Fever & respiratory symptoms*	23 (1.3)	9 (5.4)	32 (1.6)
Only respiratory symptoms*	56 (3.0)	2 (1.2)	58 (2.8)
Asymptomatic or Other Symptoms			
Fatigue	3 (0.2)	0 (0.0)	3 (0.2)
Muscle Ache (Myalgia)	9 (0.5)	0 (0.0)	9 (0.5)
Headache	30 (1.7)	2 (1.4)	32 (1.7)
Nausea/Vomiting	1 (0.1)	0 (0.0)	1 (0.1)
Abdominal Pain	2 (0.1)	0 (0.0)	2 (0.1)
Diarrhea	1 (0.1)	0 (0.0)	1 (0.1)
No Symptoms at All	1,701 (100.0)	146 (98.6)	1,847 (97.8)

*includes cough, sore throat, shortness of breath, chest pain, wheezing, runny nose

Supplementary Table 2: Seroprevalence posterior mean and 95% equal-tailed credible interval estimates based on the second survey for both districts.

Gender	Age	District East	District Malir
Female	0-4	0.139 (0.04, 0.25)	0.080 (0.02, 0.14)
Female	5-9	0.128 (0.04, 0.22)	0.088 (0.04, 0.15)
Female	10-18	0.130 (0.05, 0.21)	0.086 (0.04, 0.14)
Female	19-39	0.149 (0.07, 0.23)	0.101 (0.05, 0.16)
Female	40-59	0.155 (0.07, 0.26)	0.084 (0.03, 0.14)
Female	60+	0.149 (0.05, 0.27)	0.099 (0.04, 0.19)
Male	0-4	0.139 (0.04, 0.25)	0.079 (0.02, 0.14)
Male	5-9	0.128 (0.03, 0.23)	0.082 (0.03, 0.14)
Male	10-18	0.170 (0.09, 0.28)	0.098 (0.04, 0.17)
Male	19-39	0.153 (0.08, 0.24)	0.079 (0.03, 0.13)
Male	40-59	0.217 (0.10, 0.40)	0.083 (0.03, 0.14)
Male	60+	0.145 (0.05, 0.26)	0.087 (0.03, 0.15)

Supplementary Figure 1: Map showing GIS location of enrolled and refused households in the study sites (selected Union Councils of District Malir and District East, Karachi).



Supplementary Figure 2. Comparison of COVID-19 positive cases trend from the District East and study sampling sites with an estimation of seroprevalence in phase 1 and 2

