

SUPPLEMENTARY MATERIAL

Molecular profiling of neonatal dried blood spots reveals changes in innate and adaptive immunity following fetal inflammatory response

Daniel Costa^{1,2}, Nuria Bonet³, Amanda Solé^{2,4}, Jose Manuel González de Aledo⁵, Eduard Sabidó^{2,4,6}, Ferran Casals³, Carlota Rovira⁷, Alfons Nadal⁵, Jose Luis Marin⁵, Teresa Cobo^{5,*}, Robert Castelo^{2,8,*}

¹Department of Pediatrics, Hospital de Figueres, Figueres, Spain; ²Department of Experimental and Health Sciences, Universitat Pompeu Fabra (UPF), Barcelona, Spain; ³Genomics Core Facility, Department of Experimental and Health Sciences, Universitat Pompeu Fabra (UPF), Barcelona, Spain; ⁴Proteomics Unit, Centre de Regulació Genómica (CRG), Barcelona, Spain; ⁵Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Centre for Biomedical Research on Rare Diseases (CIBER-ER), Barcelona, Spain; ⁶Barcelona Institute of Science and Technology (BIST), Barcelona, Spain; ⁷Hospital Sant Joan de Déu; ⁸Research Programme on Biomedical Informatics, Institut Hospital del Mar d'Investigacions Biomèdiques (IMIM), Barcelona, Spain; *Correspondence: Robert Castelo (robert.castelo@upf.edu), Teresa Cobo (TCOBO@clinic.cat)

May 22, 2019

Supplementary Figure 1









		Control	I	II	III	
	Stage					FIR
III		2	4	5	1	
II		1	0	0	0	
I		2	0	0	0	
Control		6	0	0	0	
	MIR					

Figure S1 Pathological examination of the placenta. Number of neonates cross-classified by the diagnosis of the stage of the maternal (MIR) and the fetal (FIR) inflammatory response.

Supplementary Figure 2




Figure S2 Distribution of raw expression values from the US ELGAN cohort microarray data. (a, c) Expression values from background probes. (b, d) Expression values from foreground probes. (a, b) Samples colored by array identifier. (c, d) Samples colored by scanning date.

Supplementary Figure 3




Figure S3 Distribution of background-corrected and quantile-normalized expression values from the US ELGAN cohort microarray data.

Supplementary Figure 4




Figure S4 Sample differences from the US ELGAN cohort microarray data, in terms of fold-changes between samples in base-2 logarithmic scale. (a) Sample labels are colored by storage condition (FZ: frozen; RT: room temperature). (b) Sample labels are colored by microarray scanning date. There is a clear effect of storage condition, an effect of scanning date mostly in frozen samples and no apparent sex or FIR effect.

Supplementary Figure 5




Figure S5 Differential expression between FIR and non-FIR samples from the US ELGAN cohort microarray data. (a) Distribution of raw p-values (b) Quantile-quantile plot of moderated t -statistics.

Supplementary Figure 6




Figure S6 Combined MIR and FIR linear effect on gene expression. Top-20 genes with largest fold change among DE genes with $FDR < 5\%$ and minimum-50% fold change throughout the following combined 4 stages of MIR and FIR: 0 no MIR and no FIR; 1 MIR but no FIR; 2 MIR stage III and FIR stage I; 3 MIR stage III and FIR stage II or III. Adjusted expression values in log-CPM units are plotted in the y -axis as function of the combined stages in the x -axis.