

Supplementary material

Cohort Profile: the Oxford Parkinson's Disease Centre Discovery Cohort Magnetic Resonance Imaging sub-study (OPDC-MRI)

Griffanti et al.




Figure S1. Resting state fMRI: basal ganglia network, part of the template used in dual regression to extract subject-specific maps. Full template available online¹

Discontinued sequences

Supplementary Table S1 illustrates the main acquisition parameters and the number of available subjects per group for each of the sequences that were acquired between 2010 and 2016.

The diffusion-weighted sequence with reduced FOV centered on the substantia nigra (SN) was inspired by the study by Vaillancourt and colleagues², who found reduced FA in the SN of subjects with Parkinson's compared with controls, especially in the caudal compared with the rostral region of interest. We acquired a similar sequence in our first 50 participants, but did not find any difference between Parkinson's and controls (unpublished). We therefore decided to discontinue the sequence.

DESPOT1 and DESPOT2 (driven equilibrium single pulse observation of T1 and T2, respectively) are a set of rapid combined T1 and T2 mapping approaches developed by Deoni and colleagues to be acquired in a clinically acceptable time³. They also demonstrated that by acquiring DESPOT2 in addition to DESPOT1 it is possible to retrospectively calculate 'synthetic' weighted images with arbitrary T1 or T2 weighting, since the underlying relaxation parameters are known. A study using quantitative MRI (qMRI) in early Parkinson's⁴ found widespread reduction of midbrain T1 values contralateral to the clinically more severely affected limbs and

suggested that T1 may be a valuable marker for the monitoring of progressive neuronal loss in Parkinson's. T1 qMRI probes the chemical composition of the tissues, providing complementary information to the microstructural one available with diffusion imaging. In light of these findings we experimented with this sequence in OPDC-MRI. Quantitative T1 maps were generated using DESPOT-HiFi (dual SPGR, 144 slices) with two IR-SPGR images to assess B1 field inhomogeneity ³. While we were able to use the T1 maps obtained from DESPOT1 ⁵, the quality of DESPOT2 obtained was not deemed sufficient to perform further analyses.

The study by Baudrexel and colleagues ⁴ also acquired multi-echo T2* and performed T2* mapping of the midbrain, which revealed a bilateral decrease of T2* values restricted to the SN, indicating a local increase in total iron content. They conclude that nigral T2* reductions might be associated with an increased general vulnerability for the development of the disorder. We therefore included this small FOV, multi ECHO T2* sequence to explore its diagnostic usefulness given that SN iron accumulation seems implicated in Parkinson's and T2* correlates with iron content.

This sequence was subsequently replaced with whole brain SWI because, despite not being able to provide T2* mapping (non multi-echo), it has a much shorter acquisition time and covers the whole brain, allowing, for example, to quantify iron content in the basal ganglia (see main text for details).

Table S1. Discontinued MRI experimental sequences: parameters used in the study and number of available datasets for each modality.

	SN dMRI	qT1	qT2	SN GRE
Sequence type	EPI	3D, DESPOT-HiFi	3D, DESPOT-HiFi	2D, T2*-weighted
Period of acquisition	2010-2013	2010-2014	2010-2014	2010-2016
TR (ms)	3000	8	55	648
TE (ms)	82	3.7	2.8	5 to 60 [#]
TI (ms)	--	--	--	--
Flip angle (degrees)	90	4, 18	15, 60	45
Voxel size (mm)	0.8x0.8x4	1.1x1.1x1.1	1.1x1.1x1.1	1 x 1 x 2
FoV read (mm)	60	270	270	256
Fov phase (%)	41.7	83.3	83.3	100
Base Resolution	72	240	240	256

Phase Resolution (%)	100	100	100	100
Bandwidth (Hz/Px)	938	210	605	260
Orientation	Transversal	Sagittal	Sagittal	Transversal
N volumes	30 directions + 1 b0 x 10 averages (?)	--	--	--
Other sequence-specific characteristics	b-value = 600 s/mm; reduced FOV, 15 slices	dual SPGR, two IR-SPGR images [§]	RF Phase Increment = 0, 180	#12 TEs, 5 ms increment, reduced FOV, 10 slices
Acquisition time	(10 x) 1 m 36 s	12 m 56 s	(4 x) 1m 42 s	13 m 51 s
Number of subjects				
N iPD	24	46	46	72
N PD-LRRK2	1	5	3	2
N PD-GBA	3	8	7	7
N RBD	0	15	15	20
N RBD-GBA	0	1	1	1
N aGBA	0	8	8	8
N HC	19	26	26	25
N total	47	109	106	135

Legend: iPD = idiopathic Parkinson's patients; PD-LRRK2 = Parkinson's patients with mutation of the LRRK2 gene; PD-GBA = Parkinson's patients with mutation of the GBA gene; RBD = patients with REM sleep behavior disorder (RBD); RBD-GBA = RBD patients with pathogenic mutation of the GBA gene; aLRRK2 = asymptomatic carriers of a pathogenic mutation of the LRRK2 gene; aGBA = asymptomatic carriers of a pathogenic mutation of the GBA gene; HC = Healthy controls. [§] Two IR-SPGR images to assess B1 field inhomogeneity (72 slices, 2.3 x 1.1 x 2.2 mm³, TE 3.7 ms, TR 8 ms, TI 350 ms and 450 ms).

References

- OPDC MRI analysis [<https://ora.ox.ac.uk/objects/uuid:8200af66-f438-4a7b-ad14-e8b032f0a9e7>].
- Vaillancourt DE, Spraker MB, Prodoehl J, et al. High-resolution diffusion tensor imaging in the substantia nigra of de novo Parkinson disease. *Neurology* 2009;72(16):1378-84. doi: 10.1212/01.wnl.0000340982.01727.6e
- Deoni SC, Peters TM, Rutt BK. High-resolution T1 and T2 mapping of the brain in a clinically acceptable time with DESPOT1 and DESPOT2. *Magnetic resonance in medicine : official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* 2005;53(1):237-41. doi: 10.1002/mrm.20314

4. Baudrexel S, Nurnberger L, Rub U, et al. Quantitative mapping of T1 and T2* discloses nigral and brainstem pathology in early Parkinson's disease. *NeuroImage* 2010;51(2):512-20. doi: 10.1016/j.neuroimage.2010.03.005
5. Klein JC, Rolinski M, Griffanti L, et al. Cortical structural involvement and cognitive dysfunction in early Parkinson's disease. *NMR in biomedicine* 2018;31(4):e3900. doi: 10.1002/nbm.3900