

Supplemental Material

Supplemental Tables

Supplemental Table S1 - Fit indexes of the unidimensional models for irritability, outbursts and impairment Confirmatory Factor Analysis Models

Supplemental Table S2 – Factor Loadings and Category Threshold parameters for items in the Irritable Mood section

Supplemental Table S3 - Factor Loadings and Category Threshold parameters for items in the Temper Outbursts section

Supplemental Table S4 - Factor Loadings and Category Threshold parameters for items in the Impairment section

Supplemental Table S5 – Latent Class Model Fit Indexes and Entropy for each four latent class solutions for irritable mood, outbursts and impairment items

Supplemental Table S6 - Fit indexes of the unidimensional, correlated and bifactor models of irritable mood and temper outbursts

Supplemental Table S7 - Irritable mood, outburst and impairment items and their categories of response as described in DMDD section of DAWBA questionnaire

Supplemental Table S8 - Left Censored regression comparing DMDD groups in SDQ scores

Supplemental Table S9 - Psychiatric comorbidities in Disruptive Mood Dysregulation Disorder defined using different combination rules for the irritable mood and temper outburst domains

Supplemental Figures

Supplemental Figure S1 – Latent Class Clinical Indicator endorsement distribution between classes with low and high levels of irritable mood, outbursts and impairment

Supplemental Figure S2 – Receiver Operating Curve for predicting latent classes of subjects with high levels of irritable mood, outbursts and impairment

Supplemental Figure S3 – Comparison between Strengths and Difficulties Questionnaire Scores between Typically Developing Children, Children with other disorders, with irritable mood, outbursts and combined Disruptive Mood Dysregulation Disorder.

Supplemental Figure S4 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder and Matched controls for psychiatric comorbidity using the OR rule.

Supplemental Figure S5 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder and Matched controls for psychiatric comorbidity using the AND rule.

Supplemental Text

Fit Indexes for Confirmatory Factor Analysis

For all CFA unidimensional models, we used delta parameterization and weighted least-square parameters using a diagonal weight matrix with standard errors and with mean and variance-adjusted chi-square test statistics (WLSMV) estimators. The measures of goodness of fit were assessed through the following fit indices: chi-square, WRMR (weighted root mean square residual), CFI (comparative fit index), TLI (Tucker-Lewis Index) and RMSEA, (root mean square error of approximation). To demonstrate good fit to the data, an estimated model should have a WRMR near or below 0.9³¹, a RMSEA of near or below 0.06 and CFI and TLI near or above 0.95³². Details about the estimators and the model fit used in this study can be found in Supplemental Table S1, which provides the fit indexes of the unidimensional models for irritability, outbursts and impairment items.

Supplemental Table S1 - Fit indices of the unidimensional irritable mood, outbursts and impairment Confirmatory Factor Analysis Models

Fit indexes	Irritability	Outbursts	Impairment
FP	36	75	16
X2	3254.236 (df=28; p<0.001)	3752.504 (df=91; p<0.001)	2865.135 (df=6; p<0.001)
RMSEA	0.057	0.057	0.032
CI 90%	0.040- 0.074	0.047-0.068	0.000-0.088
CFI	0.989	0.968	1.000
TLI	0.984	0.961	0.999
WRMR	1.095	0.085	0.019

Note: FP: Free Parameters; X2: Robust Chi Square Difference Test; df: degrees of freedom; RMSEA: Root Mean Square Error of Approximation; CI: Confidence Interval; WRMR: Weighted Root Mean Square Residual.

Supplemental Table S2 – Factor Loadings and Category Threshold parameters for items in the Irritable Mood section

Items (item code in DAWBA)	Factor Loadings	T1	T2	T3	T4	Location (mean)	Location rank
<i>Frequency/duration</i>							
Frequency of irritable/angry mood		Occasionally	1-2/week	\geq 3/week	Every day		
	0.501			0.078	1.143	0.611	4
Long duration of irritability		<1hr	Few hours	Most/all of the day			
	0.444	0.053	0.98	1.726		0.920	6
<i>Characteristics</i>							
Easily irritated	0.901	-0.138	0.775			0.319	3
Intense irritability	0.904	0.281	0.973			0.627	5
<i>Settings</i>							
At home	0.626	-2.123	0.303			-0.910	1
In the classroom	0.292	0.471	1.521			0.996	7

With peers	0.423	0.438	1.608		1.023	8
			A medium			
Irritability evident to others		A little	amount	A great deal		
	0.427	-1.345	-0.095	1.256	-0.061	2

Note: T1= First threshold; T2 = Second threshold; T3 = Third threshold; T4 = Fourth threshold.

Supplemental Table S3 - Factor Loadings and Category Threshold parameters for items in the Temper Outbursts section

Items (item code in DAWBA)	Factor Loadings	T1	T2	T3	T4	Location (mean)	Location rank
<i>Frequency/duration</i>							
		Occasiona	1-2/week	≥ 3/week	Every day		
Frequency of outbursts	0.565	lly		-0.068	1.129	-0.424	5
<i>Characteristics</i>							
		A little	A lot				
Slamming doors	0.525	-0.893	0.333			-0.274	6
Shouting	0.555	-1.004	0.062			-0.472	4
Swearing	0.693	-1.129	-0.015			-0.572	3
Saying mean things to others	0.743	-0.548	0.259			-0.143	7
Saying negative things about himself	0.481	0.633	1.316			0.981	10
Physical aggression to others	0.619	0.050	1.086			0.576	9

Deliberate self-harm	0.527	1.211	2.030	1.626	14
Breaking things	0.572	0.562	1.471	1.023	11
<i>Settings</i>		A little	A lot		
At home	0.722	-2.597	0.068	-1.264	1
In the classroom	0.351	0.604	1.628	1.123	13
With peers	0.372	0.448	1.585	1.024	12
<i>Triggers</i>					
Recognizable triggers		Perhaps	Definitely		
	-0.193	-0.815	-0.403	-0.610	2
Easily triggered		A little	A lot		
	0.664	-0.109	0.784	0.347	9

Note: T1 = First threshold; T2= Second threshold; T3= Third threshold; T4 = Fourth threshold.

Supplemental Table S4 - Factor Loadings and Category Threshold parameters for items in the Impairment section

Items	Factor Loadings	T1	T2	T3	Location (mean)	Location rank
		A little	A medium amount	A great deal		
Impact on family life	0.709	0.018	1.036	2.109	1.054	1
Impact on friendship	0.903	0.545	1.434	2.376	1.452	3
Impact on learning	0.825	0.601	1.238	2.443	1.427	2
Impact on leisure	0.826	0.672	1.521	2.757	1.650	4

Note: T1 = First threshold; T2 = Second threshold; T3 = Third threshold.

Table S5 – Latent Class Model Fit Indexes and Entropy for each of the four latent class solutions for irritable mood, outbursts and impairment items

Irritable Mood							
Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-1.296.125	120	2.634.154	2.611.943	2.641.154	32.466.270	-
Model 2	-1.191.353	112	2.472.502	2.424.906	2.487.502	11.511.877	0.886
Model 3	-1.173.152	104	2.483.992	2.411.012	2.506.992	7.871.695	0.98
Model 4	-1.167.479	96	2.520.538	2.422.173	2.551.538	6.737.120	0.948
Outbursts							
Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-1.501.228	247	3.050.347	3.024.963	3.058.347	3.351.590	-
Model 2	-1.435.849	238	2.973.468	2.919.527	2.990.468	2.044.022	0.665
Model 3	-1.424.724	229	3.005.096	2.922.597	3.031.096	1.821.518	0.634
Model 4	-1.412.391	220	3.034.308	2.923.251	3.069.308	1.574.852	0.719
Impairment							
Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-7.231.254	11	1.470.197	1.457.504	1.474.197	1,98E+08	-

Model 2	-6.302.546	6	1.314.387	1.285.830	1.323.387	1,22E+07	0.736
Model 3	-6.260.462	1	1.335.903	1.291.480	1.349.903	3,78E+06	0.646
Model 4	-6.241.585	-4	1.362.060	1.301.772	1.381.060	1,62E-03	0.547

Note: BIC, Bayesian Information Criterion; aBIC, adjusted Bayesian Information Criteria; cAIC,

Consistent Akaike Information Criterion;

Supplemental Table S6 - Fit indices of the unidimensional, correlated and bifactor models of irritable mood and temper outbursts

Fit indexes	Unidimensional	Correlated with two dimensions
FP	31	32
X2	241.4	231.625
RMSEA	0.066	0.064
CI 90%	0.056-0.076	0.054-0.074
CFI	0.926	0.930
TLI	0.847	0.917
SRMR	0.136	0.134

Note: FP: Free Parameters; X2: Robust Chi Square Difference Test; df: degrees of freedom; RMSEA: Root Mean Square Error of Approximation; CI: Confidence Interval; WRMR: Standardized Root Mean Square Residual.

Supplemental Table S7 - Irritable mood, outburst and impairment items and their response categories as described in DMDD section of the DAWBA questionnaire

DAWBA Questionnaire Items	Response categories				
Irritable Mood Section					
Frequency of irritable/angry mood	Never	Occasionally	1-2/week	≥ 3/week	Every day
Easily irritated	No	A little	A lot		
Intense irritability	No	A little	A lot		
Long duration of irritability *	No more than a few minutes	Less than an hour	A few hours	A medium amount	
Irritability evident to others	Not at all	A little	A great deal		
At home	No	A little	A lot		
In the classroom	No	A little	A lot		
With peers	No	A little	A lot		
Angry weeks (irritable most of the day, nearly every day)*	No	Yes			

Outbursts Section

Frequency of outbursts	Never	Occasionally	1-2/week	≥ 3/week	Every day
Slamming doors	No	A little	A lot		
Shouting	No	A little	A lot		
Swearing	No	A little	A lot		
Saying mean things to others	No	A little	A lot		
Saying negative things about himself	No	A little	A lot		
Physical aggression to others	No	A little	A lot		
Deliberate self-harm	No	A little	A lot		
Breaking things	No	A little	A lot		
At home	No	A little	A lot		
In the classroom	No	A little	A lot		
With peers	No	A little	A lot		
Recognizable triggers	No	Perhaps	Definitely		
Easily triggered	No	A little	A lot		
	Less than a	Less than a	Less than a		More than 3
Outburst-free gap in the last year*	day	week	month	1-3 months	months

Impairment Section

			A medium	
Impact on family life	Not at all	A little	amount	A great deal
			A medium	
Impact on friendship	Not at all	A little	amount	A great deal
			A medium	
Impact on learning	Not at all	A little	amount	A great deal
			A medium	
Impact on leisure	Not at all	A little	amount	A great deal

Note: *The items “long duration of irritability” and “irritable most of the day, nearly every day” have the same category of response, so we included only the first one in the analysis.

**Time interval between outbursts, a criterion included in the current DMDD diagnosis, was not included in the analysis because it was meant to be exclusionary with Bipolar Disorder.

Supplemental Table S8 - Left Censored regression comparing DMDD groups in SDQ

scores

	Contrast with TDC			Contrast with Other DSM-5 Disorders		
	MD	SE	p-value	MD	SE	p-value
SDQ: Total						
Other disorders	9.25	0.35	<0.001	-	-	-
Irritable mood only	11.0	0.70	<0.001	1.75	0.88	0.047
Outbursts only	13.16	1.10	<0.001	3.91	1.31	0.003
Combined	14.43	0.78	<0.001	5.19	0.96	<0.001
SDQ: Emotional						
Other disorders	2.04	0.17	<0.001	-	-	-
Irritable mood only	2.78	0.33	<0.001	-0.05	0.41	0.887
Outbursts only	2.68	0.52	<0.001	-0.09	0.60	0.885
Combined	3.03	0.37	<0.001	0.25	0.44	0.571
SDQ: Hyperkinetic						
Other disorders	3.65	0.22	<0.001	-	-	-
Irritable mood only	3.90	0.43	<0.001	0.23	0.44	0.590
Outbursts only	5.21	0.67	<0.001	1.54	0.65	0.018

Combined	5.03	0.47	<0.001	1.37	0.48	0.004
----------	------	------	--------	------	------	-------

SDQ: Conduct

Other disorders	2.65	0.166	<0.001	-	-	-
-----------------	------	-------	--------	---	---	---

Irritable mood only	3.97	0.324	<0.001	1.32	0.36	<0.001
---------------------	------	-------	--------	------	------	--------

Outbursts only	5.15	0.509	<0.001	2.50	0.53	<0.001
----------------	------	-------	--------	------	------	--------

Combined	5.64	0.358	<0.001	2.99	0.39	<0.001
----------	------	-------	--------	------	------	--------

SDQ: Peer relationships

Other disorders	2.40	0.17	<0.001	-	-	-
-----------------	------	------	--------	---	---	---

Irritable mood only	3.01	0.34	<0.001	0.62	0.39	0.109
---------------------	------	------	--------	------	------	-------

Outbursts only	2.72	0.53	<0.001	0.33	0.58	0.569
----------------	------	------	--------	------	------	-------

Combined	3.37	0.38	<0.001	0.97	0.42	0.02
----------	------	------	--------	------	------	------

SDQ: Impact

Other disorders	4.10	0.22	<0.001	-	-	-
-----------------	------	------	--------	---	---	---

Irritable mood only	4.74	0.39	<0.001	0.61	0.38	0.105
---------------------	------	------	--------	------	------	-------

Outbursts only	4.40	0.60	<0.001	0.30	0.57	0.599
----------------	------	------	--------	------	------	-------

Combined	5.81	0.42	<0.001	1.67	0.40	<0.001
----------	------	------	--------	------	------	--------

Note: TDC, Typically Developing Comparisons; SDQ, Strengths and Difficulties

Questionnaire; MD, Mean Difference.

Supplemental Table S9 – Psychiatric comorbidities in Disruptive Mood Dysregulation Disorder defined using different combination rules for the irritable mood and temper outburst domains


Psychiatric Comorbidities	Current DMDD (n=86)	Irritable Mood (n= 65)	Outbursts (n=26)	AND Rule (n=53)	OR Rule (n=144)
	Prevalence Estimation (%)				
<i>Any disorder (except DMDD)</i>	62.8	56.9	53.8	77.4	63.9
<i>Any Anxiety Disorder</i>	23.3	18.5	11.5	26.4	20.1
Generalized Anxiety Disorder	5.8	4.6	0	7.5	4.9
Separation Anxiety Disorder	7	3.1	3.8	11.3	6.2
Agoraphobia	3.5	1.5	0	3.8	2.1
Panic Disorder	1.2	0	0	1.9	0.7
Social Anxiety Disorder	3.5	6.2	0	3.8	4.2
Specific Phobia	11.6	4.6	7.7	13.2	8.3
Other Specified Anxiety Disorder	1.2	0	0	0	0
<i>Any Mood Disorder</i>	11.6	13.8	3.8	13.2	11.8
Major Depressive Disorder	8.1	9.2	3.8	9.4	8.3
Bipolar Disorder/Mania	3.5	6.2	0	1.9	3.5
Other Specified Depressive Disorder	1.2	1.5	0	1.9	1.4

<i>Any Attention-Deficit/Hyperactivity Disorder</i>	44.2	23.1	34.6	43.4	32.6
Combined	29.1	12.3	19.2	28.3	19.4
Hyperactive/impulsive	3.5	0	11.5	3.8	3.5
Inattentive	9.3	7.7	0	9.4	6.9
Another Specified ADHD	2.3	3.1	3.8	1.9	2.8
<i>Any Disruptive Behavior Disorder</i>	20.9	21.5	26.9	43.4	30.6
Oppositional Defiant Disorder	0	16.9	19.2	17.0	17.4
Conduct Disorder	18.6	10.8	11.5	24.5	16
Other Disruptive Behavior Disorder	2.3	0	3.8	5.7	2.8
<i>Other Disorders</i>					
Obsessive-Compulsive Disorder	1.2	1.5	0	1.9	1.4
Posttraumatic Stress Disorder	3.5	0	0	3.8	1.4
Autism Spectrum Disorder	3.5	0	3.8	5.7	2.8
Eating Disorders	1.2	0	0	1.9	0.7
<i>Current DMDD</i>	-	33.8	42.3	71.7	49.3


Note: DMDD, Disruptive Mood Dysregulation Disorder

Supplemental Figure S1 – Latent Class Clinical Indicator endorsement distribution between classes with low and high levels of irritable mood, outbursts and impairment


Irritable Mood



Outbursts




Impairment




Note: Graphs represents the chance of endorsement of each irritability, outburst and impairment clinical indicators between the two latent classes.


Supplemental Figure S2 – Receiver Operating Curve for predicting latent classes of subjects with high levels of irritable mood, outbursts and impairment



Supplemental Figure S3 – Comparison of Strengths and Difficulties Questionnaire Scores among Typically Developing Children, Children with non-DMDD disorders, with irritable mood only, outbursts only and both irritable mood and outbursts (combined).



Supplemental Figure S4 – Comparison of Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder, defined using the OR rule, and subjects matched for psychiatric comorbidity.



Supplemental Figure S5 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder, defined using the AND rule, and subjects matched for psychiatric comorbidity.

