

## S8 Appendix - Statistical analysis

Analysis of covariance (ANCOVA) was performed to verify if the variation in the average of viable larvae collected during the NVC male mosquitoes release period was statistically significant. At the end of the NVC releases, a significant reduction in the number of viable larvae in the treated area was observed, according to the results of the analysis performed in program R (p value =  $1.37e-11$ , Figure S8A).

ANCOVA analysis was also used to compare the dengue cases originating in the treated and control area during the period of NVC releases. As shown in Figure S8B, the number of dengue cases originating in the treated area was significantly lower than the control area (p value equal to 0.0139).

Linear regression was carried out to evaluate the correlation between dengue cases and larvae mean in both study areas (control and treated). Analysis was performed using the sum of weekly mean viable larvae and the accumulated dengue cases that occurred in the control and treated areas from March 6 to April 22 (2019), which is the period of NVC releases with reported dengue cases. As shown in Figure S8C-a and S8C-b, there is a strong dependence between dengue cases and the number of viable larvae, both in control and treated areas ( $p=0.000124$  and  $p=4.41e-5$ , respectively). Figure S8C-c shows the graph resulting from the linear regression analysis, evidencing the clear difference between the trends of both groups.

The multivariate test (MANOVA) was used to verify the correlation between NVC male mosquito releases, reduction of larvae numbers and the lowest incidence of dengue cases in the treated area. First, a normality test of the data was carried out. As shown in Figure S8D-a, analysis of mean of viable larvae provided a p value = 0.06773, demonstrating that these data follow a normal distribution, whereas data referring to the cases of dengue did not present a normal distribution (p value = 0.01787). Then, a new normality test was performed after control and treated areas were split in three neighborhoods. This new analysis showed a non-significant p value for both dengue cases ( $p=0.4633$ ) and mean of viable larvae ( $p=0.3942$ ), indicating a normal distribution of both sets of data (Figure S8D-b), which enabled MANOVA analysis. The MANOVA test was performed considering both the average of viable larvae collected by trap and dengue cases originating in control and treated study areas. As shown in Figure S8D-c, the test demonstrated a significant influence of treatment (NVC male mosquito releases) on the reduction of the average of viable larvae and on the number of dengue cases (p value = 0.002588).

```
R R Console
> data<-read.table("ancova.txt", header=TRUE)
> week<-data[,1]
> area<-data[,2]
> larvae_mean<-data[,3]
> dengue_cases<-data[,4]
> ancova_larvae<-aov(larvae_mean~week+area,data=data)
> summary(ancova_larvae)
              Df Sum Sq Mean Sq F value    Pr(>F)
week           1   4324    4324    21.48 2.24e-05 ***
area           1  14526   14526    72.18 1.37e-11 ***
Residuals     55  11069     201
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

**Figure S8A. ANCOVA performed in R version 3.5.2 software for viable progeny changes.** Analysis was carried out to verify the variation of viable progeny means over the weeks, from *Ae. aegypti* eggs collected both in control and treated areas. The significative reduction of viable progeny in treated area was determined based on p value ( $p=1.37e-11$ ).

```


R Console
> data<-read.table("ancova.txt", header=TRUE)
> week<-data[,1]
> area<-data[,2]
> larvae_mean<-data[,3]
> dengue_cases<-data[,4]
> ancova_dengue<-aov(dengue_cases~week+area)
> summary(ancova_dengue)

              Df Sum Sq Mean Sq F value    Pr(>F)
week             1  491.8   491.8  18.039 8.41e-05 ***
area             1  175.9   175.9   6.452  0.0139 *
Residuals       55 1499.3    27.3

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

**Figure S8B. ANCOVA performed in R version 3.5.2 software for Dengue cases.** Analysis was carried out to verify the difference on the number of Dengue cases originated in control and treated areas. The number of Dengue cases was significantly lower in treated area compared with control area (p value=0.0139).



**Figure S8C. Linear regression.** Analysis was performed to verify the dependence between mean of eggs collected or mean of viable progeny and number of Dengue cases. Results of linear regression between mean of eggs collected and dengue cases in control area (A) and treated area (B) are shown. Results of linear regression between mean of viable larvae and dengue cases in control area (C) and treated area (D) are shown. E. Graph resulting from the analysis of sum of weekly eggs collected mean and the accumulated dengue cases that occurred in control and treated area from October 3<sup>rd</sup> (2018) to May 29<sup>th</sup> (2019), with linear regression lines. F. Graph resulting from the analysis of sum of weekly larvae viable mean and the accumulated dengue cases that occurred in control and treated area from October 3<sup>rd</sup> (2018) to May 29<sup>th</sup> (2019), with linear regression lines.

**A**

```
R Console
> data_manova<-read.table("manova.txt",header=TRUE)
> data_manova
  area larvae_mean dengue_cases
1 control    38.376          57
2 control    44.508          46
3 control    44.616           6
4 treated     8.962           2
5 treated    12.053           1
6 treated    14.868           5
> shapiro.test(data_manova[,2])

      Shapiro-Wilk normality test

data:  data_manova[, 2]
W = 0.80689, p-value = 0.06773

> shapiro.test(data_manova[,3])

      Shapiro-Wilk normality test

data:  data_manova[, 3]
W = 0.7452, p-value = 0.01787
```

**B**

```
R Console
> shapiro.test(data_manova[1:3,3])

      Shapiro-Wilk normality test

data:  data_manova[1:3, 3]
W = 0.90271, p-value = 0.3942

> shapiro.test(data_manova[4:6,3])

      Shapiro-Wilk normality test

data:  data_manova[4:6, 3]
W = 0.92308, p-value = 0.4633
```

**C**

```
R Console
> data_manova<-read.table("manova.txt",header=TRUE)
> data_manova
  area larvae_mean dengue_cases
1 control    38.376          57
2 control    44.508          46
3 control    44.616           6
4 treated     8.962           2
5 treated    12.053           1
6 treated    14.868           5
> mv_area<-data_manova$area
> mv_larvae<-data_manova$larvae_mean
> mv_dengue<-data_manova$dengue_cases
> manv<-manova(cbind(mv_larvae,mv_dengue)~mv_area, data = data_manova)
> summary(manv)

              Df  Pillai approx F num Df den Df  Fr(>F)
mv_area      1 0.98115   78.081     2     3 0.002588 **
Residuals    4

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

**Figure S8D. MANOVA performed in R version 3.5.2 software.** A. Shapiro-Wilk test was applied to data in order to verify if viable larvae data and Dengue cases data from control and treated areas followed a normal distribution. B. Control and treated areas were split in three neighborhood each, then the Shapiro-Wilk test was applied again and provided a non-significant p value, indicating normal distribution of data. C. Multivariate test was used to verify the influence of NVC male mosquito releases in viable progeny and Dengue incidence during the study period.