

# Ideal Cardiovascular Health in Urban Jamaica: Prevalence Estimates and Relationship to Community Property Value, Household Assets and Educational Attainment (Supplement)

---

## **AUTHORS:**

Joette A. McKenzie<sup>1</sup>, Novie O. Younger-Coleman<sup>1</sup>, Marshall K. Tulloch-Reid<sup>1</sup>, Ishtar Govia<sup>1</sup>, Nadia R. Bennett<sup>1</sup>, Shelly R. McFarlane<sup>1</sup>, Renee Walters<sup>1</sup>, Damian K. Francis<sup>2</sup>, Karen Webster-Kerr<sup>3</sup>, Andriene Grant<sup>3</sup>, Tamu Davidson<sup>3</sup>, Rainford J. Wilks<sup>1</sup>, David R. Williams<sup>4</sup>, Trevor S. Ferguson<sup>1</sup>

## **INSTITUTIONS**

<sup>1</sup>Caribbean Institute for Health Research, The University of the West Indies, Mona, Kingston, Jamaica

<sup>2</sup>School of Health and Human Performance, Georgia College and State University, Milledgeville, GA, USA

<sup>3</sup>Ministry of Health and Wellness, Jamaica

<sup>4</sup>Department of Social and Behavioral Sciences, Harvard T. H. Chan School of Public Health, Boston, MA, USA

## **CORRESPONDENCE:**

Dr Trevor Ferguson

Epidemiology Research Unit

Caribbean Institute for Health Research

The University of the West Indies, Mona

Kingston 7, Jamaica

Tel: 927-2471

Email: [trevor.ferguson02@uwimona.edu.jm](mailto:trevor.ferguson02@uwimona.edu.jm)

## Supplementary File

**Table S1: Items Included in Household Assets List**

Included Household Assets	
1.	Gas stove/Electric Stove
2.	Refrigerator or freezer
3.	Microwave oven
4.	Air conditioner
5.	Fan
6.	Telephone (land line or cell)
7.	Radio/cassette player /Stereo equipment/component set
8.	Electronic Gaming Equipment
9.	Video cassette recorder/ DVD
10.	Washing machine
11.	Clothes Dryer
12.	TV sets
13.	Cable TV
14.	Water Heater (Solar or Electric)
15.	Water Tank
16.	Bicycle
17.	Motorbike
18.	Car or another vehicle
19.	Computer/ Tablet
20.	Computer Accessories /Printer/Fax/Scanner
21.	Smart Phone
22.	Internet service

**Table S2: Number and Proportion of Missing values for variables included in the analyses**

Variable	Number of missing	Number of non-missing values	% missing
Height	88	937	8.6
Weight	91	934	8.9
Body Mass Index	108	917	10.5
Systolic Blood Pressure	69	956	6.7
Diastolic Blood Pressure	69	956	6.7
Fasting glucose	261	764	25.5
Fasting cholesterol	296	729	28.9
Healthy diet	3	1,022	0.3
Current non-smoker	324	701	31.6
Physical Activity	64	961	6.2
Education	33	992	3.2
Household Assets	4	1,021	0.4
Median Land Value	56	969	5.5

**Table S3: Summary Characteristics of Urban Participants in Jamaica Health and Lifestyle Survey III by median land value tertiles**

<b>Characteristic</b>	<b>n<sup>1</sup></b>	<b>Lower Tertile n = 368</b>	<b>Middle Tertile n = 368</b>	<b>Upper Tertile n = 233</b>
<i>Mean for Continuous Variables</i>				
		<i>Mean ± SD</i>	<i>Mean ± SD</i>	<i>Mean ± SD</i>
Age (years)	969	45.8 ± 17.5	46.8 ± 17.4	49.2 ± 17.7
Height (cm)*	886	164.8 ± 9.2	166.2 ± 9.5	164.1 ± 9.9
Weight (kg)	886	80.5 ± 29.2	78.2 ± 20.6	78.1 ± 25.1
BMI (kg/m <sup>2</sup> )	869	29.6 ± 11.0	28.5 ± 8.0	29.2 ± 9.7
Systolic Blood Pressure (mmHg)	905	130.8 ± 22.6	130.6 ± 22.2	130.3 ± 21.5
Diastolic Blood Pressure	905	84.3 ± 13.0	83.8 ± 13.1	83.3 ± 12.5
Fasting Glucose	718	5.8 ± 1.8	5.7 ± 1.9	5.9 ± 2.1
Total Cholesterol	684	4.3 ± 1.1	4.5 ± 1.1	4.6 ± 1.0
<i>Proportions for Categorical Variables</i>				
		% (n)	% (n)	% (n)
Cigarette Smoking	636	13.9 (33)	10.4 (25)	7.6 (12)
Education***	942			
<i>Less than High School</i>		21.8 (78)	24.4 (87)	26.4 (60)
<i>High School</i>		60.1 (215)	46.8 (167)	44.1 (100)
<i>More than High School</i>		18.2 (65)	28.9 (103)	29.5 (67)
Household Assets***	965			
<i>Lower Tertile (≤ 9 items)</i>		43.3 (159)	36.3 (133)	41.4 (96)
<i>Middle Tertile (10 – 12 items)</i>		36.0 (132)	30.9 (113)	25.0 (58)
<i>Upper Tertile (13 – 22 items)</i>		20.7 (76)	32.8 (120)	33.6 (78)

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001 for MLV tertile differences

<sup>1</sup>n = number of available values for each variable

IQR = inter-quartile range

**Table S4A: Mean number of Ideal Cardiovascular Health (ICH) Characteristics, Prevalence of Five or More ICH Characteristics and Individual ICH Components among of Urban Participants in Jamaica Health and Lifestyle Survey III by Median Land Value Category**

<b>Characteristic</b>	<b>Lower Tertile of MLV n = 385</b>	<b>Middle Tertile of MLV n = 384</b>	<b>Upper Tertile of MLV n = 256</b>
Mean number of ICH characteristics	3.5 (3.4, 3.7)	3.6 (3.4, 3.7)	3.7 (3.4, 3.9)
5 or more ICH characteristics	20.2 (14.6, 25.9)	21.4 (15.1, 27.8)	28.7 (21.0, 36.3)
All 4 ICH – Health Behaviours	1.2 (0, 2.7) **	3.8 (1.0, 6.6)	8.1 (3.7, 12.4)
All 4 ICH – Health Factors	15.0 (9.2, 20.8)	13.2 (7.7, 18.7)	14.8 (9.5, 20.2)
Normal Blood Pressure (<120/80 mmHg)	31.6 (24.9, 38.3)	27.3 (20.7, 33.9)	27.3 (21.2, 33.5)
Normal BMI (<25 kg/m <sup>2</sup> )	27.0 (20.2, 33.9)	32.5 (24.7, 40.2)	33.7 (25.8, 41.5)
Normal Glucose (<5.6 mmol/l)	54.3 (45.2, 63.5)	52.4 (43.0, 61.8)	55.0 (45.9, 64.1)
Normal Cholesterol (<5.2 mmol/l)	83.5 (75.2, 91.9) *	77.7 (70.7, 84.7)	69.5 (62.1, 76.8)
Physical Activity ≥150 min/week	58.1 (49.2, 66.9)	57.5 (49.6, 65.4)	62.8 (54.4, 71.2)
Healthy Diet (≥3 of four dietary factors)	13.6 (8.8, 18.5) **	21.7 (18.2, 25.2)	26.0 (19.7, 32.4)
Current Non-Smoker	83.4 (77.6, 89.2)	86.4 (79.5, 93.3)	90.8 (85.8, 95.7)

Values reported are given as percentages with 95% confidence intervals in brackets except for mean number of ICH characteristics which is given as number with 95% CI in brackets

MLV = Median Land Value

ICH = ideal cardiovascular health; ICH includes 7 characteristics (current non-smoking, healthy diet, moderate physical activity, and normal body mass index, blood pressure, glucose, and cholesterol)

ICH – Health Behaviours include adequate physical activity, non-smoker, normal BMI and healthy diet; ICH Health Factors include normal blood pressure, normal glucose, normal cholesterol and current-non-smoker).

Healthy diet defined as meeting 3 or 4 of criteria: (1) adequate fruit and vegetable intake, (2) adequate fish intake (3) low salt diet, (4) low sugar sweetened beverage consumption

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001 for difference in proportion compared to the Upper MLV Tertile

**Table S4B: Mean number of Ideal Cardiovascular Health Characteristics, Prevalence of Five or More ICH Characteristics and Individual ICH Components among of Urban Participants in Jamaica Health and Lifestyle Survey III by Tertiles of Household Assets**

Characteristic	Lower Tertile of Household Assets	Middle Tertile of Household Assets	Upper Tertile of Household Assets
	n = 423	n = 317	n = 285
Mean number of ICH characteristics	3.4 (3.2, 3.6)	3.7 (3.5, 3.9)	3.6 (3.4, 3.8)
5 or more ICH characteristics	18.9 (13.7, 24.1)	27.9 (20.0, 35.7)	22.4 (16.2, 28.7)
All 4 ICH – Health Behaviours	2.6 (0.7, 4.5)	6.5 (1.5, 11.5)	3.4 (1.1, 5.8)
All 4 ICH – Health Factors	13.0 (7.8, 18.2)	14.9 (9.2, 20.6)	14.7 (8.3, 21.1)
Normal Blood Pressure (<120/80 mmHg)	26.1 (20.1, 32.1)	29.7 (23.9, 35.5)	30.2 (23.3, 37.1)
Normal BMI (<25 kg/m <sup>2</sup> )	27.4 (20.4, 34.4)	37.9 (29.5, 46.4)	28.4 (21.0, 35.9)
Normal Glucose (<5.6 mmol/l)	54.4 (46.2, 62.6)	52.6 (43.6, 61.5)	53.9 (44.9, 63.0)
Normal Cholesterol (<5.2 mmol/l)	76.6 (69.7, 83.6)	79.0 (71.9, 86.1)	76.6 (68.8, 84.4)
Physical Activity ≥150 min/week	60.2 (53.4, 67.1)	66.0 (58.2, 73.9) *	51.5 (41.3, 61.7)
Healthy Diet (≥3 of four dietary factors)	16.8 (12.4, 21.2) *	16.8 (11.5, 22.1) *	27.3 (23.0, 31.6)
Current Non-Smoker	80.4 (74.2, 86.5) *	86.9 (81.0, 92.9)	92.6 (87.2, 98.1)

Values reported are given as percentages with 95% confidence intervals in brackets except for mean number of ICH characteristics which is given as number with 95% CI in brackets

ICH = ideal cardiovascular health; ICH includes 7 characteristics (current non-smoking, healthy diet, moderate physical activity, and normal body mass index, blood pressure, glucose, and cholesterol)

ICH – Health Behaviours include adequate physical activity, non-smoker, normal BMI and healthy diet; ICH Health Factors include normal blood pressure, normal glucose, normal cholesterol and current-non-smoker). Healthy diet defined as meeting 3 or 4 of criteria: (1) adequate fruit and vegetable intake, (2) adequate fish intake (3) low salt diet, (4) low sugar sweetened beverage consumption

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001 for difference in proportion compared to the Upper Household Assets Tertile

**Table S4C: Mean number of Ideal Cardiovascular Health (ICH) Characteristics, Prevalence of Five or More ICH Characteristics and Individual ICH Components among of Urban Participants in Jamaica Health and Lifestyle Survey III by Educational Attainment Category**

Characteristic	Less than High School	High School	More than High School
<b>Male</b>	<b>n = 111</b>	<b>n = 182</b>	<b>n = 67</b>
Mean number of ICH characteristics	3.2 (2.9, 3.6)	3.8 (3.5, 4.1)	3.4 (3.0, 3.8)
5 or more ICH characteristics	16.8 (7.3, 26.2)	30.9 (20.1, 41.8)	15.5 (4.8, 26.1)
All 4 ICH – Health Behaviours	3.0 (0, 6.3)	7.7 (2.2, 13.1)	6.1 (0, 12.7)
All 4 ICH – Health Factors	1.4 (0, 4.3)	14.9 (5.6, 24.1)	7.5 (0, 15.0)
Normal Blood Pressure (<120/80 mmHg)	8.2 (1.4, 15.1)	26.2 (16.2, 36.3)	17.2 (6.9, 27.5)
Normal BMI (<25 kg/m <sup>2</sup> )	36.9 (25.4, 48.3)	44.9 (33.1, 56.8)	30.1 (15.8, 44.3)
Normal Glucose (<5.6 mmol/l)	44.0 (30.3, 57.6)	56.3 (42.5, 70.0)	52.8 (37.5, 68.0)
Normal Cholesterol (<5.2 mmol/l)	67.0 (52.7, 81.2)	83.3 (74.1, 92.6)	76.4 (61.7, 91.2)
Physical Activity ≥150 min/week	69.4 (59.8, 79.1)	69.5 (63.6, 75.5)	61.7 (45.4, 78.1)
Healthy Diet (≥3 of four dietary factors)	25.6 (14.8, 36.4) *	19.0 (12.6, 25.4)	13.7 (6.2, 21.2)
Current Non-Smoker	71.0 (58.6, 83.3)	78.5 (68.0, 89.0)	88.7 (77.8, 99.6)
<b>Female</b>	<b>n = 143</b>	<b>n = 343</b>	<b>n = 179</b>
Mean number of ICH characteristics	3.2 (2.8, 3.5) **	3.5 (3.4, 3.7) *	3.8 (3.6, 3.9)
5 or more ICH characteristics	12.5 (3.9, 21.1) *	18.8 (13.8, 23.8) *	29.0 (21.2, 36.9)
All 4 ICH – Health Behaviours	1.5 (0, 3.7)	0.6 (0, 1.3) *	4.1 (0.6, 7.6)
All 4 ICH – Health Factors	5.7 (0, 11.5) *	18.1 (11.7, 24.4)	21.4 (13.2, 29.6)
Normal Blood Pressure (<120/80 mmHg)	14.1 (5.9, 22.4) ***	38.3 (31.5, 45.2)	41.5 (34.2, 48.8)
Normal BMI (<25 kg/m <sup>2</sup> )	26.0 (16.7, 35.4)	21.2 (16.0, 26.4)	25.2 (17.5, 33.0)
Normal Glucose (<5.6 mmol/l)	42.4 (29.5, 55.2) *	55.0 (47.3, 62.7)	58.0 (48.4, 67.7)
Normal Cholesterol (<5.2 mmol/l)	63.5 (51.1, 75.9) *	78.8 (73.4, 84.3)	78.5 (70.1, 86.9)
Physical Activity ≥150 min/week	52.6 (39.1, 66.2)	52.3 (44.9, 59.7)	50.2 (42.6, 57.7)
Healthy Diet (≥3 of four dietary factors)	24.5 (15.2, 33.8)	16.5 (11.7, 21.3) *	26.9 (20.8, 33.0)
Current Non-Smoker	93.5 (87.2, 100.0)	90.3 (85.6, 94.9)	95.2 (90.5, 99.9)

Values reported are given as percentages with 95% confidence intervals in brackets except for mean number of ICH characteristics which is given as number with 95% CI in brackets

ICH = ideal cardiovascular health; ICH includes 7 characteristics (current non-smoking, healthy diet, moderate physical activity, and normal body mass index, blood pressure, glucose, and cholesterol)

ICH – Health Behaviours include adequate physical activity, non-smoker, normal BMI and healthy diet; ICH Health Factors include normal blood pressure, normal glucose, normal cholesterol and current-non-smoker). Healthy diet defined as meeting 3 or 4 of criteria: (1) adequate fruit and vegetable intake, (2) adequate fish intake (3) low salt diet, (4) low sugar sweetened beverage consumption

\*p<0.05; \*\*p<0.01; \*\*\*p<0.001 for difference in proportion compared to the 'More than High School' Education

**Table S5: Odds Ratio for  $\geq 5$  Ideal Cardiovascular Health Characteristics among of Urban Participants in Jamaica Health and Lifestyle Survey using multilevel models**

Characteristic	Males		Females	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Median Land Value				
Upper Tertile (>2 – 9 million JMD)	1.0		1.0	
Middle Tertile (1-2 million JMD)	0.61 (0.29 – 1.28)	0.188	0.72 (0.41 – 1.24)	0.240
Lower Tertile (<1 million JMD)	0.47 (0.21 – 1.05)	0.067	0.59 (0.33 – 1.05)	0.071
Household Assets				
Upper Tertile (13-22 items)	1.0		1.0	
Middle Tertile (10-12 items)	1.65 (0.77 – 3.53)	0.197	1.10 (0.59 – 2.02)	0.769
Lower Tertile ( $\leq 9$ items)	1.08 (0.48 – 2.45)	0.856	1.08 (0.59 – 1.97)	0.802
Education Category				
More than High School	1.0		1.0	
High School	2.16 (0.94 – 4.96)	0.069	0.60 (0.36 – 1.00)	0.050
Less than High School	2.01 (0.67 – 6.03)	0.214	0.57 (0.25 – 1.27)	0.171
Age (years)	0.97 (0.95 – 0.99)	0.002	0.97 (0.95 – 0.98)	<0.001

Separate models created for males and female participants in two level multi-level model with individuals nested in communities and using multiple imputation to account for missing data. The model for males included 349 participants while the model for females included 637 participants.



**Table S6: Regression coefficients for Number of Ideal Cardiovascular Health Characteristics among of Urban Participants in Jamaica Health and Lifestyle Survey (ANCOVA models)**

Characteristic	Males		Females	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Median Land Value				
Upper Tertile (>2 – 9 million JMD)	Referent		Referent	
Middle Tertile (1-2 million JMD)	-0.36 (-0.82, 0.09)	0.116	0.01 (-0.27, 0.28)	0.968
Lower Tertile (<1 million JMD)	-0.57 (-0.99, -0.15)	0.008	0.17 (-0.26, 0.30)	0.901
Household Assets				
Upper Tertile (13-22 items)	Referent		Referent	
Middle Tertile (10-12 items)	0.10 (-0.50, 0.70)	0.737	0.05 (-0.22, 0.32)	0.728
Lower Tertile ( $\leq 9$ items)	-0.07 (-0.64, 0.52)	0.821	-0.08 (-0.31, 0.15)	0.504
Education Category				
More than High School	Referent		Referent	
High School	0.46 (-0.07, 0.98)	0.087	-0.27 (-0.49, -0.05)	0.016
Less than High School	0.32 (-0.34, 0.98)	0.333	-0.34 (-0.74, 0.07)	0.101
Age (years)	-0.02 (-0.03, -0.01)	0.001	-0.02 (-0.03, -0.01)	0.001

Separate models created for males and female participants in ANCOVA model weighted for survey design and using multiple imputation to account for missing data. The model for males included 360 participants while the model for females included 665 participants.

**Table S7: Odds Ratio for ≥5 Ideal Cardiovascular Health Characteristics among of Urban Participants in Jamaica Health and Lifestyle Survey (multivariable models using complete cases only)**

Characteristic	Males		Females	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Median Land Value				
Upper Tertile (>2 – 9 million JMD)	1.0		1.0	
Middle Tertile (1-2 million JMD)	0.40 (0.16 – 1.01)	0.053	0.61 (0.19 – 1.94)	0.403
Lower Tertile (<1 million JMD)	0.37 (0.12 – 1.11)	0.078	0.91 (0.33 – 2.51)	0.859
Household Assets				
Upper Tertile (13-22 items)	1.0		1.0	
Middle Tertile (10-12 items)	2.01 (0.93 – 4.72)	0.073	1.29 (0.68 – 2.40)	0.427
Lower Tertile (≤ 9 items)	1.34 (0.451 – 4.00)	0.598	1.12 (0.61 – 2.06)	0.720
Education Category				
More than High School	1.0		1.0	
High School	3.19 (0.89 – 11.44)	0.032	0.79 (0.41 – 0.1.50)	0.463
Less than High School	4.87 (1.15 – 20.64)	0.075	1.12 (0.46 – 2.72)	0.806
Age (years)	0.95 (0.92 – 0.99)	0.018	0.96 (0.93 – 0.99)	0.007

Separate models created for males and female participants in logistic regression models weighted for survey design with listwise deletion of missing data. The model for males included 336 participants while the model for females included 602 participants.

**Table S8: Prevalence of Ideal Cardiovascular Health if Various Population**

First Author (year published)	Population	Sample Size	Age group	Prevalence of all 7 characteristics	Prevalence of all 6- 7 characteristics	Prevalence of all ≥5 characteristics
Harrison (2019)	Quebec	777	18-65 years	0.5	16.2	36.9
Jankovic (2019)	Serbia	13,100	≥20 years	0.1%	4.1	19.9
Zeng (2012)	China (urban)	9962	≥20 – 85 years	0.5	6.4	26.9
Kim (2013)	New Ulm, Minnesota	4754		1.0	7.5	21.9
Short (2013)	Mississippi	2003 Black, 5,125 White	≥18 years	1.3 (Black) 2.6 (White)	-	-
Moghaddam (2014)	Tehran	2,861 women, 2,004 men	≥20 years	One woman No men	-	19.7 (women) 10.3 (men)
Lu (2015)	China (southeast)	11,996	≥20 years	1.1	8.9	27.6
Djousse' (2015)	Mississippi (Jackson)	4132	Mean 55.3 years	0 (men) 0 (women)	0.3 (men) 0.6 (women)	3.9 (men) 4.3 (women)
Gupta (2015)	India	6198	20-75 years	0.1 (men) 0.1 (women)	2.5 (men) 3.6 (women)	15.1 (men) 21.3 (women)
Valasquez (2015)	Brazil	34,362	≥18 years	0.34	-	-
Younis (2016)	USA (review of 14 studies)	-	-	-	0.3 - 15%	-
Empana (2016)	Paris	8916	5-75	-	-	9.98
Ren (2016)	China (Shandong)	7683 men 7667 women	18-69	0.05	2.95	23.3
Benziger (2018)	Peru	3058	≥35	0	-	12.7
Seron (2018)	Argentina, Chile, Uruguay	5458	35-75	0.1	-	8.5%
van Nieuwenhuizen (2018)	Ghana	3510	20-70	0.3	-	25.2 (women) 27.0 (men)