

Multimodal neuroimaging and suicidality in a US population-based sample of school-aged children

Supplemental Material

eMethods

1. ABCD Study
2. Determination of Suicidality
3. Individual and Environmental Characteristics

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics

4. Neuroimaging
 - 4.1. Brain structural imaging data
 - 4.2. Resting-state functional imaging data
 - 4.3. Task-based functional imaging data
 - Monetary incentive delay task (MID)
 - Stop signal task (SST)
 - Emotional n-back (EN-back)

eFigure 1. Sample selection flowchart

- 4.4. Imaging data preparation for analysis
- 4.5. Statistical analysis of neuroimaging data.
 - 4.5.1. Testing for difference and equivalence
 - 4.5.2. Testing for predictive value

eResults

1. Prevalence of suicidality

eTable 2. Suicidal ideation, plans and attempts in unrelated ABCD participants

2. Summary of findings

eTable 3. Number and percentage of cortical regions found for each possible combined outcome of the Welch's t-test and the Equivalence test by imaging modality and suicidality group comparison – FDR-corrected results

eTable 4. Number and percentage of cortical regions found for each possible combined outcome of the Welch's t-test and the Equivalence test by imaging modality and suicidality group comparison – Uncorrected results

3. Brain Structural Imaging

eTable 5. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the child-reported suicidality group: Welch's t-test

eTable 6. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

eTable 7. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

eFigure 2. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the child-reported suicidality group.

eFigure 3. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the child-reported suicidality group.

eFigure 4. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group.

eFigure 5. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the caregiver-reported suicidality group.

eFigure 6. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group.

eFigure 7. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the concordantly-reported suicidality group.

4. Resting-State Functional Imaging

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

eFigure 8. Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group.

eFigure 9. Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group.

eFigure 10. Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly-reported suicidality group.

5. Task-Based Functional Imaging

Monetary Incentive Delay task (MID)

Anticipation of large reward vs neutral

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

eTable 12. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward: Welch's t-test

eTable 13. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for anticipation of reward: Welch's t-test

eFigure 11. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward

eFigure 12. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward

eFigure 13. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward

eFigure 14. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward

eFigure 15. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward

eFigure 16. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward

Anticipation of large loss vs neutral

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

eTable 16. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for anticipation of loss: Welch's t-test

eFigure 17. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss

eFigure 18. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss

eFigure 19. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss

eFigure 20. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss

eFigure 21. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss

eFigure 22. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss

Positive vs negative feedback of reward

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

eTable 19. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for reward feedback: Welch's t-test

eFigure 23. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback

eFigure 24. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback

eFigure 25. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback

eFigure 26. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback

eFigure 27. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback

eFigure 28. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback

Positive vs negative feedback of loss

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

eTable 22. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for loss feedback: Welch's t-test

eFigure 29. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback

eFigure 30. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback

eFigure 31. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback

eFigure 32. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback

eFigure 33. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback

eFigure 34. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback

Stop Signal Task (SST)

Incorrect stop vs correct go

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

eFigure 35. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for failed stops

eFigure 36. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for failed stops

eFigure 37. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops

eFigure 38. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops

eFigure 39. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops

eFigure 40. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops

Correct stop vs correct go

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

eTable 28. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for correct stops: Welch's t-test

eFigure 41. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for correct stops

eFigure 42. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for correct stops

eFigure 43. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops

eFigure 44. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops

eFigure 45. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops

eFigure 46. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops

Emotional n-back

2-back vs 0-back

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

eTable 31. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for 2-back vs 0-back: Welch's t-test

eFigure 47. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back

eFigure 48. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back

eFigure 49. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back

eFigure 50. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back

eFigure 51. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back

eFigure 52. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back

Emotional vs neutral faces

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

eTable 34. Differences in functional activity between the never-suicidal group and the concordantly -reported suicidality group for emotional faces: Welch's t-test

eFigure 53. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces

eFigure 54. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces

eFigure 55. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces

eFigure 56. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces

eFigure 57. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces

eFigure 58. Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces

eFigure 59. Equivalence testing for mean differences in functional activation during the MID task, the SST and EN-back task relating to suicidality

eTable 35. Range of effect sizes (Cohen's *d*) and 90% confidence intervals bounds for each imaging modality and suicidality group comparison

eTable 36. Results with effect sizes (Cohen's *d*) largest than $|0.15|$ for each imaging modality in child- and caregiver-suicidality group comparison

eReferences

eMethods

1. The ABCD study

The ABCD study is a 22-site study that aims to follow 9-10-year-olds annually for up to 10 years. The study sample was designed to match the sociodemographic variation of the US population based on a within-site school-based sampling strategy of local randomization and representativeness ¹. All caregivers provided written informed consent and all children provided assent.

2. Determination of childhood Suicidality

Suicidality in children and caregivers was assessed with the KSADS-5 ². The KSADS-5 assesses the following aspects of suicidality: passive suicidal ideation, active but nonspecific suicidal ideation, suicidal ideation with a specific method, active suicidal ideation with an intent, active suicidal ideation with a plan, preparatory actions toward imminent suicidal behavior, interrupted suicidal attempts, aborted suicidal attempts and suicidal attempts. Children were considered as having experienced suicidality if they or their caregiver endorsed any of the above items as present either in the past or at the time of the assessment. Items relating to self-injurious behavior without suicidal intent were not included. In the ABCD database, child-reported items are in txt file=abcd_ksad501 and caregiver-reported items are in txt file= abcd_ksad01. We defined suicidality as endorsing at least one of the following KSADS-5 items:

- (1) Current passive suicidal ideation: child-report item: ksads_23_946_t; caregiver-report item: ksads_23_946_p; 1=endorsed
- (2) Past passive suicidal ideation: Youth report item: ksads_23_957_t; Caregiver-report item: ksads_23_957_p ; 1=endorsed
- (3) Current active but nonspecific suicidal ideation - Youth report item: ksads_23_947_t; Caregiver-report item: ksads_23_947_p; 1=endorsed
- (4) Past active but nonspecific suicidal ideation; child-report item: ksads_23_958_t; caregiver-report item: ksads_23_958_p; 1=endorsed
- (5) Current active suicidal ideation with a specific method: child-report item: ksads_23_948_t; Caregiver-report item: ksads_23_948_p; 1=endorsed
- (6) Past active suicidal ideation with a specific method: child-report item: ksads_23_959_t; caregiver-report item: ksads_23_959_p 1=endorsed
- (7) Current active suicidal ideation with an intent: child-report item: ksads_23_949_t; caregiver-report item: ksads_23_949_p; 1=endorsed

- (8) Past active suicidal ideation with an intent: child-report item: ksads_23_960_t; caregiver-report item: ksads_23_960_p; 1=endorsed
- (9) Current active suicidal ideation with a plan: child-report item: ksads_23_950_t; caregiver-report item: ksads_23_950_p 1=endorsed
- (10) Past active suicidal ideation with a plan: child-report item: ksads_23_961_t; caregiver-report item: ksads_23_961_p 1=endorsed
- (11) Current preparatory Actions toward imminent suicidal behavior: child-report item: ksads_23_951_t; caregiver-report item: ksads_23_951_p; 1=endorsed
- (12) Past preparatory Actions toward imminent suicidal behavior: child-report item: ksads_23_962_t; caregiver-report item: ksads_23_962_p; 1=endorsed
- (13) Current interrupted suicidal attempt: child-report item: ksads_23_952_t; caregiver-report item: ksads_23_952_p ; 1=endorsed
- (14) Past interrupted suicidal attempt: child-report item: ksads_23_963_t; caregiver-report item: ksads_23_963_p; 1=endorsed
- (15) Current aborted suicidal attempt: child-report item: ksads_23_953_t; caregiver-report item: ksads_23_953_p; 1=endorsed
- (16) Past aborted suicidal attempt: child-report item: ksads_23_964_t; caregiver-report item: ksads_23_964_p; 1=endorsed
- (17) Current suicidal attempt: child-report item: ksads_23_954_t; caregiver-report item: ksads_23_954_p; 1=endorsed
- (18) Past suicidal attempt: child-report item: ksads_23_965_t; caregiver-report item: ksads_23_965_p; 1=endorsed

Items not included:

- (19) Self-injurious behavior without suicidal intent: child-report item: ksads_23_945_t ksads_23_956_t; caregiver-report item: ksads_23_945_p ksads_23_956_p;

3. Individual and Environmental Characteristics

Details of the assessments protocols in the ABCD study have been published previously³⁻⁵.and consist of study specific instruments and as well as widely used instruments.

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics		
Variable	ABCD Study assessment	Description
Individual characteristics		

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics

Variable	ABCD Study assessment	Description
Sex	ABCD Parent Demographics Survey	Responses coded as: 0=Female 1=Male
Race	ABCD Parent Demographics Survey	Participants answered the following question: <i>“What race do you consider the child to be? Please check all that apply”</i> Responses coded as: White (0=No, 1=Yes) Black/African American Asian (includes Asian Indian, Chinese, Filipino, Japanese, Korean, and Vietnamese; 0=No, 1=Yes) Multiple races (if participant checked more than one; 0=No, 1=Yes) Other races (0=No, 1=Yes)
Ethnicity	ABCD Parent Demographics Survey	Participants answered the following question: <i>“Do you consider the child Hispanic/Latino/Latina?”</i> Responses coded as: 0=No 1=Yes
Intelligence	ABCD NIH Toolbox: Cognition Total Composite	Age-Corrected Standard Score
Exposure to stressful events (Caregiver report)	Based on the KSADS-5 Section on Traumatic Events	Responses coded as: 0=Never exposed 1= Exposed
Common childhood medical conditions (Caregiver report)	ABCD Medical History Questionnaire	Participants answered the following question: <i>“Has the child ever been to a doctor for any of these things? Bronchitis, Allergies, Asthma?”</i> Responses coded as: 0=Never had any common childhood medical condition 1= Had at least one common childhood medical condition
Pubertal development (Caregiver report)	ABCD Parent Pubertal Development Scale and Menstrual Cycle Survey History (PDMS)	Total score (range 1-5) by sex
Pubertal development (Child report)	ABCD Youth Pubertal Development Scale and Menstrual Cycle Survey History (PDMS)	Total score (range 1-5) by sex

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics

Variable	ABCD Study assessment	Description
Environmental, family and school characteristics		
Income level	ABCD Parent Demographics Survey	<p>Participants answered the following question: “<i>What is your TOTAL COMBINED FAMILY INCOME for the past 12 months?</i>”</p> <p>Score ranging from 1 (less than \$5,000) to 10 (\$200,000 and greater).</p>
Economic problems in the last year	ABCD Parent Demographics Survey	<p>Participants answered the following question: “<i>In the past 12 months, has there been a time when you and your immediate family experienced any of the following: 1) Needed food but couldn't afford to buy it or couldn't afford to go out to get it. 2) Were without telephone service because you could not afford it. 3) Were evicted from your home for not paying the rent or mortgage. 4) Had services turned off by the gas or electric company, or the oil company wouldn't deliver oil because payments were not made. 5) Had someone who needed to see a doctor or go to the hospital but didn't go because you could not afford it. 6) Had someone who needed a dentist but couldn't go because you could not afford it?</i>”</p> <p>Responses coded as: 0=Did not experience any of the problems above 1= Experienced at least one of the problems above</p>
Primary caregiver level of education	ABCD Parent Demographics Survey	<p>Participants answered the following question: “<i>What is the highest grade or level of school you have completed or the highest degree you have received?</i>”</p> <p>Score ranging from 0 (Never attended or Kindergarten only) to 21 (Doctoral degree).</p>
Primary caregiver marital/partnership status	ABCD Parent Demographics Survey	<p>Participants answered the following question: “<i>Are you now married, widowed, divorced, separated, never married or living with a partner?</i>”</p> <p>Responses coded as: 0= Married/ Living with partner</p>

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics		
Variable	ABCD Study assessment	Description
		1=Widowed/Divorced/Separated/Never married
Family Conflict (Caregiver report)	ABCD Family Environment Scale: Family Conflict Subscale Modified from PhenX	The number of times events indicating family conflict occurred in the past year are summed for all questions; higher scores indicating greater conflict
Family Conflict (Child report)	ABCD Family Environment Scale: Family Conflict Subscale Modified from PhenX	The number of times events indicating family conflict occurred in the past year are summed for all questions; higher scores indicating greater conflict
Positive school environment (Child report)	ABCD School Risk and Protective Factors Survey: School Environment Subscale	Score ranging 1-24; higher scores indicating a more perceived positive school environment
Individual clinical characteristics		
Child general psychopathology (Caregiver report)	Child Behavior Checklist (ASEBA)	T-scores for total problems
Psychiatric disorders (Caregiver report)	ABCD Parent Diagnostic Interview for DSM-5 Full (KSADS-5)	Responses coded as: Any current depressive disorder (includes major depression disorder, dysthymia, and unspecified depressive disorder; 0=No 1=Yes) Any current anxiety disorder (includes generalized anxiety disorder, panic disorder, agoraphobia, separation anxiety disorder, social anxiety disorder, specific phobia, and other specified anxiety disorder, ; 0=No 1=Yes) Current ADHD (includes attention-deficit hyperactivity disorder [ADHD] and unspecified ADHD; 0=No 1=Yes) Current ODD (includes oppositional defiant disorder [ODD]; 0=No 1=Yes) CD (includes conduct disorder; 0=No 1=Yes) Current PTSD (includes post-traumatic stress disorder, and other specified trauma-and stressor-related disorder; 0=No 1=Yes)
Family clinical characteristics		
Caregiver general	Adult Self-Report (ASEBA)	T-scores for total problems

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics

Variable	ABCD Study assessment	Description
psychopathology (Caregiver report)		
Parental use of mental health services (Caregiver report)	ABCD Family History Assessment	Participants answered the following questions: <i>“Has the biological mother/father of your child ever been to a doctor or a counselor about any emotional or mental problems, or problems with alcohol or drugs?”</i> Responses coded as: 0=No 1= Yes
Parental hospitalization due to mental health problems (Caregiver report)	ABCD Family History Assessment	Participants answered the following questions: <i>“Has the biological mother/father of your child ever been hospitalized because of emotional or mental problems, or drug or alcohol problems?”</i> Responses coded as: 0=No 1= Yes
Parental history of depression (Caregiver report)	ABCD Family History Assessment	Participants answered the following questions: <i>“Has the biological mother/father of your child ever suffered from depression, that is, have they felt so low for a period of at least two weeks that they hardly ate or slept or couldn't work or do whatever they usually do?”</i> Responses coded as: 0=No 1= Yes
Parental history of attempted or committed suicide (Caregiver report)	ABCD Family History Assessment	Participants answered the following questions: <i>“Has the biological mother/father of your child ever attempted or committed suicide?”</i> Responses coded as: 0=No 1= Yes
Maternal alcohol/substance use during pregnancy (Caregiver report)	ABCD Developmental History Questionnaire	Participants answered the following question: <i>“Once the biological mother knew she was pregnant, was she using any of the following? Tobacco, Alcohol, Marijuana, Cocaine, Heroin/Morphine, Oxycontin”</i>

eTable 1. Definition of variables assessing individual, environmental (family and school) and clinical characteristics

Variable	ABCD Study assessment	Description
		Responses were coded as 0= No 1= Yes if any use
Further details can also be found at https://nda.nih.gov/data_dictionary.html		

4. Neuroimaging

Participants underwent magnetic resonance imaging (MRI) scanning to obtain 3D T₁- and 3D T₂-weighted images, diffusion weighted images and resting state- and task-functional data using standard adult-size multi-channel coils capable of multiband echo planar imaging (EPI) acquisitions. The data from different MR modalities were acquired in a fixed order as follows: 3D T₁-weighted images, 2 runs of resting state fMRI (rs-fMRI), diffusion weighted images, 3D T₂-weighted images, 1–2 more runs of resting state fMRI and the task-based fMRI, which included a modified monetary incentive delay task (MID)⁶, stop signal task (SST)⁷ and emotional n-back task (EN-back)⁸. The order of the three fMRI tasks was randomized across participants. Scan sessions typically required ~2 hours to complete, including a mid-session rest break if the child requested one. The acquisition protocol was harmonized across three 3T scanner systems (Siemens Prisma, General Electric 750 and Philips). Before the scan, participants completed a screening questionnaire for MR contraindications and were familiarized to the scanner environment with a simulator using motion compliance training to decrease motion artefacts. Details of the acquisition protocols and procedures, as well as a detailed description of the image processing and analyses have been published in Casey et al (2018)⁹ and Hagler et al (2019)¹⁰, respectively. Below we summarize this information and describe the selection criteria of participants for the current study.

Note: On December 2019, an ABCD Study release on Known Issues informed of incorrect post-processing on all task and resting-state fMRI data obtained in Phillips scanners. Specifically, the field map direction for these data was mistakenly flipped, which led to increased distortion in processed fMRI images. For this reason, participants with fMRI data collected in Phillips scanners were excluded from fMRI analyses in the current study.

4.1 Brain structural imaging data

Following pre-processing, cortical reconstruction and volumetric segmentation of structural datasets was conducted using FreeSurfer image analysis suite (version 5.3.0; <http://surfer.nmr.mgh.harvard.edu/>). For each individual, we considered FreeSurfer-derived regional cortical thickness measures (n=68) and regional subcortical volume measures (n=18). Regional volumetric and cortical thickness measures were respectively adjusted for total intracranial volume and mean cortical thickness.

ABCD unrelated participants with complete suicidality data were excluded from the analysis of structural data for the following technical and medical reasons:

(1) Poor image quality according to the Quality Control and Recommended Inclusion Criteria for structural data provided in the ABCD documentation (**n=671**). These are outlined in a pdf document as part of the Release Notes Imaging Instruments folder. According exclusions were based on the rating of the following items:

Txt file= abcd_mrfindings01; scan excluded if mrif_score ≥ 3

Txt file= ABCD MRI QC Raw Part 1; item= iqc_t1_ok_ser; scan excluded if 0

(2) Incomplete or poorly Freesurfer deconstruction (**n=580**)

Txt file= FreeSurfer QC; item= fsqc_qc; scan excluded if 0

(3) Severe medical conditions likely to affect brain structure (**n=505**): We excluded participants diagnosed with traumatic brain injury with loss of consciousness; muscular dystrophy, multiple sclerosis, sickle cell anemia, neoplastic disorders; cerebral palsy, diabetes, epilepsy, lead Poisoning, cardiac and kidney disorders. The relevant items in the ABCD documentation are:

Text file= ABCD Sum Scores Traumatic Brain Injury: item= tbi_ss_ntbiloc; scan excluded if >0

Txt file=ABCD Parent Medical History Questionnaire (MHX): medhx_2l, medhx_2m, medhx_2p, medhx_2e, medhx_2f, medhx_2g, medhx_2h, medhx_2k, medhx_2o, medhx_2j; for all items, scan excluded if 1

4.2 Resting-state functional imaging data

During collection of rs-fMRI data, children viewed a crosshair for 20 minutes. To quantify connectivity strength during rs-fMRI, a seed-based correlational approach was applied on parcellated regions¹¹⁻¹³. Using a functional atlas¹¹ we grouped the cortical-surface regions into 12 predefined large-scaled networks: auditory, cingulo-opercular, cingulo-parietal, default-mode, dorsal-attention, fronto-parietal, retrosplenial-temporal, salience, sensorimotor-hand, sensorimotor-mouth, ventral-

attention, and visual networks. We computed functional connectivity estimates using Fisher r to z transformations of the mean correlations between pairs of regions examined. Specifically, we computed the within-network functional connectivity ($n=12$), the between-network functional connectivity ($n=66$) and the functional connectivity between each network and 19 subcortical regions ($n=228$).

ABCD unrelated participants with complete suicidality data were excluded from the rs-fMRI analyses for the following technical and medical reasons:

- (1) Incidental findings (e.g. hydrocephalus, herniation), unsatisfactory quality control (QC) for rs-fMRI for at least 4 runs (ABCD QC criteria recommendation requires at least 1 only), unsatisfactory T1, had less than 375 data points worth of 1 run, incomplete or poorly Freesurfer deconstruction, or missing information in these indicators (**$n=2282$**)

Txt file=MR Findings; item = abcd_mrfindings01; scan excluded if $mrif_score \geq 3$, $mrif_hydrocephalus="yes"$, $mrif_herniation="yes"$.

Txt file= ABCD MRI QC Raw Part 1: item= iqc_rsfmri_ok_ser; scan excluded if <4

Txt file= ABCD MRI QC Raw Part 1; item= iqc_t1_ok_ser; scan excluded if 0

Txt file= ABCD rsfMRI Gordon Network Correlations: included if $rsfmri_c_ngd_ntpoints > 375$

Txt file= FreeSurfer QC; item= fsqc_qc; scan excluded if 0

- (2) Missing ROI-based tabulated data (**$n=1029$**)
- (3) Phillips scan data (**$n=527$**)

4.3 Task-based functional imaging data

We examined activations evoked by three tasks: a modified monetary incentive delay task (MID) ⁶, stop signal task (SST) ⁷ and emotional n-back task (EN-back) ^{8,14}, selected to probe reward processing, inhibitory control, and working memory, respectively ⁹, with the latter also including affective processing. The processes underlying these tasks have been implicated in suicidal behavior in clinical samples of older youth and adults ¹⁵⁻²⁵. For each task, task-based changes in the blood-oxygen-dependent (BOLD) signal were computed at the individual subject level using a general linear model (GLM) implemented in AFNI's 3dDeconvolve ²⁶. Average time courses were calculated for 167 regions-of-interest (ROI) derived from the Freesurfer cortical parcellation ($n=148$) ²⁷ and subcortical segmentation ($n=19$) ²⁸.

The MID task started with a cue, indicating possible earnings: large reward (\$5), small reward (\$0.20), no money (\$0), small loss (\$0.20), and large loss (\$5). After a variable period following the cue, the children were shown a brief target. To either earn reward or avoid loss, children needed to respond before the target disappeared. Following the response, children saw the outcome feedback of the trial. For the MID we considered measures derived from the following contrasts: *anticipation of large reward vs neutral condition* (i.e, no gain or loss), *anticipation of large loss vs neutral condition*, *reward positive feedback vs reward negative feedback* and *loss positive feedback vs loss negative feedback*. For reward anticipation, we focused on large reward, since it has been shown to maximize this process ²⁹.

In the SST, each trial begins with the presentation of a leftward or rightward pointing arrow (“Go” stimulus), and participants are instructed to indicate the direction of the arrow, responding “as quickly and accurately as possible”. When the arrow is followed unpredictably by a “Stop Signal” (an up-right arrow presented for 300 ms) participants are required to withhold or interrupt the motor response to the “Go” stimulus. For the SST we considered measures derived from the following contrasts that are relevant to inhibitory control: *incorrect stop vs correct go* and *correct stop vs correct go*.

For the EN-back task, depending on the condition, children needed to respond whether the stimulus was the same as (1) the one shown 2 trials earlier (2-back) or (2) the target stimulus shown at the beginning (0-back). Stimuli included houses and emotional (happy or fearful) and non-emotional faces. For the EN-back task we considered measures derived from the *2-back vs 0-back* (i.e., sensorimotor control) contrast regardless of stimulus valence as an index of working memory and measures derived from the contrast *emotional vs neutral faces* as an index of emotional valence processing independent of working-memory.

ABCD unrelated participants with complete suicidality data were excluded from the task-based fMRI analyses for the following technical and medical reasons:

- **Monetary incentive delay task (MID)**

(1) Incidental findings (e.g. hydrocephalus, herniation), unsatisfactory quality control (QC) for MID fMRI, unsatisfactory T1, incomplete or poorly Freesurfer deconstruction, excessive motion, unacceptable performance, or missing information in these indicators (**n=824**)

Txt file= abcd_mrfindings01; scan excluded if `mrif_score ≥ 3`, `mrif_hydrocephalus="yes"`, `mrif_herniation="yes"`.

Txt file= ABCD MRI QC Raw Part 2; item= `iqc_mid_ok_ser`; scan excluded if 0

Txt file= ABCD MRI QC Raw Part 1; item= `iqc_t1_ok_ser`; scan excluded if 0

Txt file= FreeSurfer QC; item= fsqc_qc; scan excluded if 0

Txt file= ABCD Task fMRI MID Average Beta Weights Part 1: include if
tfMRI_mid_all_b_dof > 200

Txt file= ABCD Task fMRI MID Behavior: include if tfMRI_mid_beh_performflag = 1 and
tfMRI_mid_all_beh_t_nt = 100

(2) Missing ROI-based tabulated data (**n=1932**)

(3) Phillips scan data (**n=630**)

- **Stop signal task (SST)**

(1) Incidental findings (e.g. hydrocephalus, herniation), unsatisfactory quality control (QC) for SST fMRI, unsatisfactory T1, incomplete or poorly Freesurfer deconstruction, excessive motion, unacceptable performance, or missing information in these indicators (**n=1047**)

Txt file = abcd_mrfindings01; scan excluded if mrif_score ≥ 3 , mrif_hydrocephalus="yes",
mrif_herniation="yes".

Txt file= ABCD MRI QC Raw Part 2: item= iqc_sst_ok_ser; scan excluded if 0

Txt file= ABCD MRI QC Raw Part 1; item= iqc_t1_ok_ser; scan excluded if 0

Txt file= FreeSurfer QC; item= fsqc_qc; scan excluded if 0

Txt file= ABCD Task fMRI SST Average Beta Weights Part 1: include if tfMRI_sst_all_b_dof
> 200

Txt file= ABCD Task fMRI SST Behavior: include if tfMRI_sst_beh_performflag = 1

(2) Missing ROI-based tabulated data (**n=2088**)

(3) Phillips scan data (**n=576**)

- **Emotional n-back (EN-back)**

(1) Incidental findings (e.g. hydrocephalus, herniation), unsatisfactory quality control (QC) for SST fMRI, unsatisfactory T1, incomplete or poorly Freesurfer deconstruction, excessive motion, unacceptable performance, or missing information in these indicators (**n=1272**)

Txt file = abcd_mrfindings01; scan excluded if mrif_score ≥ 3 , mrif_hydrocephalus="yes",
mrif_herniation="yes".

Txt file= ABCD MRI QC Raw Part 2: item= iqc_nback_ok_ser; scan excluded if 0


Txt file= ABCD MRI QC Raw Part 1; item= iqc_t1_ok_ser; scan excluded if 0

Txt file= FreeSurfer QC; item= fsqc_qc; scan excluded if 0

Txt file= ABCD Task fMRI nBack Average Beta Weights Part 1: include if
 tfmri_nback_all_b_dof > 200

Txt file= ABCD Task fMRI nBack Behavior: include if tfmri_nback_beh_performflag = 1

- (2) Missing ROI-based tabulated data (**n=2119**)
- (3) Phillips scan data (**n=528**)



eFigure 1. Sample selection flowchart

4.4 Imaging data preparation for analysis

Prior to statistical analysis, the structural and resting-state imaging variables were adjusted for site/scanner effect using ComBat in R (<https://github.com/Jfortin1/ComBatHarmonization>)^{30,31}.

Initially used for batch adjustment of genetic data, ComBat uses Empirical Bayes to adjust for between-site variability while preserving biological variability. We did not apply the ComBat method on the task-based fMRI data because using within-individual contrasts alone in the ABCD task-based fMRI data can mitigate scanner-related variance³¹. Finally, for each conducted test with resting-state

and task-based fMRI measures, we calculated the interquartile range (IQR) in order to remove outlier values that were above $1.5 \times \text{IQR}$.

4.5 Statistical analyses of neuroimaging data

4.5.1 Testing for difference and equivalence

The association between suicidality and imaging measures was examined using two complementary approaches. We examined *differences* between groups endorsing suicidality and the never-suicidal group with two-independent sample t-tests; we used Welch's t-tests to account for unequal number of observations. Of note, Welch's t-test is a traditional null hypothesis significance test, in which the null hypothesis is that the true ES of the mean difference between groups is zero. Whereas with this test one can conclude that the effect is statistically different from zero if the null hypothesis is rejected, the absence of effect cannot be statistically supported. That is, after performing the test, if the *p*-value is larger than the prespecified threshold (e.g., 0.05), it is not possible to conclude that there is no effect or that the effect equals zero. It might simply be that the study had not power enough to detect differences. Moreover, regardless of whether a statistically significant effect is found or not, one might want to examine whether such an effect is large enough to be considered meaningful. In these cases, one can examine whether mean differences can be considered equivalent. For these reasons, and given the size of the current study, we examine *equivalence* performing Equivalence tests (which also included Welch's t-tests to account for unequal number of observations). Equivalence testing has been successfully used in prior studies using the ABCD cohort³² as a safeguard against Type II errors (i.e., false negative results) and against overinterpreting positive results that may arise in large samples despite group differences being minimal (i.e., statistical overpowering)³³. In equivalence testing, the observed data are statistically compared against *a priori* specified equivalence interval (δ), defined by upper (Δ_U) and lower ($-\Delta_L$) equivalence bounds. The aim of equivalence testing is to reject the null hypothesis that the observed ES (Cohen's *d*) is at least as extreme as the smallest effect size of interest (SESOI) (i.e. the 90%CI of the ES overlaps with the SESOI). Here, we used the "two one-sided tests" (TOST) procedure^{34,35} implemented with the *TOSTtwo* function from the library *TOSTER* in R. Given the current sample size and previous results in large samples³⁶, the upper (Δ_U) and lower ($-\Delta_L$) equivalence bounds were specified as $d=0.15$ and $d=-0.15$ (i.e., $\text{SESOI}=|0.15|$), which correspond to traditional notions of a "small" ES³⁷. That is, in the current study, equivalence was defined as an observed ES with 90% confidence intervals (CI) within -0.15 and 0.15.

Since null-hypothesis significant tests, such as Welch's t-test, and equivalence tests examine two different hypotheses, it is recommended to use both simultaneously³⁸. When both tests are performed, there are four possible outcomes: the effect can be 1) statistically different from zero and statistically

equivalent (this is typical of studies with large sample sizes which are powered to detect minimal differences, that are not meaningful effects), 2) not statistically different from zero and statistically equivalent (this is usually the case when the ES is practically zero, not a meaningful effect), 3) statistically different from zero and not statistically equivalent (this is when the ES of the difference is large enough to be considered meaningful), and 4) finally, if the observed ES is neither statistically different from zero nor statistically equivalent, there is insufficient data to draw conclusions and is considered an undetermined result (this is typical of studies with small sample sizes which, albeit reporting ES larger than the pre-specified SESOI, the 90%CI also includes zero). The threshold for statistical significance for both tests was set at $p < 0.05$ after applying FDR-correction for multiple comparisons.

4.5.2. Testing for predictive value

To examine the ability of neural correlates to predict suicide cases in our sample we used Precision-Recall Curves (PRC), a widely used tool in the prediction of machine learning algorithms. In contrast to Receiver Operating Characteristic (ROC) curves, PRC provide more accurate information on the performance of a prediction model in cases where there is an imbalance in the observations between the two classes (i.e., when there are many negative cases and very few positive cases, like suicidality in community-based samples)³⁹. In those cases, ROC curves can provide an inflated view of performance, and thus the use PRC is preferred. Precision can be defined as how good a model is at predicting true positive cases (i.e., positive predictive value, PPV) and is calculated as the ratio of the number of true positives divided by the sum of the true positives and false positives:

$$Precision (PPV) = TP / (TP + FP)$$

Recall can be defined as how good a model is at predicting all the true positive cases (i.e., Sensitivity) and is calculated as the ratio of the number of true positives divided by the sum of the true positives and the false negatives:

$$Recall (Sensitivity) = TP / (TP + FN)$$

PRC plots show precision values for different thresholds of sensitivity (recall) values. For all effect sizes equal or over our SESOI (i.e., $d = |0.15|$), We calculated the area under the PRC (AUPRC), which can be interpreted as the average precision across all sensitivity thresholds. The AUC of a random classifier using a ROC approach would be 0.5, given a balanced class distribution. However, with the PRC approach, the AUC for a random classifier would be close to the prevalence of positive cases in

the population calculated as $y=P/(P+N)$. On the other hand, a perfect classifier, as in ROC, would have an AUC of 1.

eResults

1. Prevalence of suicidality

eTable 2. Suicidal ideation, plans and attempts in unrelated ABCD participants (N=7994)			
KSADS-5 Items	Child-reported suicidality	Caregiver-reported suicidality	Agreement
Past	n (%)	n (%)	n (%) ^a
Passive suicidal ideation	469 (5.9)	550 (6.9)	123 (13.7)
Active but nonspecific suicidal ideation	294 (3.7)	270 (3.8)	54 (10.6)
Suicidal ideation with a specific method	75 (0.9)	70 (0.9)	5 (3.6)
Active suicidal ideation with an intent	47 (0.6)	30 (0.4)	2 (2.7)
Active suicidal ideation with a plan	29 (0.4)	10 (0.1)	0 (0.0)
Preparatory actions toward imminent suicidal behavior	34 (0.4)	38 (0.5)	0 (0.0)
Interrupted suicidal attempts	10 (0.1)	5 (0.1)	0 (0.0)
Aborted suicidal attempts	24 (0.3)	8 (0.1)	0 (0.0)
Suicidal attempts	59 (0.7)	23 (0.3)	9 (12.3)
Current			
Passive suicidal ideation	98 (1.2)	48 (0.6)	6 (4.3)
Active but nonspecific suicidal ideation	126 (1.6)	93 (1.2)	10 (4.8)
Suicidal ideation with a specific method	47 (0.6)	14 (0.2)	1 (1.7)
Active suicidal ideation with an intent	25 (0.3)	7 (0.1)	1 (3.2)
Active suicidal ideation with a plan	20 (0.3)	3 (0.0)	0 (0.0)
Preparatory actions towards imminent suicidal behavior	17 (0.2)	6 (0.1)	0 (0.0)
Interrupted suicidal attempts	4 (0.1)	0 (0.0)	0 (0.0)
Aborted suicidal attempts	16 (0.2)	1 (0.0)	0 (0.0)
Suicidal attempts	15 (0.2)	5 (0.1)	1 (5.3)
Total (any endorsement)	684 (8.6)	654 (8.2)	198 (17.4)
Child or caregivers could have endorsed more than one item; KSADS-5=Kiddie Schedule for Affective Disorders and Schizophrenia for DSM5			
^a Percentage of positive agreement between children and caregivers that endorsed the item			

Neuroimaging

2. Summary of findings

eTable 3. Number and percentage of cortical regions found for each possible combined outcome of the Welch’s t-test and the Equivalence test by imaging modality and suicidality group comparison – FDR-corrected results ^a

Task	Contrast	Suicidality informant	Statistically equivalent <i>and</i> no statistically different, n (%)	Statistically equivalent <i>and</i> statistically different, n (%)	No statistically equivalent <i>and</i> no statistically different, n (%)	No statistically equivalent <i>and</i> statistically different, n (%)	Overlap of findings across groups with no evidence of equivalence <i>and</i> different (uncorrected)
sMRI							
-		Child	62 (72.1)	0 (0)	24 (27.9)	0 (0)	Thickness None (↓left bank of the superior temporal sulcus, only in caregiver analysis)
		Caregiver	76 (88.4)	0 (0)	9 (10.5)	1 (1.2)	
		Concordant	0 (0)	0 (0)	86 (100)	0 (0)	
rs-fMRI							
-		Child	170 (55.6)	0 (0)	136 (44.4)	0 (0)	Connectivity None
		Caregiver	0 (0)	0 (0)	306 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	306 (100)	0 (0)	
Task-based fMRI							
MID							
<i>Reward anticipation</i>		Child	53 (31.7)	0 (0)	114 (68.3)	0 (0)	Activation None
		Caregiver	87 (52.1)	0 (0)	80 (47.9)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
<i>Loss anticipation</i>		Child	120 (71.9)	0 (0)	47 (28.1)	0 (0)	None
		Caregiver	79 (47.3)	0 (0)	88 (52.7)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
<i>Reward feedback</i>		Child	57 (34.1)	0 (0)	110 (65.9)	0 (0)	None
		Caregiver	116 (69.5)	0 (0)	51 (30.5)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
<i>Loss feedback</i>		Child	126 (75.4)	0 (0)	41 (24.6)	0 (0)	None
		Caregiver	0 (0)	0 (0)	167 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
SST							
<i>Incorrect stop vs correct go</i>		Child	98 (58.7)	0 (0)	69 (41.3)	0 (0)	None
		Caregiver	0 (0)	0 (0)	167 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
<i>Correct stop vs correct go</i>		Child	56 (33.5)	0 (0)	111 (66.5)	0 (0)	None
		Caregiver	0 (0)	0 (0)	167 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
EN-back							
<i>2-back vs 0-back</i>		Child	0 (0)	0 (0)	167 (100)	0 (0)	None
		Caregiver	0 (0)	0 (0)	167 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
<i>Emotional vs neutral faces</i>		Child	74 (44.3)	0 (0)	93 (55.7)	0 (0)	None
		Caregiver	0 (0)	0 (0)	167 (100)	0 (0)	
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	

^a Evidence of equivalence (Equivalence test, $p_{FDR} < 0.05$, 90% confidence interval (CI) does not overlap with bounds); No evidence of equivalence (Equivalence test, $p_{FDR} \geq 0.05$, 90%CI overlaps with bound/s); No different (Welch’s t-test, $p_{FDR} < 0.05$, 95%CI includes zero), Different (Welch’s t-test, $p_{FDR} < 0.05$, 95%CI does not include zero).

eTable 4. Number and percentage of cortical regions found for each possible combined outcome of the Welch's t-test and the Equivalence test by imaging modality and suicidality group comparison – Uncorrected results ^a

Modality	Contrast	Suicidality informant	Statistically equivalent and no statistically different, n (%)	Statistically equivalent and statistically different, n (%)	No statistically equivalent and no statistically different, n (%)	No statistically equivalent and statistically different, n (%)	Overlap of findings across groups that are not statistically equivalent and statistically different (uncorrected)
sMRI							
-		Child	67 (77.9)	0 (0)	9 (10.5)	10 (11.6)	Thickness ↓lingual gyrus (ch, co)
		Caregiver	77 (89.5)	0 (0)	4 (4.7)	5 (5.8)	↓left bank of the superior temporal sulcus (ca, co)
		Concordant	9 (10.5)	0 (0)	72 (83.7)	5 (5.8)	↑medial orbitofrontal gyrus (ca, co)
rs-fMRI							
-		Child	219 (71.6)	0 (0)	75 (24.5)	12 (3.9)	Connectivity ↑Cingulo parietal – left hippocampus (ch, ca)
		Caregiver	162 (52.9)	0 (0)	118 (38.6)	26 (8.5)	↓Default – right hippocampus (ch, ca, co)
		Concordant	0 (0)	0 (0)	294 (96.1)	12 (3.9)	↑Dorsal attention – left cerebellum (ca, co) ↓Salience – left amygdala (ch, co) ↓Salience – left accumbens area (ch, co)
Task-based fMRI							
MID							
	<i>Reward anticipation</i>	Child	97 (58.1)	0 (0)	51 (30.5)	19 (11.4)	Activation ↓ left cuneus gyrus (ch, co)
		Caregiver	120 (71.9)	0 (0)	44 (26.3)	3 (1.8)	↓left temporal superior plan temporal gyrus (ch, co)
		Concordant	0 (0)	0 (0)	160 (95.8)	7 (4.2)	↓bilateral calcarine sulcus (ch, ca, co) ↓bilateral parieto-occipital sulcus (ch, co) ↓right occipito-temporal medial lingual gyrus (ch, co)
	<i>Loss anticipation</i>	Child	127 (76.0)	0 (0)	37 (22.2)	3 (1.8)	↓ left paracentral gyrus and sulcus (ca, co)
		Caregiver	113 (67.7)	0 (0)	43 (25.7)	11 (6.6)	↓left temporal superior plan polar gyrus (ca, co)
		Concordant	0 (0)	0 (0)	158 (94.6)	9 (5.4)	↓left temporal pole (ch, ca) ↓left central sulcus (ca, co)
	<i>Reward feedback</i>	Child	109 (65.3)	0 (0)	54 (32.3)	4 (2.4)	↓left middle posterior cingulate gyrus and sulcus (ch, co)
		Caregiver	108 (64.7)	0 (0)	52 (31.1)	7 (4.2)	↓left postcentral gyrus (ca, co)
		Concordant	0 (0)	0 (0)	134 (80.2)	33 (19.8)	↓bilateral central sulcus (ca, co) ↓bilateral superior circular insula sulcus (ca, co) ↓right subcentral gyrus and sulcus (ca, co)
	<i>Loss feedback</i>	Child	136 (81.4)	0 (0)	30 (18.0)	1 (0.6)	
		Caregiver	115 (68.9)	0 (0)	45 (26.9)	7 (4.2)	None
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
SST							
	<i>Incorrect stop vs correct go</i>	Child	127 (76.0)	0 (0)	36 (21.6)	4 (2.4)	
		Caregiver	110 (65.9)	0 (0)	53 (31.7)	4 (2.4)	↓ right inferior temporal sulcus (ch, ca)
		Concordant	0 (0)	0 (0)	167 (100)	0 (0)	
	<i>Correct stop vs correct go</i>	Child	104 (62.3)	0 (0)	57 (34.1)	6 (3.6)	
		Caregiver	95 (56.9)	0 (0)	64 (38.3)	8 (4.8)	↓ right inferior frontal orbital gyrus (ca, co)
		Concordant	0 (0)	0 (0)	164 (98.2)	3 (1.8)	
EN-back							
	<i>2-back vs 0-back</i>	Child	105 (62.9)	0 (0)	57 (34.1)	5 (3.0)	↓ left transverse temporal sulcus (ch, co)
		Caregiver	102 (61.2)	0 (0)	60 (35.9)	5 (3.0)	↓right lateral occipito-temporal sulcus (ca, co)
		Concordant	0 (0)	0 (0)	161 (96.4)	6 (3.6)	
	<i>Emotional</i>	Child	110 (65.9)	0 (0)	54 (32.3)	3 (1.8)	↓bilateral circular insula sulcus (ch, co)

<i>vs neutral faces</i>	Caregiver Concordant	94 (56.3) 0 (0)	0 (0) 0 (0)	64 (38.3) 155 (92.8)	9 (5.4) 12 (7.2)	↓right lateral occipito-temporal sulcus (ch, co)
-------------------------	----------------------	--------------------	----------------	-------------------------	---------------------	--

^aEvidence of equivalence (Equivalence test, $p_{unc} < 0.05$, 90% confidence interval (CI) does not overlap with bounds); No evidence of equivalence (Equivalence test, $p_{unc} \geq 0.05$, 90%CI overlaps with bound/s); No different (Welch's t-test, $p_{unc} < 0.05$, 95%CI includes zero), Different (Welch's t-test, $p_{unc} < 0.05$, 95%CI does not include zero). MID, Monetary incentive delay task. SST, Stop signal task. EN-back, Emotional n-back task. Ch, child-reported suicidality. Ca, caregiver-reported suicidality. Co, concordantly-reported suicidality.

3. Brain Structural Imaging

eTable 5. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the child-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
Cortical thickness					
lh.Banks.of.Superior.Temporal.Sulcus	-0.010	-0.022, 0.003	-1.53	0.127	0.408
lh.caudalanteriorcingulate	0.023	0.003, 0.042	2.30	0.022	0.222
lh.caudalmiddlefrontal	-0.005	-0.014, 0.004	-1.04	0.299	0.658
lh.cuneus	-0.004	-0.016, 0.008	-0.60	0.549	0.777
lh.entorhinal	-0.006	-0.034, 0.022	-0.45	0.656	0.811
lh.fusiform	0.005	-0.004, 0.014	1.12	0.263	0.641
lh.inferiorparietal	-0.004	-0.011, 0.004	-0.98	0.329	0.658
lh.inferiortemporal	0.000	-0.01, 0.01	0.05	0.961	0.974
lh.isthmuscingulate	0.006	-0.011, 0.022	0.70	0.486	0.752
lh.lateraloccipital	-0.002	-0.01, 0.006	-0.42	0.676	0.814
lh.lateralorbitofrontal	-0.002	-0.012, 0.007	-0.49	0.625	0.804
lh.lingual	-0.011	-0.02, -0.002	-2.33	0.020	0.222
lh.medialorbitofrontal	0.015	0.002, 0.028	2.28	0.023	0.222
lh.middletemporal	-0.005	-0.017, 0.007	-0.86	0.392	0.696
lh.parahippocampal	-0.029	-0.052, -0.006	-2.48	0.013	0.222
lh.paracentral	-0.007	-0.017, 0.002	-1.55	0.121	0.408
lh.parsopercularis	0.000	-0.009, 0.01	0.07	0.946	0.974
lh.parsorbitalis	0.017	0.001, 0.032	2.15	0.032	0.272
lh.parstriangularis	0.001	-0.01, 0.012	0.23	0.816	0.895
lh.pericalcarine	-0.005	-0.016, 0.006	-0.92	0.357	0.693
lh.postcentral	-0.002	-0.012, 0.007	-0.52	0.604	0.804
lh.posteriorcingulate	0.012	0, 0.024	1.90	0.058	0.327
lh.precentral	-0.008	-0.016, 0	-1.95	0.051	0.316
lh.precuneus	-0.004	-0.012, 0.003	-1.12	0.264	0.641
lh.rostralanteriorcingulate	0.013	-0.003, 0.03	1.59	0.113	0.408
lh.rostralmiddlefrontal	0.006	-0.001, 0.014	1.63	0.103	0.408
lh.superiorfrontal	0.007	-0.001, 0.015	1.71	0.088	0.408
lh.superiorparietal	-0.004	-0.011, 0.002	-1.27	0.205	0.537
lh.superiortemporal	-0.001	-0.011, 0.009	-0.10	0.918	0.974
lh.supramarginal	-0.004	-0.012, 0.004	-0.90	0.369	0.696
lh.frontalpole	0.021	-0.006, 0.047	1.54	0.124	0.408
lh.temporalpole	-0.001	-0.024, 0.022	-0.08	0.932	0.974
lh.transversetemporal	-0.013	-0.029, 0.003	-1.64	0.101	0.408
lh.insula	0.005	-0.005, 0.016	1.00	0.318	0.658

eTable 5. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the child-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
rh.Banks.of.Superior.Temporal.Sulcus	0.007	-0.006, 0.02	1.02	0.308	0.658
rh.caudalanteriorcingulate	0.017	-0.003, 0.036	1.68	0.094	0.408
rh.caudalmiddlefrontal	0.004	-0.006, 0.014	0.80	0.423	0.719
rh.cuneus	-0.005	-0.016, 0.006	-0.87	0.386	0.696
rh.entorhinal	0.008	-0.024, 0.039	0.47	0.639	0.805
rh.fusiform	-0.004	-0.013, 0.004	-0.99	0.321	0.658
rh.inferiorparietal	0.002	-0.005, 0.009	0.54	0.592	0.804
rh.inferiortemporal	-0.003	-0.013, 0.006	-0.65	0.514	0.752
rh.isthmuscingulate	0.016	0, 0.032	1.98	0.049	0.316
rh.lateraloccipital	-0.006	-0.015, 0.002	-1.45	0.147	0.434
rh.lateralorbitofrontal	0.007	-0.004, 0.017	1.30	0.195	0.537
rh.lingual	-0.003	-0.012, 0.006	-0.64	0.520	0.752
rh.medialorbitofrontal	0.017	0.004, 0.03	2.50	0.013	0.222
rh.middletemporal	0.008	-0.002, 0.018	1.51	0.132	0.408
rh.parahippocampal	-0.025	-0.046, -0.005	-2.45	0.015	0.222
rh.paracentral	-0.004	-0.014, 0.006	-0.84	0.399	0.696
rh.parsopercularis	0.008	-0.002, 0.018	1.51	0.132	0.408
rh.parsorbitalis	0.016	0.001, 0.031	2.06	0.040	0.301
rh.parstriangularis	0.003	-0.006, 0.013	0.67	0.503	0.752
rh.pericalcarine	-0.002	-0.013, 0.01	-0.29	0.773	0.861
rh.postcentral	-0.002	-0.012, 0.008	-0.39	0.694	0.814
rh.posteriorcingulate	0.003	-0.01, 0.015	0.41	0.683	0.814
rh.precentral	-0.003	-0.011, 0.005	-0.74	0.460	0.752
rh.precuneus	-0.002	-0.01, 0.006	-0.54	0.592	0.804
rh.rostralanteriorcingulate	0.000	-0.018, 0.019	0.05	0.959	0.974
rh.rostralmiddlefrontal	0.005	-0.004, 0.013	1.09	0.276	0.647
rh.superiorfrontal	0.010	0.001, 0.018	2.29	0.023	0.222
rh.superiorparietal	-0.001	-0.008, 0.006	-0.29	0.772	0.861
rh.superiortemporal	0.003	-0.006, 0.013	0.65	0.518	0.752
rh.supramarginal	-0.001	-0.01, 0.007	-0.31	0.759	0.861
rh.frontalpole	-0.016	-0.041, 0.009	-1.27	0.205	0.537
rh.temporalpole	0.006	-0.019, 0.031	0.49	0.627	0.804
rh.transversetemporal	-0.005	-0.021, 0.01	-0.68	0.499	0.752
rh.insula	0.000	-0.012, 0.012	0.03	0.974	0.974
Subcortical volumes					
left.lateral.ventricle	-56.5	-274.1, 161.1	-0.42	0.676	1.000
left.cerebellum.cortex	218.3	-168.6, 605.2	1.15	0.252	1.000
left.thalamus.proper	25.2	-18.6, 68.9	1.04	0.299	1.000
left.caudate	-5.8	-45.5, 33.9	-0.47	0.640	1.000
left.putamen	-45.6	-99.5, 8.3	-1.74	0.082	1.000
left.pallidum	-18.3	-38, 1.3	-2.04	0.042	1.000
left.hippocampus	-17.7	-49.8, 14.4	-0.66	0.511	1.000
left.amygdala	-11.3	-27.7, 5.1	-0.97	0.332	1.000

eTable 5. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the child-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
left.accumbens.area	-0.2	-8.8, 8.5	-0.10	0.923	1.000
right.lateral.ventricle	-55.1	-267.3, 157.1	-0.37	0.710	1.000
right.cerebellum.cortex	182.9	-214.2, 580	1.03	0.302	1.000
right.thalamus.proper	23.3	-20.2, 66.9	0.78	0.437	1.000
right.caudate	-4.5	-44.1, 35	-0.62	0.535	1.000
right.putamen	-38.3	-85, 8.3	-1.79	0.075	1.000
right.pallidum	-3.1	-18.9, 12.6	-0.58	0.564	1.000
right.hippocampus	-6.8	-36.9, 23.3	-0.29	0.772	1.000
right.amygdala	-4.7	-20.5, 11	-0.20	0.843	1.000
right.accumbens.area	-0.4	-8, 7.2	-0.13	0.897	1.000

Bold denotes group differences at $p < 0.05$ uncorrected

eTable 6. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
Cortical thickness					
lh.Banks.of.Superior.Temporal.Sulcus	-0.024	-0.036, -0.011	-3.66	0.000	0.019
lh.caudalanteriorcingulate	0.013	-0.008, 0.034	1.24	0.216	0.984
lh.caudalmiddlefrontal	-0.002	-0.011, 0.007	-0.43	0.666	0.984
lh.cuneus	-0.001	-0.012, 0.01	-0.20	0.843	0.984
lh.entorhinal	0.009	-0.02, 0.039	0.64	0.523	0.984
lh.fusiform	0.004	-0.004, 0.012	1.01	0.311	0.984
lh.inferiorparietal	-0.003	-0.01, 0.004	-0.77	0.441	0.984
lh.inferiortemporal	0.001	-0.009, 0.011	0.18	0.854	0.984
lh.isthmuscingulate	0.003	-0.013, 0.019	0.35	0.726	0.984
lh.lateraloccipital	0.002	-0.007, 0.011	0.40	0.692	0.984
lh.lateralorbitofrontal	-0.005	-0.015, 0.006	-0.90	0.369	0.984
lh.lingual	-0.003	-0.013, 0.007	-0.61	0.541	0.984
lh.medialorbitofrontal	0.005	-0.008, 0.018	0.71	0.477	0.984
lh.middletemporal	-0.005	-0.016, 0.007	-0.76	0.447	0.984
lh.parahippocampal	0.005	-0.019, 0.03	0.42	0.673	0.984
lh.paracentral	-0.013	-0.024, -0.002	-2.37	0.018	0.616
lh.parsopercularis	-0.004	-0.014, 0.006	-0.83	0.405	0.984
lh.parsorbitalis	0.006	-0.011, 0.023	0.70	0.483	0.984
lh.parsstriangularis	-0.001	-0.012, 0.01	-0.21	0.833	0.984
lh.pericalcarine	-0.008	-0.02, 0.004	-1.37	0.172	0.984
lh.postcentral	0.005	-0.004, 0.014	1.08	0.282	0.984
lh.posteriorcingulate	0.000	-0.013, 0.013	0.02	0.984	0.984
lh.precentral	-0.003	-0.011, 0.005	-0.72	0.474	0.984

eTable 6. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
lh.precuneus	-0.004	-0.012, 0.004	-0.96	0.336	0.984
lh.rostralanteriorcingulate	0.011	-0.007, 0.028	1.21	0.228	0.984
lh.rostralmiddlefrontal	0.003	-0.005, 0.011	0.74	0.462	0.984
lh.superiorfrontal	-0.001	-0.009, 0.008	-0.16	0.876	0.984
lh.superiorparietal	0.001	-0.006, 0.008	0.36	0.719	0.984
lh.superiortemporal	0.001	-0.01, 0.012	0.17	0.869	0.984
lh.supramarginal	-0.003	-0.012, 0.005	-0.73	0.465	0.984
lh.frontalpole	0.007	-0.021, 0.034	0.49	0.625	0.984
lh.temporalpole	0.012	-0.012, 0.036	0.99	0.324	0.984
lh.transversetemporal	-0.010	-0.026, 0.006	-1.23	0.218	0.984
lh.insula	0.007	-0.004, 0.019	1.31	0.191	0.984
rh.Banks.of.Superior.Temporal.Sulcus	0.000	-0.014, 0.014	0.03	0.977	0.984
rh.caudalanteriorcingulate	-0.003	-0.023, 0.017	-0.28	0.782	0.984
rh.caudalmiddlefrontal	0.004	-0.005, 0.014	0.87	0.387	0.984
rh.cuneus	0.001	-0.01, 0.012	0.11	0.913	0.984
rh.entorhinal	0.006	-0.027, 0.038	0.35	0.727	0.984
rh.fusiform	-0.002	-0.01, 0.006	-0.57	0.569	0.984
rh.inferiorparietal	0.005	-0.003, 0.012	1.25	0.213	0.984
rh.inferiortemporal	0.001	-0.009, 0.011	0.18	0.858	0.984
rh.isthmuscingulate	0.013	-0.005, 0.031	1.44	0.149	0.984
rh.lateraloccipital	0.000	-0.008, 0.009	0.10	0.917	0.984
rh.lateralorbitofrontal	0.004	-0.006, 0.015	0.82	0.410	0.984
rh.lingual	-0.001	-0.011, 0.009	-0.23	0.815	0.984
rh.medialorbitofrontal	0.009	-0.005, 0.022	1.26	0.207	0.984
rh.middletemporal	-0.002	-0.013, 0.008	-0.43	0.670	0.984
rh.parahippocampal	-0.003	-0.024, 0.018	-0.26	0.794	0.984
rh.paracentral	-0.009	-0.019, 0.001	-1.78	0.076	0.984
rh.parsopercularis	0.000	-0.011, 0.01	-0.06	0.956	0.984
rh.parsorbitalis	0.011	-0.005, 0.027	1.38	0.168	0.984
rh.parstriangularis	-0.001	-0.011, 0.01	-0.10	0.920	0.984
rh.pericalcarine	-0.005	-0.017, 0.007	-0.88	0.380	0.984
rh.postcentral	0.002	-0.009, 0.013	0.42	0.677	0.984
rh.posteriorcingulate	0.000	-0.013, 0.012	-0.06	0.949	0.984
rh.precentral	-0.002	-0.011, 0.006	-0.50	0.615	0.984
rh.precuneus	0.000	-0.008, 0.008	-0.11	0.909	0.984
rh.rostralanteriorcingulate	0.001	-0.017, 0.02	0.14	0.885	0.984
rh.rostralmiddlefrontal	0.005	-0.004, 0.013	1.08	0.280	0.984
rh.superiorfrontal	0.002	-0.006, 0.011	0.57	0.572	0.984
rh.superiorparietal	0.001	-0.006, 0.009	0.40	0.691	0.984
rh.superiortemporal	0.007	-0.003, 0.017	1.37	0.172	0.984
rh.supramarginal	-0.004	-0.013, 0.006	-0.76	0.450	0.984
rh.frontalpole	-0.003	-0.031, 0.024	-0.22	0.827	0.984
rh.temporalpole	0.028	0.003, 0.053	2.20	0.028	0.638

eTable 6. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
rh.transversetemporal	0.000	-0.016, 0.016	0.05	0.957	0.984
rh.insula	0.008	-0.005, 0.021	1.25	0.210	0.984
Subcortical volumes					
left.lateral.ventricle	69.0	-161.9, 300	0.59	0.557	0.836
left.cerebellum.cortex	-278.8	-660.8, 103.1	-1.43	0.152	0.547
left.thalamus.proper	-45.8	-90, -1.6	-2.03	0.042	0.417
left.caudate	-6.0	-45.1, 33.1	-0.30	0.764	0.879
left.putamen	0.0	-55.6, 55.6	0.00	1.000	1.000
left.pallidum	-12.6	-33.3, 8.1	-1.20	0.231	0.594
left.hippocampus	-28.2	-60.9, 4.5	-1.69	0.091	0.546
left.amygdala	-5.5	-22.1, 11	-0.66	0.510	0.836
left.accumbens.area	3.1	-5.5, 11.7	0.71	0.478	0.836
right.lateral.ventricle	42.6	-196.5, 281.7	0.35	0.726	0.879
right.cerebellum.cortex	-252.0	-654.6, 150.5	-1.23	0.219	0.594
right.thalamus.proper	-35.1	-80.4, 10.3	-1.52	0.129	0.547
right.caudate	-9.7	-49.4, 30.1	-0.48	0.633	0.876
right.putamen	1.9	-45.3, 49	0.08	0.938	0.993
right.pallidum	-2.3	-18.2, 13.7	-0.28	0.781	0.879
right.hippocampus	-32.4	-64.3, -0.5	-2.00	0.046	0.417
right.amygdala	-5.3	-21.7, 11.1	-0.63	0.526	0.836
right.accumbens.area	-3.1	-11, 4.8	-0.77	0.441	0.836

Bold denotes group differences at p<0.05 uncorrected and corrected

eTable 7. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test


FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
Cortical thickness					
lh.Banks.of.Superior.Temporal.Sulcus	-0.033	-0.055, -0.01	-2.84	0.005	0.181
lh.caudalanteriorcingulate	0.035	-0.003, 0.073	1.82	0.071	0.606
lh.caudalmiddlefrontal	-0.006	-0.021, 0.009	-0.79	0.428	0.823
lh.cuneus	-0.003	-0.023, 0.017	-0.31	0.759	0.886
lh.entorhinal	0.015	-0.035, 0.065	0.59	0.557	0.838
lh.fusiform	0.005	-0.01, 0.02	0.66	0.510	0.838
lh.inferiorparietal	-0.005	-0.018, 0.008	-0.76	0.448	0.823
lh.inferiortemporal	0.000	-0.019, 0.019	0.01	0.993	0.993
lh.isthmuscingulate	-0.018	-0.046, 0.009	-1.32	0.190	0.806
lh.lateraloccipital	-0.006	-0.021, 0.009	-0.82	0.411	0.823
lh.lateralorbitofrontal	0.005	-0.013, 0.023	0.57	0.567	0.838
lh.lingual	-0.021	-0.037, -0.004	-2.49	0.014	0.310

eTable 7. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test


FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
lh.medialorbitofrontal	0.019	-0.003, 0.041	1.73	0.086	0.629
lh.middletemporal	-0.009	-0.029, 0.011	-0.89	0.374	0.823
lh.parahippocampal	-0.009	-0.051, 0.032	-0.45	0.654	0.855
lh.paracentral	-0.016	-0.034, 0.003	-1.69	0.093	0.629
lh.parsopercularis	-0.007	-0.024, 0.01	-0.80	0.427	0.823
lh.parsorbitalis	0.028	-0.002, 0.059	1.83	0.069	0.606
lh.parstriangularis	-0.007	-0.027, 0.012	-0.75	0.452	0.823
lh.pericalcarine	-0.011	-0.032, 0.011	-0.97	0.332	0.806
lh.postcentral	-0.007	-0.022, 0.009	-0.87	0.384	0.823
lh.posteriorcingulate	0.000	-0.02, 0.019	-0.02	0.983	0.993
lh.precentral	0.002	-0.011, 0.015	0.29	0.769	0.886
lh.precuneus	-0.006	-0.019, 0.007	-0.87	0.388	0.823
lh.rostralanteriorcingulate	0.001	-0.03, 0.032	0.09	0.930	0.973
lh.rostralmiddlefrontal	0.010	-0.004, 0.023	1.40	0.163	0.806
lh.superiorfrontal	0.001	-0.014, 0.015	0.10	0.922	0.973
lh.superiorparietal	-0.001	-0.013, 0.011	-0.11	0.915	0.973
lh.superiortemporal	-0.004	-0.023, 0.016	-0.38	0.702	0.884
lh.supramarginal	0.001	-0.012, 0.015	0.22	0.823	0.933
lh.frontalpole	0.029	-0.018, 0.077	1.24	0.218	0.806
lh.temporalpole	0.039	0.001, 0.078	2.02	0.045	0.606
lh.transversetemporal	-0.013	-0.042, 0.016	-0.86	0.390	0.823
lh.insula	0.007	-0.012, 0.026	0.72	0.472	0.823
rh.Banks.of.Superior.Temporal.Sulcus	0.014	-0.008, 0.036	1.22	0.223	0.806
rh.caudalanteriorcingulate	0.013	-0.022, 0.049	0.74	0.463	0.823
rh.caudalmiddlefrontal	0.005	-0.011, 0.022	0.62	0.534	0.838
rh.cuneus	-0.010	-0.028, 0.008	-1.10	0.275	0.806
rh.entorhinal	0.031	-0.029, 0.09	1.02	0.307	0.806
rh.fusiform	0.004	-0.011, 0.018	0.49	0.626	0.851
rh.inferiorparietal	0.010	-0.002, 0.023	1.63	0.106	0.654
rh.inferiortemporal	-0.004	-0.021, 0.012	-0.51	0.610	0.851
rh.isthmuscingulate	0.001	-0.028, 0.03	0.06	0.953	0.982
rh.lateraloccipital	-0.008	-0.024, 0.007	-1.06	0.293	0.806
rh.lateralorbitofrontal	0.017	0.002, 0.033	2.25	0.026	0.436
rh.lingual	-0.010	-0.027, 0.007	-1.18	0.240	0.806
rh.medialorbitofrontal	0.034	0.01, 0.057	2.83	0.005	0.181
rh.middletemporal	0.011	-0.008, 0.03	1.13	0.259	0.806
rh.parahippocampal	-0.006	-0.044, 0.031	-0.34	0.733	0.886
rh.paracentral	-0.019	-0.039, 0.001	-1.92	0.057	0.606
rh.parsopercularis	0.005	-0.013, 0.022	0.50	0.616	0.851
rh.parsorbitalis	0.013	-0.013, 0.039	1.01	0.314	0.806
rh.parstriangularis	0.003	-0.014, 0.02	0.39	0.697	0.884
rh.pericalcarine	-0.005	-0.027, 0.016	-0.51	0.614	0.851

eTable 7. Differences in cortical thickness and subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test


FreeSurfer Derived Regions	Mean	95%CI	t	p (unc)	p FDR
rh.postcentral	-0.003	-0.019, 0.014	-0.35	0.725	0.886
rh.posteriorcingulate	0.002	-0.019, 0.023	0.18	0.854	0.952
rh.precentral	0.004	-0.01, 0.019	0.62	0.537	0.838
rh.precuneus	-0.004	-0.018, 0.009	-0.60	0.547	0.838
rh.rostralanteriorcingulate	-0.005	-0.039, 0.029	-0.30	0.767	0.886
rh.rostralmiddlefrontal	0.001	-0.012, 0.015	0.15	0.879	0.965
rh.superiorfrontal	0.007	-0.006, 0.021	1.06	0.291	0.806
rh.superiorparietal	0.007	-0.006, 0.02	1.14	0.257	0.806
rh.superiortemporal	0.011	-0.006, 0.028	1.29	0.198	0.806
rh.supramarginal	-0.005	-0.022, 0.011	-0.63	0.527	0.838
rh.frontalpole	-0.021	-0.063, 0.02	-1.00	0.317	0.806
rh.temporalpole	0.030	-0.016, 0.076	1.27	0.206	0.806
rh.transversetemporal	-0.014	-0.041, 0.014	-0.98	0.329	0.806
rh.insula	0.006	-0.018, 0.029	0.46	0.645	0.855
Subcortical volumes					
left.lateral.ventricle	-112.3	-468.6, 244.1	-0.62	0.535	0.959
left.cerebellum.cortex	-65.1	-713.2, 583.1	-0.20	0.843	0.959
left.thalamus.proper	19.4	-57.8, 96.5	0.50	0.621	0.959
left.caudate	21.3	-47.6, 90.3	0.61	0.542	0.959
left.putamen	14.1	-84, 112.2	0.28	0.777	0.959
left.pallidum	-30.5	-62.3, 1.4	-1.89	0.061	0.959
left.hippocampus	-30.8	-85.3, 23.7	-1.12	0.266	0.959
left.amygdala	-1.9	-32.9, 29.1	-0.12	0.906	0.959
left.accumbens.area	6.7	-8.7, 22.1	0.86	0.390	0.959
right.lateral.ventricle	-151.8	-485, 181.4	-0.90	0.369	0.959
right.cerebellum.cortex	-7.8	-682.4, 666.9	-0.02	0.982	0.982
right.thalamus.proper	22.9	-56, 101.8	0.57	0.567	0.959
right.caudate	18.9	-48.3, 86.2	0.56	0.579	0.959
right.putamen	12.5	-74.3, 99.3	0.28	0.776	0.959
right.pallidum	1.9	-28.8, 32.5	0.12	0.903	0.959
right.hippocampus	10.6	-42.6, 63.8	0.39	0.695	0.959
right.amygdala	3.4	-25.8, 32.6	0.23	0.819	0.959
right.accumbens.area	2.2	-10.8, 15.2	0.33	0.740	0.959
Bold denotes group differences at $p < 0.05$ uncorrected					




eFigure 2. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 3. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 4. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 5. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 6. Results of the Equivalence test for differences in cortical thickness between the never-suicidal group and the concordantly-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 7. Results of the Equivalence test for differences in subcortical volumes between the never-suicidal group and the concordantly-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

4. Resting -State Functional Imaging

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
AuditoryLeftCerebellumCortex	0.008	-0.005, 0.022	1.24	0.215	0.998
AuditoryLeftThalamusProper	0.008	-0.004, 0.02	1.37	0.171	0.998
AuditoryLeftCaudate	-0.012	-0.028, 0.005	-1.35	0.179	0.998
AuditoryLeftPutamen	-0.002	-0.013, 0.01	-0.26	0.793	0.998
AuditoryLeftPallidum	-0.002	-0.011, 0.008	-0.33	0.743	0.998
AuditoryBrainStem	0.000	-0.012, 0.012	0.00	0.997	0.998
AuditoryLeftHippocampus	0.001	-0.006, 0.007	0.13	0.894	0.998
AuditoryLeftAmygdala	-0.008	-0.023, 0.006	-1.14	0.254	0.998
AuditoryLeftAccumbensArea	0.002	-0.01, 0.014	0.27	0.791	0.998
AuditoryLeftVentraldc	-0.002	-0.02, 0.017	-0.16	0.870	0.998
AuditoryRightCerebellumCortex	0.003	-0.015, 0.02	0.30	0.763	0.998
AuditoryRightThalamusProper	0.008	-0.003, 0.019	1.49	0.138	0.998
AuditoryRightCaudate	-0.010	-0.025, 0.004	-1.40	0.162	0.998
AuditoryRightPutamen	-0.006	-0.02, 0.008	-0.82	0.412	0.998
AuditoryRightPallidum	-0.006	-0.018, 0.005	-1.07	0.286	0.998
AuditoryRightHippocampus	0.008	-0.01, 0.025	0.85	0.396	0.998
AuditoryRightAmygdala	0.007	-0.003, 0.017	1.40	0.163	0.998
AuditoryRightAccumbensArea	-0.004	-0.014, 0.006	-0.81	0.421	0.998
AuditoryRightVentraldc	0.001	-0.009, 0.01	0.13	0.900	0.998
CinguloOpercularLeftCerebellumCortex	0.001	-0.005, 0.008	0.44	0.662	0.998
CinguloOpercularLeftThalamusProper	-0.001	-0.016, 0.014	-0.15	0.879	0.998
CinguloOpercularLeftCaudate	-0.003	-0.016, 0.011	-0.37	0.709	0.998
CinguloOpercularLeftPutamen	0.006	-0.013, 0.024	0.60	0.550	0.998
CinguloOpercularLeftPallidum	-0.003	-0.018, 0.013	-0.34	0.733	0.998
CinguloOpercularBrainStem	0.005	-0.007, 0.016	0.80	0.425	0.998
CinguloOpercularLeftHippocampus	-0.005	-0.017, 0.008	-0.72	0.474	0.998
CinguloOpercularLeftAmygdala	0.006	-0.007, 0.019	0.96	0.336	0.998
CinguloOpercularLeftAccumbensArea	-0.008	-0.02, 0.003	-1.42	0.157	0.998
CinguloOpercularLeftVentraldc	0.003	-0.016, 0.021	0.29	0.775	0.998
CinguloOpercularRightCerebellumCortex	0.002	-0.009, 0.013	0.40	0.689	0.998
CinguloOpercularRightThalamusProper	0.006	-0.004, 0.017	1.21	0.226	0.998
CinguloOpercularRightCaudate	0.009	-0.001, 0.018	1.71	0.087	0.998
CinguloOpercularRightPutamen	-0.001	-0.007, 0.005	-0.29	0.769	0.998
CinguloOpercularRightPallidum	-0.005	-0.019, 0.009	-0.70	0.484	0.998
CinguloOpercularRightHippocampus	0.001	-0.011, 0.013	0.22	0.827	0.998
CinguloOpercularRightAmygdala	-0.003	-0.02, 0.014	-0.32	0.750	0.998
CinguloOpercularRightAccumbensArea	0.000	-0.017, 0.016	-0.04	0.967	0.998
CinguloOpercularRightVentraldc	0.001	-0.011, 0.012	0.12	0.905	0.998
CinguloParietalLeftCerebellumCortex	-0.009	-0.022, 0.004	-1.31	0.190	0.998
CinguloParietalLeftThalamusProper	0.001	-0.012, 0.014	0.20	0.841	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
CinguloParietalLeftCaudate	0.001	-0.01, 0.012	0.16	0.874	0.998
CinguloParietalLeftPutamen	-0.008	-0.025, 0.009	-0.93	0.352	0.998
CinguloParietalLeftPallidum	-0.006	-0.017, 0.005	-1.14	0.257	0.998
CinguloParietalBrainStem	-0.002	-0.011, 0.007	-0.46	0.644	0.998
CinguloParietalLeftHippocampus	0.011	0.002, 0.02	2.31	0.021	0.998
CinguloParietalLeftAmygdala	0.000	-0.005, 0.006	0.15	0.879	0.998
CinguloParietalLeftAccumbensArea	-0.006	-0.021, 0.01	-0.71	0.478	0.998
CinguloParietalLeftVentraldc	0.002	-0.01, 0.014	0.28	0.782	0.998
CinguloParietalRightCerebellumCortex	0.001	-0.016, 0.018	0.10	0.924	0.998
CinguloParietalRightThalamusProper	0.000	-0.017, 0.017	0.03	0.979	0.998
CinguloParietalRightCaudate	-0.001	-0.012, 0.011	-0.12	0.904	0.998
CinguloParietalRightPutamen	-0.004	-0.017, 0.009	-0.57	0.568	0.998
CinguloParietalRightPallidum	-0.001	-0.014, 0.012	-0.19	0.847	0.998
CinguloParietalRightHippocampus	0.004	-0.007, 0.014	0.72	0.474	0.998
CinguloParietalRightAmygdala	-0.006	-0.021, 0.01	-0.70	0.482	0.998
CinguloParietalRightAccumbensArea	-0.002	-0.012, 0.008	-0.43	0.667	0.998
CinguloParietalRightVentraldc	0.006	-0.003, 0.015	1.31	0.191	0.998
DefaultLeftCerebellumCortex	0.007	-0.002, 0.016	1.59	0.113	0.998
DefaultLeftThalamusProper	0.004	-0.002, 0.01	1.34	0.181	0.998
DefaultLeftCaudate	0.010	-0.004, 0.025	1.43	0.155	0.998
DefaultLeftPutamen	0.003	-0.009, 0.014	0.49	0.623	0.998
DefaultLeftPallidum	0.001	-0.017, 0.018	0.06	0.953	0.998
DefaultBrainStem	0.001	-0.015, 0.016	0.06	0.951	0.998
DefaultLeftHippocampus	0.002	-0.008, 0.012	0.41	0.681	0.998
DefaultLeftAmygdala	0.002	-0.01, 0.014	0.31	0.757	0.998
DefaultLeftAccumbensArea	0.003	-0.009, 0.016	0.50	0.619	0.998
DefaultLeftVentraldc	0.002	-0.009, 0.013	0.31	0.760	0.998
DefaultRightCerebellumCortex	-0.010	-0.027, 0.007	-1.13	0.259	0.998
DefaultRightThalamusProper	-0.003	-0.013, 0.007	-0.58	0.560	0.998
DefaultRightCaudate	-0.007	-0.016, 0.002	-1.44	0.150	0.998
DefaultRightPutamen	-0.001	-0.011, 0.008	-0.26	0.793	0.998
DefaultRightPallidum	-0.002	-0.009, 0.006	-0.42	0.677	0.998
DefaultRightHippocampus	-0.015	-0.031, 0	-2.00	0.047	0.998
DefaultRightAmygdala	-0.004	-0.016, 0.007	-0.77	0.440	0.998
DefaultRightAccumbensArea	0.008	-0.009, 0.025	0.90	0.370	0.998
DefaultRightVentraldc	-0.004	-0.021, 0.013	-0.49	0.628	0.998
DorsalAttentionLeftCerebellumCortex	0.009	-0.002, 0.02	1.60	0.110	0.998
DorsalAttentionLeftThalamusProper	-0.004	-0.017, 0.008	-0.70	0.481	0.998
DorsalAttentionLeftCaudate	-0.001	-0.013, 0.012	-0.11	0.915	0.998
DorsalAttentionLeftPutamen	0.008	-0.004, 0.021	1.28	0.201	0.998
DorsalAttentionLeftPallidum	-0.018	-0.036, 0	-1.94	0.054	0.998
DorsalAttentionBrainStem	-0.010	-0.021, 0.001	-1.75	0.081	0.998
DorsalAttentionLeftHippocampus	0.012	0.002, 0.022	2.33	0.020	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
DorsalAttentionLeftAmygdala	0.001	-0.008, 0.01	0.22	0.826	0.998
DorsalAttentionLeftAccumbensArea	-0.001	-0.007, 0.006	-0.16	0.875	0.998
DorsalAttentionLeftVentraldc	-0.013	-0.03, 0.004	-1.50	0.135	0.998
DorsalAttentionRightCerebellumCortex	-0.001	-0.012, 0.01	-0.26	0.795	0.998
DorsalAttentionRightThalamusProper	0.001	-0.015, 0.016	0.10	0.923	0.998
DorsalAttentionRightCaudate	-0.008	-0.024, 0.008	-0.97	0.332	0.998
DorsalAttentionRightPutamen	0.004	-0.007, 0.014	0.72	0.473	0.998
DorsalAttentionRightPallidum	0.000	-0.012, 0.011	-0.05	0.962	0.998
DorsalAttentionRightHippocampus	0.001	-0.011, 0.012	0.15	0.880	0.998
DorsalAttentionRightAmygdala	-0.003	-0.014, 0.008	-0.57	0.567	0.998
DorsalAttentionRightAccumbensArea	-0.014	-0.029, 0.002	-1.75	0.081	0.998
DorsalAttentionRightVentraldc	-0.006	-0.016, 0.004	-1.26	0.207	0.998
FrontoParietalLeftCerebellumCortex	0.004	-0.005, 0.013	0.83	0.407	0.998
FrontoParietalLeftThalamusProper	-0.002	-0.011, 0.007	-0.39	0.698	0.998
FrontoParietalLeftCaudate	-0.003	-0.01, 0.003	-0.94	0.346	0.998
FrontoParietalLeftPutamen	-0.013	-0.027, 0.001	-1.82	0.070	0.998
FrontoParietalLeftPallidum	0.000	-0.011, 0.011	0.02	0.986	0.998
FrontoParietalBrainStem	-0.002	-0.018, 0.014	-0.24	0.812	0.998
FrontoParietalLeftHippocampus	-0.009	-0.026, 0.007	-1.11	0.269	0.998
FrontoParietalLeftAmygdala	0.002	-0.008, 0.011	0.31	0.755	0.998
FrontoParietalLeftAccumbensArea	0.000	-0.011, 0.012	0.06	0.951	0.998
FrontoParietalLeftVentraldc	0.005	-0.006, 0.015	0.84	0.400	0.998
FrontoParietalRightCerebellumCortex	0.009	-0.001, 0.019	1.81	0.071	0.998
FrontoParietalRightThalamusProper	-0.009	-0.025, 0.008	-1.03	0.305	0.998
FrontoParietalRightCaudate	-0.001	-0.012, 0.01	-0.23	0.820	0.998
FrontoParietalRightPutamen	0.006	-0.004, 0.016	1.24	0.216	0.998
FrontoParietalRightPallidum	0.003	-0.006, 0.013	0.73	0.465	0.998
FrontoParietalRightHippocampus	0.000	-0.006, 0.007	0.13	0.899	0.998
FrontoParietalRightAmygdala	-0.001	-0.015, 0.013	-0.13	0.894	0.998
FrontoParietalRightAccumbensArea	0.003	-0.008, 0.013	0.47	0.641	0.998
FrontoParietalRightVentraldc	-0.004	-0.019, 0.01	-0.59	0.558	0.998
RetrosplenialTemporalLeftCerebellumCortex	0.006	-0.005, 0.017	1.02	0.309	0.998
RetrosplenialTemporalLeftThalamusProper	-0.001	-0.012, 0.009	-0.25	0.806	0.998
RetrosplenialTemporalLeftCaudate	-0.002	-0.013, 0.008	-0.41	0.683	0.998
RetrosplenialTemporalLeftPutamen	-0.007	-0.013, 0	-2.01	0.045	0.998
RetrosplenialTemporalLeftPallidum	0.005	-0.01, 0.019	0.63	0.528	0.998
RetrosplenialTemporalBrainStem	-0.001	-0.014, 0.011	-0.20	0.843	0.998
RetrosplenialTemporalLeftHippocampus	0.003	-0.014, 0.02	0.38	0.701	0.998
RetrosplenialTemporalLeftAmygdala	0.000	-0.017, 0.016	-0.03	0.978	0.998
RetrosplenialTemporalLeftAccumbensArea	0.002	-0.009, 0.013	0.36	0.721	0.998
RetrosplenialTemporalLeftVentraldc	0.001	-0.013, 0.016	0.20	0.844	0.998
RetrosplenialTemporalRightCerebellumCortex	-0.005	-0.021, 0.01	-0.68	0.495	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
RetrosplenialTemporalRightThalamusProper	-0.013	-0.025, -0.001	-2.06	0.040	0.998
RetrosplenialTemporalRightCaudate	-0.001	-0.019, 0.017	-0.12	0.904	0.998
RetrosplenialTemporalRightPutamen	0.000	-0.01, 0.011	0.04	0.965	0.998
RetrosplenialTemporalRightPallidum	-0.006	-0.016, 0.004	-1.25	0.211	0.998
RetrosplenialTemporalRightHippocampus	0.000	-0.01, 0.009	-0.05	0.961	0.998
RetrosplenialTemporalRightAmygdala	0.002	-0.004, 0.008	0.60	0.550	0.998
RetrosplenialTemporalRightAccumbensArea	0.000	-0.014, 0.015	0.04	0.965	0.998
RetrosplenialTemporalRightVentraldc	-0.005	-0.019, 0.008	-0.75	0.452	0.998
SensorimotorHandLeftCerebellumCortex	-0.001	-0.02, 0.017	-0.13	0.894	0.998
SensorimotorHandLeftThalamusProper	-0.007	-0.023, 0.009	-0.86	0.391	0.998
SensorimotorHandLeftCaudate	0.006	-0.006, 0.017	0.91	0.363	0.998
SensorimotorHandLeftPutamen	0.000	-0.012, 0.013	0.05	0.961	0.998
SensorimotorHandLeftPallidum	0.002	-0.011, 0.015	0.31	0.758	0.998
SensorimotorHandBrainStem	-0.012	-0.024, 0	-2.02	0.044	0.998
SensorimotorHandLeftHippocampus	-0.006	-0.024, 0.012	-0.62	0.535	0.998
SensorimotorHandLeftAmygdala	0.000	-0.011, 0.011	-0.04	0.971	0.998
SensorimotorHandLeftAccumbensArea	0.002	-0.009, 0.012	0.30	0.762	0.998
SensorimotorHandLeftVentraldc	0.005	-0.005, 0.015	1.03	0.306	0.998
SensorimotorHandRightCerebellumCortex	-0.001	-0.008, 0.005	-0.38	0.706	0.998
SensorimotorHandRightThalamusProper	-0.011	-0.025, 0.003	-1.60	0.111	0.998
SensorimotorHandRightCaudate	0.005	-0.008, 0.017	0.70	0.482	0.998
SensorimotorHandRightPutamen	-0.003	-0.02, 0.014	-0.36	0.721	0.998
SensorimotorHandRightPallidum	0.003	-0.013, 0.018	0.34	0.731	0.998
SensorimotorHandRightHippocampus	0.002	-0.009, 0.013	0.31	0.760	0.998
SensorimotorHandRightAmygdala	0.003	-0.01, 0.017	0.47	0.636	0.998
SensorimotorHandRightAccumbensArea	-0.002	-0.015, 0.011	-0.27	0.784	0.998
SensorimotorHandRightVentraldc	0.001	-0.01, 0.012	0.16	0.871	0.998
SensorimotorMouthLeftCerebellumCortex	-0.001	-0.017, 0.015	-0.15	0.882	0.998
SensorimotorMouthLeftThalamusProper	-0.008	-0.019, 0.003	-1.44	0.150	0.998
SensorimotorMouthLeftCaudate	-0.002	-0.012, 0.007	-0.48	0.629	0.998
SensorimotorMouthLeftPutamen	0.006	-0.003, 0.015	1.27	0.203	0.998
SensorimotorMouthLeftPallidum	-0.001	-0.007, 0.005	-0.37	0.709	0.998
SensorimotorMouthBrainStem	0.002	-0.013, 0.016	0.23	0.822	0.998
SensorimotorMouthLeftHippocampus	-0.004	-0.015, 0.008	-0.59	0.557	0.998
SensorimotorMouthLeftAmygdala	-0.007	-0.024, 0.009	-0.85	0.394	0.998
SensorimotorMouthLeftAccumbensArea	0.000	-0.016, 0.017	0.03	0.978	0.998
SensorimotorMouthLeftVentraldc	-0.001	-0.012, 0.011	-0.11	0.909	0.998
SensorimotorMouthRightCerebellumCortex	-0.004	-0.017, 0.008	-0.65	0.515	0.998
SensorimotorMouthRightThalamusProper	-0.002	-0.015, 0.011	-0.33	0.742	0.998
SensorimotorMouthRightCaudate	-0.004	-0.015, 0.008	-0.62	0.537	0.998
SensorimotorMouthRightPutamen	-0.001	-0.017, 0.014	-0.18	0.859	0.998
SensorimotorMouthRightPallidum	0.000	-0.01, 0.009	-0.06	0.956	0.998
SensorimotorMouthRightHippocampus	-0.005	-0.013, 0.004	-1.01	0.312	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
SensorimotorMouthRightAmygdala	0.007	-0.002, 0.016	1.60	0.111	0.998
SensorimotorMouthRightAccumbensArea	0.003	-0.003, 0.009	1.00	0.320	0.998
SensorimotorMouthRightVentraldc	-0.002	-0.015, 0.012	-0.27	0.788	0.998
SaliencLeftCerebellumCortex	0.004	-0.007, 0.015	0.71	0.477	0.998
SaliencLeftThalamusProper	-0.001	-0.018, 0.016	-0.13	0.899	0.998
SaliencLeftCaudate	0.001	-0.015, 0.016	0.09	0.932	0.998
SaliencLeftPutamen	0.007	-0.003, 0.016	1.41	0.158	0.998
SaliencLeftPallidum	-0.007	-0.019, 0.005	-1.16	0.248	0.998
SaliencBrainStem	0.001	-0.01, 0.013	0.24	0.813	0.998
SaliencLeftHippocampus	0.006	-0.005, 0.018	1.09	0.277	0.998
SaliencLeftAmygdala	-0.022	-0.039, -0.004	-2.45	0.015	0.998
SaliencLeftAccumbensArea	-0.012	-0.023, -0.001	-2.06	0.040	0.998
SaliencLeftVentraldc	0.007	-0.003, 0.017	1.33	0.184	0.998
SaliencRightCerebellumCortex	0.002	-0.008, 0.011	0.36	0.721	0.998
SaliencRightThalamusProper	0.004	-0.002, 0.011	1.29	0.199	0.998
SaliencRightCaudate	-0.010	-0.027, 0.007	-1.19	0.236	0.998
SaliencRightPutamen	0.003	-0.008, 0.014	0.59	0.555	0.998
SaliencRightPallidum	-0.002	-0.018, 0.015	-0.18	0.855	0.998
SaliencRightHippocampus	-0.006	-0.022, 0.011	-0.70	0.481	0.998
SaliencRightAmygdala	-0.003	-0.014, 0.007	-0.61	0.545	0.998
SaliencRightAccumbensArea	0.003	-0.009, 0.015	0.53	0.597	0.998
SaliencRightVentraldc	-0.008	-0.02, 0.003	-1.41	0.160	0.998
VentralAttentionLeftCerebellumCortex	-0.004	-0.015, 0.007	-0.80	0.427	0.998
VentralAttentionLeftThalamusProper	-0.006	-0.023, 0.01	-0.75	0.454	0.998
VentralAttentionLeftCaudate	-0.003	-0.013, 0.008	-0.50	0.616	0.998
VentralAttentionLeftPutamen	-0.001	-0.01, 0.008	-0.18	0.855	0.998
VentralAttentionLeftPallidum	0.004	-0.005, 0.013	0.83	0.406	0.998
VentralAttentionBrainStem	-0.001	-0.007, 0.006	-0.16	0.873	0.998
VentralAttentionLeftHippocampus	-0.015	-0.031, 0.001	-1.91	0.057	0.998
VentralAttentionLeftAmygdala	-0.002	-0.013, 0.008	-0.42	0.674	0.998
VentralAttentionLeftAccumbensArea	-0.009	-0.024, 0.006	-1.14	0.255	0.998
VentralAttentionLeftVentraldc	0.003	-0.013, 0.019	0.35	0.728	0.998
VentralAttentionRightCerebellumCortex	-0.003	-0.013, 0.007	-0.66	0.510	0.998
VentralAttentionRightThalamusProper	0.005	-0.007, 0.017	0.83	0.408	0.998
VentralAttentionRightCaudate	0.005	-0.006, 0.016	0.89	0.374	0.998
VentralAttentionRightPutamen	0.004	-0.006, 0.014	0.77	0.445	0.998
VentralAttentionRightPallidum	-0.004	-0.021, 0.012	-0.54	0.591	0.998
VentralAttentionRightHippocampus	0.003	-0.007, 0.014	0.62	0.538	0.998
VentralAttentionRightAmygdala	0.000	-0.009, 0.009	-0.02	0.981	0.998
VentralAttentionRightAccumbensArea	-0.001	-0.009, 0.008	-0.17	0.862	0.998
VentralAttentionRightVentraldc	0.003	-0.004, 0.009	0.81	0.421	0.998
VisualLeftCerebellumCortex	0.007	-0.009, 0.022	0.85	0.396	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
VisualLeftThalamusProper	-0.003	-0.013, 0.008	-0.51	0.608	0.998
VisualLeftCaudate	0.006	-0.009, 0.022	0.81	0.418	0.998
VisualLeftPutamen	0.004	-0.011, 0.018	0.48	0.631	0.998
VisualLeftPallidum	-0.001	-0.011, 0.009	-0.18	0.853	0.998
VisualBrainStem	-0.002	-0.014, 0.01	-0.36	0.716	0.998
VisualLeftHippocampus	0.010	-0.003, 0.023	1.58	0.116	0.998
VisualLeftAmygdala	0.008	-0.004, 0.019	1.27	0.204	0.998
VisualLeftAccumbensArea	-0.003	-0.021, 0.014	-0.38	0.701	0.998
VisualLeftVentraldc	-0.011	-0.021, -0.001	-2.14	0.033	0.998
VisualRightCerebellumCortex	-0.006	-0.016, 0.003	-1.31	0.190	0.998
VisualRightThalamusProper	-0.007	-0.016, 0.003	-1.45	0.148	0.998
VisualRightCaudate	0.001	-0.006, 0.007	0.20	0.845	0.998
VisualRightPutamen	0.002	-0.012, 0.017	0.33	0.742	0.998
VisualRightPallidum	0.000	-0.012, 0.011	-0.08	0.936	0.998
VisualRightHippocampus	0.002	-0.014, 0.019	0.28	0.780	0.998
VisualRightAmygdala	0.004	-0.013, 0.021	0.48	0.634	0.998
VisualRightAccumbensArea	0.010	0, 0.019	1.95	0.052	0.998
VisualRightVentraldc	0.004	-0.009, 0.017	0.62	0.536	0.998
WithinAuditory	0.001	-0.009, 0.011	0.25	0.802	0.998
WithinCinguloOpercula	-0.006	-0.015, 0.003	-1.38	0.170	0.998
WithinCinguloParietal	0.012	-0.015, 0.038	0.84	0.399	0.998
WithinDefault	-0.001	-0.009, 0.006	-0.33	0.745	0.998
WithinDorsalAttention	0.004	-0.004, 0.011	0.94	0.349	0.998
WithinFrontoParietal	0.004	-0.003, 0.012	1.08	0.281	0.998
WithinRetrosplenialTemporal	-0.002	-0.019, 0.015	-0.27	0.788	0.998
WithinSensorimotorHand	0.001	-0.009, 0.01	0.14	0.889	0.998
WithinSensorimotorMouth	0.008	-0.011, 0.028	0.84	0.399	0.998
WithinSalience	0.000	-0.017, 0.017	-0.01	0.994	0.998
WithinVentralAttention	0.004	-0.005, 0.012	0.86	0.393	0.998
WithinVisual	-0.013	-0.025, -0.002	-2.26	0.025	0.998
AuditoryCinguloOpercular	-0.001	-0.008, 0.006	-0.24	0.812	0.998
AuditoryCinguloParietal	0.003	-0.006, 0.012	0.59	0.555	0.998
AuditoryDefault	0.000	-0.006, 0.006	-0.08	0.938	0.998
AuditoryDorsalAttention	-0.002	-0.007, 0.003	-0.78	0.434	0.998
AuditoryFrontoParietal	0.001	-0.005, 0.007	0.32	0.751	0.998
AuditoryRetrosplenialTemporal	-0.001	-0.01, 0.007	-0.34	0.736	0.998
AuditorySensorimotorHand	0.002	-0.006, 0.009	0.40	0.690	0.998
AuditorySensorimotorMouth	-0.001	-0.012, 0.01	-0.20	0.841	0.998
AuditorySalience	0.000	-0.009, 0.01	0.07	0.945	0.998
AuditoryVentralAttention	0.002	-0.004, 0.009	0.71	0.477	0.998
AuditoryVisual	0.002	-0.006, 0.009	0.45	0.655	0.998
CinguloOpercularCinguloParietal	-0.001	-0.009, 0.008	-0.19	0.849	0.998
CinguloOpercularDefault	-0.002	-0.008, 0.005	-0.49	0.624	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
CinguloOpercularDorsalAttention	0.002	-0.003, 0.008	0.85	0.395	0.998
CinguloOpercularFrontoParietal	0.001	-0.005, 0.006	0.26	0.792	0.998
CinguloOpercularRetrosplenialTemporal	0.004	-0.005, 0.012	0.82	0.413	0.998
CinguloOpercularSensorimotorHand	0.004	-0.002, 0.01	1.37	0.171	0.998
CinguloOpercularSensorimotorMouth	0.000	-0.009, 0.009	0.00	0.998	0.998
CinguloOpercularSalience	-0.003	-0.011, 0.005	-0.66	0.508	0.998
CinguloOpercularVentralAttention	0.003	-0.004, 0.009	0.86	0.391	0.998
CinguloOpercularVisual	-0.004	-0.011, 0.004	-1.00	0.319	0.998
CinguloParietalDefault	0.001	-0.008, 0.011	0.23	0.821	0.998
CinguloParietalDorsalAttention	-0.001	-0.01, 0.008	-0.19	0.846	0.998
CinguloParietalFrontoParietal	-0.013	-0.022, -0.004	-2.95	0.003	0.998
CinguloParietalRetrosplenialTemporal	0.002	-0.011, 0.015	0.32	0.749	0.998
CinguloParietalSensorimotorHand	0.002	-0.008, 0.011	0.32	0.750	0.998
CinguloParietalSensorimotorMouth	0.003	-0.009, 0.016	0.54	0.587	0.998
CinguloParietalSalience	0.002	-0.011, 0.015	0.29	0.769	0.998
CinguloParietalVentralAttention	0.004	-0.004, 0.012	0.98	0.327	0.998
CinguloParietalVisual	-0.003	-0.014, 0.008	-0.52	0.604	0.998
DefaultDorsalAttention	0.003	-0.004, 0.009	0.87	0.387	0.998
DefaultFrontoParietal	0.003	-0.003, 0.008	0.90	0.371	0.998
DefaultRetrosplenialTemporal	-0.003	-0.011, 0.005	-0.63	0.531	0.998
DefaultSensorimotorHand	-0.004	-0.009, 0.002	-1.21	0.226	0.998
DefaultSensorimotorMouth	0.006	-0.002, 0.014	1.55	0.122	0.998
DefaultSalience	0.000	-0.009, 0.008	-0.10	0.921	0.998
DefaultVentralAttention	-0.001	-0.007, 0.006	-0.16	0.876	0.998
DefaultVisual	0.005	-0.002, 0.011	1.37	0.172	0.998
DorsalAttentionFrontoParietal	-0.002	-0.007, 0.004	-0.60	0.546	0.998
DorsalAttentionRetrosplenialTemporal	-0.001	-0.009, 0.006	-0.30	0.761	0.998
DorsalAttentionSensorimotorHand	0.004	-0.001, 0.009	1.42	0.156	0.998
DorsalAttentionSensorimotorMouth	-0.001	-0.008, 0.007	-0.20	0.840	0.998
DorsalAttentionSalience	0.001	-0.007, 0.008	0.14	0.885	0.998
DorsalAttentionVentralAttention	-0.001	-0.007, 0.005	-0.41	0.686	0.998
DorsalAttentionVisual	-0.003	-0.009, 0.004	-0.74	0.460	0.998
FrontoParietalRetrosplenialTemporal	-0.005	-0.012, 0.001	-1.64	0.101	0.998
FrontoParietalSensorimotorHand	0.000	-0.006, 0.006	-0.01	0.989	0.998
FrontoParietalSensorimotorMouth	0.002	-0.006, 0.009	0.50	0.617	0.998
FrontoParietalSalience	0.008	0.001, 0.016	2.18	0.030	0.998
FrontoParietalVentralAttention	0.002	-0.003, 0.007	0.65	0.519	0.998
FrontoParietalVisual	-0.003	-0.008, 0.003	-0.88	0.379	0.998
RetrosplenialTemporalSensorimotorHand	-0.005	-0.012, 0.003	-1.25	0.213	0.998
RetrosplenialTemporalSensorimotorMouth	-0.005	-0.015, 0.006	-0.87	0.385	0.998
RetrosplenialTemporalSalience	-0.003	-0.013, 0.007	-0.55	0.582	0.998
RetrosplenialTemporalVentralAttention	-0.003	-0.011, 0.005	-0.73	0.465	0.998

eTable 8. Differences in functional connectivity between the never-suicidal group and the child-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
RetrosplenialTemporalVisual	0.002	-0.008, 0.011	0.39	0.693	0.998
SensorimotorHandSensorimotorMouth	0.005	-0.005, 0.015	0.97	0.335	0.998
SensorimotorHandSalience	0.002	-0.006, 0.009	0.41	0.679	0.998
SensorimotorHandVentralAttention	0.000	-0.006, 0.006	0.00	0.997	0.998
SensorimotorHandVisual	0.004	-0.003, 0.011	1.21	0.229	0.998
SensorimotorMouthSalience	-0.008	-0.019, 0.003	-1.39	0.167	0.998
SensorimotorMouthVentralAttention	0.002	-0.005, 0.009	0.57	0.572	0.998
SensorimotorMouthVisual	-0.005	-0.015, 0.004	-1.07	0.286	0.998
SalienceVentralAttention	-0.001	-0.009, 0.007	-0.22	0.825	0.998
SalienceVisual	-0.008	-0.017, 0.001	-1.71	0.089	0.998
VentralAttentionVisual	0.000	-0.007, 0.006	-0.05	0.961	0.998

Bold denotes group differences at p<0.05 uncorrected

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
AuditoryLeftCerebellumCortex	-0.006	-0.019, 0.009	-0.77	0.444	0.885
AuditoryLeftThalamusProper	-0.003	-0.015, 0.01	-0.39	0.698	0.943
AuditoryLeftCaudate	-0.003	-0.021, 0.015	-0.30	0.761	0.956
AuditoryLeftPutamen	0.003	-0.009, 0.015	0.45	0.654	0.928
AuditoryLeftPallidum	-0.002	-0.013, 0.009	-0.32	0.748	0.956
AuditoryBrainStem	0.009	-0.003, 0.021	1.50	0.135	0.689
AuditoryLeftHippocampus	0.002	-0.005, 0.009	0.48	0.630	0.928
AuditoryLeftAmygdala	0.008	-0.006, 0.022	1.10	0.273	0.815
AuditoryLeftAccumbensArea	-0.016	-0.029, -0.003	-2.36	0.019	0.458
AuditoryLeftVentraldc	-0.002	-0.021, 0.016	-0.24	0.814	0.962
AuditoryRightCerebellumCortex	-0.011	-0.028, 0.006	-1.33	0.185	0.782
AuditoryRightThalamusProper	0.008	-0.003, 0.02	1.39	0.164	0.750
AuditoryRightCaudate	-0.001	-0.017, 0.014	-0.17	0.866	0.966
AuditoryRightPutamen	0.002	-0.013, 0.017	0.24	0.807	0.962
AuditoryRightPallidum	-0.006	-0.019, 0.007	-0.91	0.363	0.834
AuditoryRightHippocampus	-0.005	-0.024, 0.015	-0.50	0.618	0.928
AuditoryRightAmygdala	0.007	-0.005, 0.018	1.13	0.258	0.807
AuditoryRightAccumbensArea	0.003	-0.007, 0.013	0.63	0.528	0.928
AuditoryRightVentraldc	0.013	0.003, 0.023	2.55	0.011	0.458
CinguloOpercularLeftCerebellumCortex	0.000	-0.006, 0.007	0.05	0.958	0.994
CinguloOpercularLeftThalamusProper	-0.009	-0.025, 0.008	-1.04	0.299	0.815
CinguloOpercularLeftCaudate	-0.005	-0.019, 0.008	-0.75	0.451	0.885
CinguloOpercularLeftPutamen	-0.007	-0.026, 0.012	-0.70	0.483	0.895
CinguloOpercularLeftPallidum	-0.006	-0.023, 0.011	-0.74	0.459	0.885
CinguloOpercularBrainStem	0.002	-0.01, 0.014	0.31	0.759	0.956
CinguloOpercularLeftHippocampus	0.000	-0.014, 0.014	-0.04	0.965	0.994

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
CinguloOpercularLeftAmygdala	-0.004	-0.019, 0.01	-0.61	0.543	0.928
CinguloOpercularLeftAccumbensArea	-0.012	-0.024, 0.001	-1.85	0.065	0.570
CinguloOpercularLeftVentraldc	-0.001	-0.019, 0.018	-0.06	0.952	0.994
CinguloOpercularRightCerebellumCortex	0.006	-0.006, 0.017	0.95	0.341	0.815
CinguloOpercularRightThalamusProper	-0.012	-0.022, -0.001	-2.16	0.031	0.458
CinguloOpercularRightCaudate	0.008	-0.002, 0.017	1.55	0.122	0.653
CinguloOpercularRightPutamen	0.006	0, 0.012	1.84	0.066	0.570
CinguloOpercularRightPallidum	0.002	-0.014, 0.018	0.26	0.792	0.956
CinguloOpercularRightHippocampus	-0.002	-0.015, 0.011	-0.28	0.778	0.956
CinguloOpercularRightAmygdala	-0.008	-0.026, 0.01	-0.91	0.365	0.834
CinguloOpercularRightAccumbensArea	-0.003	-0.02, 0.014	-0.35	0.723	0.950
CinguloOpercularRightVentraldc	-0.006	-0.018, 0.006	-0.96	0.338	0.815
CinguloParietalLeftCerebellumCortex	-0.009	-0.023, 0.006	-1.20	0.233	0.782
CinguloParietalLeftThalamusProper	0.001	-0.014, 0.015	0.10	0.922	0.990
CinguloParietalLeftCaudate	-0.002	-0.014, 0.01	-0.38	0.706	0.943
CinguloParietalLeftPutamen	-0.018	-0.036, 0	-1.96	0.051	0.559
CinguloParietalLeftPallidum	0.005	-0.007, 0.016	0.76	0.449	0.885
CinguloParietalBrainStem	0.001	-0.009, 0.011	0.21	0.833	0.963
CinguloParietalLeftHippocampus	0.017	0.007, 0.027	3.45	0.001	0.194
CinguloParietalLeftAmygdala	0.002	-0.004, 0.009	0.75	0.452	0.885
CinguloParietalLeftAccumbensArea	-0.006	-0.023, 0.011	-0.68	0.497	0.900
CinguloParietalLeftVentraldc	-0.001	-0.014, 0.012	-0.17	0.864	0.966
CinguloParietalRightCerebellumCortex	-0.006	-0.024, 0.012	-0.64	0.520	0.925
CinguloParietalRightThalamusProper	0.011	-0.006, 0.029	1.28	0.200	0.782
CinguloParietalRightCaudate	-0.001	-0.013, 0.011	-0.16	0.871	0.966
CinguloParietalRightPutamen	-0.004	-0.018, 0.01	-0.55	0.580	0.928
CinguloParietalRightPallidum	0.002	-0.012, 0.016	0.28	0.782	0.956
CinguloParietalRightHippocampus	-0.005	-0.017, 0.007	-0.78	0.438	0.885
CinguloParietalRightAmygdala	0.000	-0.017, 0.018	0.01	0.989	0.996
CinguloParietalRightAccumbensArea	0.016	0.005, 0.027	2.92	0.004	0.443
CinguloParietalRightVentraldc	-0.001	-0.011, 0.008	-0.26	0.793	0.956
DefaultLeftCerebellumCortex	0.009	-0.001, 0.019	1.84	0.067	0.570
DefaultLeftThalamusProper	0.008	0.002, 0.014	2.43	0.016	0.458
DefaultLeftCaudate	0.011	-0.005, 0.026	1.39	0.167	0.750
DefaultLeftPutamen	-0.003	-0.015, 0.01	-0.42	0.673	0.932
DefaultLeftPallidum	-0.022	-0.039, -0.005	-2.54	0.011	0.458
DefaultBrainStem	0.013	-0.003, 0.03	1.58	0.116	0.653
DefaultLeftHippocampus	0.010	-0.001, 0.02	1.87	0.063	0.570
DefaultLeftAmygdala	-0.018	-0.031, -0.005	-2.71	0.007	0.458
DefaultLeftAccumbensArea	0.005	-0.009, 0.019	0.70	0.484	0.895
DefaultLeftVentraldc	0.008	-0.003, 0.019	1.42	0.156	0.748
DefaultRightCerebellumCortex	-0.012	-0.031, 0.006	-1.30	0.196	0.782
DefaultRightThalamusProper	0.007	-0.004, 0.018	1.27	0.207	0.782

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
DefaultRightCaudate	-0.007	-0.017, 0.003	-1.44	0.150	0.748
DefaultRightPutamen	0.012	0.001, 0.022	2.21	0.028	0.458
DefaultRightPallidum	0.006	-0.002, 0.013	1.51	0.133	0.688
DefaultRightHippocampus	-0.023	-0.039, -0.007	-2.87	0.004	0.443
DefaultRightAmygdala	0.003	-0.009, 0.016	0.52	0.605	0.928
DefaultRightAccumbensArea	0.007	-0.012, 0.026	0.75	0.454	0.885
DefaultRightVentraldc	-0.002	-0.019, 0.016	-0.18	0.859	0.966
DorsalAttentionLeftCerebellumCortex	0.015	0.003, 0.026	2.41	0.016	0.458
DorsalAttentionLeftThalamusProper	-0.015	-0.029, -0.002	-2.19	0.029	0.458
DorsalAttentionLeftCaudate	0.004	-0.009, 0.016	0.57	0.572	0.928
DorsalAttentionLeftPutamen	0.011	-0.003, 0.025	1.58	0.116	0.653
DorsalAttentionLeftPallidum	0.000	-0.02, 0.019	-0.04	0.968	0.994
DorsalAttentionBrainStem	-0.003	-0.015, 0.009	-0.48	0.631	0.928
DorsalAttentionLeftHippocampus	0.002	-0.01, 0.013	0.27	0.787	0.956
DorsalAttentionLeftAmygdala	0.008	-0.002, 0.018	1.57	0.118	0.653
DorsalAttentionLeftAccumbensArea	0.000	-0.007, 0.007	0.01	0.990	0.996
DorsalAttentionLeftVentraldc	-0.011	-0.028, 0.006	-1.27	0.203	0.782
DorsalAttentionRightCerebellumCortex	0.005	-0.007, 0.017	0.77	0.442	0.885
DorsalAttentionRightThalamusProper	0.008	-0.009, 0.025	0.97	0.331	0.815
DorsalAttentionRightCaudate	-0.009	-0.027, 0.008	-1.05	0.296	0.815
DorsalAttentionRightPutamen	0.003	-0.008, 0.015	0.58	0.562	0.928
DorsalAttentionRightPallidum	-0.012	-0.026, 0.001	-1.77	0.078	0.596
DorsalAttentionRightHippocampus	0.002	-0.01, 0.015	0.37	0.715	0.948
DorsalAttentionRightAmygdala	0.003	-0.008, 0.015	0.54	0.593	0.928
DorsalAttentionRightAccumbensArea	-0.007	-0.024, 0.011	-0.74	0.460	0.885
DorsalAttentionRightVentraldc	0.005	-0.006, 0.016	0.89	0.374	0.834
FrontoParietalLeftCerebellumCortex	-0.008	-0.018, 0.003	-1.43	0.153	0.748
FrontoParietalLeftThalamusProper	0.005	-0.005, 0.014	0.95	0.340	0.815
FrontoParietalLeftCaudate	0.001	-0.006, 0.008	0.30	0.765	0.956
FrontoParietalLeftPutamen	-0.011	-0.026, 0.004	-1.42	0.156	0.748
FrontoParietalLeftPallidum	-0.001	-0.013, 0.01	-0.20	0.845	0.966
FrontoParietalBrainStem	-0.002	-0.018, 0.015	-0.19	0.852	0.966
FrontoParietalLeftHippocampus	-0.007	-0.024, 0.01	-0.78	0.434	0.885
FrontoParietalLeftAmygdala	0.010	-0.001, 0.021	1.75	0.082	0.610
FrontoParietalLeftAccumbensArea	-0.007	-0.021, 0.006	-1.06	0.292	0.815
FrontoParietalLeftVentraldc	0.000	-0.012, 0.012	0.00	0.996	0.996
FrontoParietalRightCerebellumCortex	0.003	-0.009, 0.014	0.43	0.664	0.928
FrontoParietalRightThalamusProper	0.001	-0.017, 0.018	0.09	0.930	0.991
FrontoParietalRightCaudate	-0.003	-0.014, 0.008	-0.56	0.574	0.928
FrontoParietalRightPutamen	-0.006	-0.016, 0.004	-1.13	0.261	0.807
FrontoParietalRightPallidum	0.008	-0.002, 0.017	1.64	0.101	0.653
FrontoParietalRightHippocampus	0.001	-0.006, 0.007	0.17	0.866	0.966

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
FrontoParietalRightAmygdala	0.000	-0.016, 0.015	-0.01	0.990	0.996
FrontoParietalRightAccumbensArea	0.000	-0.011, 0.011	-0.01	0.990	0.996
FrontoParietalRightVentraldc	-0.006	-0.021, 0.01	-0.75	0.453	0.885
RetrosplenialTemporalLeftCerebellumCortex	0.007	-0.005, 0.02	1.14	0.255	0.807
RetrosplenialTemporalLeftThalamusProper	0.000	-0.011, 0.011	-0.01	0.992	0.996
RetrosplenialTemporalLeftCaudate	0.012	0.001, 0.024	2.16	0.031	0.458
RetrosplenialTemporalLeftPutamen	-0.004	-0.01, 0.003	-1.02	0.308	0.815
RetrosplenialTemporalLeftPallidum	0.008	-0.008, 0.023	0.95	0.344	0.816
RetrosplenialTemporalBrainStem	-0.011	-0.023, 0.002	-1.63	0.103	0.653
RetrosplenialTemporalLeftHippocampus	-0.002	-0.02, 0.015	-0.26	0.794	0.956
RetrosplenialTemporalLeftAmygdala	0.003	-0.015, 0.021	0.36	0.720	0.950
RetrosplenialTemporalLeftAccumbensArea	0.008	-0.004, 0.019	1.25	0.211	0.782
RetrosplenialTemporalLeftVentraldc	0.002	-0.013, 0.018	0.30	0.762	0.956
RetrosplenialTemporalRightCerebellumCortex	-0.005	-0.021, 0.01	-0.70	0.486	0.895
RetrosplenialTemporalRightThalamusProper	-0.004	-0.017, 0.008	-0.69	0.493	0.900
RetrosplenialTemporalRightCaudate	0.003	-0.017, 0.022	0.28	0.778	0.956
RetrosplenialTemporalRightPutamen	0.007	-0.004, 0.017	1.17	0.242	0.789
RetrosplenialTemporalRightPallidum	-0.002	-0.012, 0.009	-0.29	0.770	0.956
RetrosplenialTemporalRightHippocampus	0.006	-0.004, 0.016	1.20	0.229	0.782
RetrosplenialTemporalRightAmygdala	0.007	0.001, 0.013	2.21	0.028	0.458
RetrosplenialTemporalRightAccumbensArea	-0.010	-0.025, 0.006	-1.21	0.229	0.782
RetrosplenialTemporalRightVentraldc	0.000	-0.015, 0.014	-0.06	0.951	0.994
SensorimotorHandLeftCerebellumCortex	-0.003	-0.022, 0.015	-0.36	0.716	0.948
SensorimotorHandLeftThalamusProper	-0.001	-0.018, 0.016	-0.14	0.888	0.977
SensorimotorHandLeftCaudate	0.003	-0.009, 0.015	0.48	0.632	0.928
SensorimotorHandLeftPutamen	-0.005	-0.019, 0.009	-0.65	0.515	0.925
SensorimotorHandLeftPallidum	-0.004	-0.019, 0.012	-0.47	0.642	0.928
SensorimotorHandBrainStem	-0.003	-0.015, 0.009	-0.53	0.598	0.928
SensorimotorHandLeftHippocampus	0.000	-0.018, 0.019	0.02	0.981	0.996
SensorimotorHandLeftAmygdala	0.005	-0.006, 0.017	0.90	0.370	0.834
SensorimotorHandLeftAccumbensArea	0.005	-0.005, 0.016	0.99	0.322	0.815
SensorimotorHandLeftVentraldc	0.010	0, 0.02	1.92	0.056	0.569
SensorimotorHandRightCerebellumCortex	0.004	-0.003, 0.01	1.10	0.272	0.815
SensorimotorHandRightThalamusProper	-0.008	-0.023, 0.008	-0.99	0.322	0.815
SensorimotorHandRightCaudate	-0.001	-0.015, 0.013	-0.14	0.892	0.977
SensorimotorHandRightPutamen	-0.015	-0.032, 0.003	-1.68	0.095	0.653
SensorimotorHandRightPallidum	0.006	-0.01, 0.022	0.75	0.453	0.885
SensorimotorHandRightHippocampus	-0.003	-0.015, 0.009	-0.44	0.661	0.928
SensorimotorHandRightAmygdala	-0.002	-0.016, 0.011	-0.34	0.733	0.956
SensorimotorHandRightAccumbensArea	0.002	-0.013, 0.016	0.23	0.822	0.963
SensorimotorHandRightVentraldc	0.010	-0.002, 0.021	1.59	0.112	0.653
SensorimotorMouthLeftCerebellumCortex	-0.004	-0.022, 0.014	-0.47	0.641	0.928
SensorimotorMouthLeftThalamusProper	0.011	-0.001, 0.023	1.81	0.072	0.595

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
SensorimotorMouthLeftCaudate	0.003	-0.007, 0.013	0.53	0.595	0.928
SensorimotorMouthLeftPutamen	0.012	0.002, 0.022	2.36	0.019	0.458
SensorimotorMouthLeftPallidum	0.007	0, 0.013	2.01	0.045	0.528
SensorimotorMouthBrainStem	-0.010	-0.025, 0.006	-1.25	0.214	0.782
SensorimotorMouthLeftHippocampus	-0.001	-0.014, 0.013	-0.13	0.894	0.977
SensorimotorMouthLeftAmygdala	-0.021	-0.039, -0.003	-2.25	0.025	0.458
SensorimotorMouthLeftAccumbensArea	0.023	0.006, 0.041	2.57	0.011	0.458
SensorimotorMouthLeftVentraldc	0.007	-0.006, 0.019	1.00	0.319	0.815
SensorimotorMouthRightCerebellumCortex	-0.014	-0.028, -0.001	-2.07	0.039	0.512
SensorimotorMouthRightThalamusProper	0.004	-0.01, 0.017	0.50	0.617	0.928
SensorimotorMouthRightCaudate	0.006	-0.006, 0.018	0.97	0.335	0.815
SensorimotorMouthRightPutamen	0.008	-0.009, 0.024	0.90	0.371	0.834
SensorimotorMouthRightPallidum	0.010	0, 0.021	1.91	0.058	0.569
SensorimotorMouthRightHippocampus	0.005	-0.005, 0.015	0.99	0.323	0.815
SensorimotorMouthRightAmygdala	0.006	-0.004, 0.015	1.23	0.220	0.782
SensorimotorMouthRightAccumbensArea	0.003	-0.004, 0.009	0.88	0.379	0.835
SensorimotorMouthRightVentraldc	0.007	-0.008, 0.022	0.96	0.338	0.815
SaliencyLeftCerebellumCortex	0.002	-0.011, 0.015	0.28	0.783	0.956
SaliencyLeftThalamusProper	-0.005	-0.022, 0.011	-0.61	0.540	0.928
SaliencyLeftCaudate	0.018	0.001, 0.035	2.04	0.042	0.512
SaliencyLeftPutamen	0.007	-0.004, 0.018	1.29	0.197	0.782
SaliencyLeftPallidum	-0.001	-0.013, 0.012	-0.11	0.913	0.983
SaliencyBrainStem	0.014	0.001, 0.027	2.09	0.038	0.512
SaliencyLeftHippocampus	0.003	-0.009, 0.015	0.49	0.626	0.928
SaliencyLeftAmygdala	-0.002	-0.02, 0.016	-0.21	0.830	0.963
SaliencyLeftAccumbensArea	-0.006	-0.018, 0.006	-1.00	0.317	0.815
SaliencyLeftVentraldc	-0.007	-0.019, 0.004	-1.26	0.208	0.782
SaliencyRightCerebellumCortex	0.004	-0.007, 0.014	0.70	0.484	0.895
SaliencyRightThalamusProper	-0.001	-0.008, 0.006	-0.27	0.791	0.956
SaliencyRightCaudate	0.002	-0.015, 0.018	0.21	0.834	0.963
SaliencyRightPutamen	0.003	-0.008, 0.015	0.60	0.551	0.928
SaliencyRightPallidum	0.010	-0.007, 0.026	1.19	0.235	0.782
SaliencyRightHippocampus	-0.015	-0.031, 0.002	-1.72	0.086	0.627
SaliencyRightAmygdala	0.004	-0.006, 0.015	0.79	0.431	0.885
SaliencyRightAccumbensArea	0.008	-0.005, 0.021	1.23	0.220	0.782
SaliencyRightVentraldc	-0.003	-0.015, 0.009	-0.45	0.651	0.928
VentralAttentionLeftCerebellumCortex	0.008	-0.004, 0.02	1.24	0.214	0.782
VentralAttentionLeftThalamusProper	-0.012	-0.03, 0.007	-1.26	0.210	0.782
VentralAttentionLeftCaudate	-0.008	-0.019, 0.003	-1.39	0.165	0.750
VentralAttentionLeftPutamen	-0.002	-0.012, 0.008	-0.41	0.684	0.942
VentralAttentionLeftPallidum	0.004	-0.006, 0.014	0.78	0.438	0.885
VentralAttentionBrainStem	-0.002	-0.009, 0.005	-0.53	0.597	0.928

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
VentralAttentionLeftHippocampus	-0.019	-0.034, -0.003	-2.34	0.020	0.458
VentralAttentionLeftAmygdala	0.000	-0.012, 0.012	0.05	0.958	0.994
VentralAttentionLeftAccumbensArea	0.004	-0.013, 0.02	0.46	0.647	0.928
VentralAttentionLeftVentraldc	0.009	-0.008, 0.026	1.05	0.295	0.815
VentralAttentionRightCerebellumCortex	0.000	-0.011, 0.011	0.01	0.994	0.996
VentralAttentionRightThalamusProper	-0.005	-0.018, 0.008	-0.72	0.474	0.895
VentralAttentionRightCaudate	0.003	-0.01, 0.015	0.42	0.673	0.932
VentralAttentionRightPutamen	0.003	-0.008, 0.014	0.56	0.577	0.928
VentralAttentionRightPallidum	0.009	-0.009, 0.026	0.99	0.325	0.815
VentralAttentionRightHippocampus	0.005	-0.007, 0.016	0.80	0.425	0.885
VentralAttentionRightAmygdala	-0.006	-0.016, 0.004	-1.19	0.235	0.782
VentralAttentionRightAccumbensArea	0.001	-0.008, 0.01	0.22	0.827	0.963
VentralAttentionRightVentraldc	-0.005	-0.011, 0.002	-1.35	0.177	0.782
VisualLeftCerebellumCortex	0.004	-0.011, 0.019	0.51	0.608	0.928
VisualLeftThalamusProper	0.008	-0.002, 0.019	1.59	0.113	0.653
VisualLeftCaudate	-0.002	-0.018, 0.015	-0.18	0.855	0.966
VisualLeftPutamen	0.010	-0.008, 0.028	1.06	0.292	0.815
VisualLeftPallidum	0.002	-0.008, 0.013	0.45	0.652	0.928
VisualBrainStem	-0.005	-0.018, 0.009	-0.68	0.497	0.900
VisualLeftHippocampus	-0.009	-0.022, 0.004	-1.40	0.162	0.750
VisualLeftAmygdala	-0.002	-0.015, 0.01	-0.34	0.735	0.956
VisualLeftAccumbensArea	-0.003	-0.02, 0.015	-0.28	0.784	0.956
VisualLeftVentraldc	-0.005	-0.016, 0.007	-0.77	0.443	0.885
VisualRightCerebellumCortex	-0.005	-0.014, 0.005	-0.96	0.339	0.815
VisualRightThalamusProper	0.005	-0.005, 0.015	1.01	0.312	0.815
VisualRightCaudate	0.003	-0.004, 0.009	0.82	0.412	0.885
VisualRightPutamen	-0.009	-0.025, 0.006	-1.16	0.248	0.797
VisualRightPallidum	0.006	-0.006, 0.018	0.96	0.339	0.815
VisualRightHippocampus	-0.014	-0.032, 0.003	-1.59	0.114	0.653
VisualRightAmygdala	-0.015	-0.032, 0.002	-1.69	0.091	0.649
VisualRightAccumbensArea	0.003	-0.008, 0.014	0.52	0.602	0.928
VisualRightVentraldc	0.007	-0.008, 0.022	0.96	0.335	0.815
WithinAuditory	0.003	-0.007, 0.013	0.65	0.517	0.925
WithinCinguloOpercula	0.003	-0.006, 0.013	0.73	0.464	0.887
WithinCinguloParietal	0.008	-0.019, 0.036	0.61	0.542	0.928
WithinDefault	-0.002	-0.01, 0.006	-0.47	0.639	0.928
WithinDorsalAttention	-0.002	-0.01, 0.006	-0.45	0.654	0.928
WithinFrontoParietal	0.002	-0.006, 0.01	0.57	0.569	0.928
WithinRetrosplenialTemporal	-0.003	-0.02, 0.014	-0.39	0.696	0.943
WithinSensorimotorHand	0.006	-0.003, 0.015	1.28	0.201	0.782
WithinSensorimotorMouth	-0.003	-0.023, 0.017	-0.33	0.743	0.956
WithinSalience	0.002	-0.016, 0.019	0.16	0.870	0.966
WithinVentralAttention	-0.006	-0.014, 0.002	-1.52	0.130	0.685

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
WithinVisual	0.008	-0.005, 0.02	1.22	0.223	0.782
AuditoryCinguloOpercular	0.006	-0.001, 0.014	1.61	0.109	0.653
AuditoryCinguloParietal	0.001	-0.009, 0.011	0.19	0.850	0.966
AuditoryDefault	-0.002	-0.008, 0.004	-0.61	0.540	0.928
AuditoryDorsalAttention	0.004	-0.002, 0.009	1.17	0.242	0.789
AuditoryFrontoParietal	0.003	-0.003, 0.009	0.94	0.347	0.816
AuditoryRetrosplenialTemporal	0.000	-0.009, 0.009	-0.04	0.968	0.994
AuditorySensorimotorHand	0.006	-0.002, 0.013	1.55	0.122	0.653
AuditorySensorimotorMouth	0.002	-0.009, 0.013	0.38	0.704	0.943
AuditorySalience	-0.001	-0.011, 0.009	-0.21	0.831	0.963
AuditoryVentralAttention	0.003	-0.004, 0.011	0.96	0.339	0.815
AuditoryVisual	0.002	-0.007, 0.01	0.40	0.690	0.943
CinguloOpercularCinguloParietal	-0.002	-0.012, 0.007	-0.48	0.631	0.928
CinguloOpercularDefault	0.002	-0.005, 0.009	0.51	0.610	0.928
CinguloOpercularDorsalAttention	0.006	0, 0.012	1.94	0.053	0.559
CinguloOpercularFrontoParietal	0.002	-0.004, 0.008	0.78	0.438	0.885
CinguloOpercularRetrosplenialTemporal	0.002	-0.007, 0.011	0.43	0.664	0.928
CinguloOpercularSensorimotorHand	0.008	0.001, 0.014	2.21	0.028	0.458
CinguloOpercularSensorimotorMouth	0.004	-0.005, 0.013	0.88	0.377	0.835
CinguloOpercularSalience	0.001	-0.008, 0.011	0.25	0.804	0.962
CinguloOpercularVentralAttention	0.001	-0.006, 0.008	0.32	0.747	0.956
CinguloOpercularVisual	-0.004	-0.012, 0.004	-1.00	0.320	0.815
CinguloParietalDefault	0.001	-0.009, 0.01	0.13	0.900	0.979
CinguloParietalDorsalAttention	0.001	-0.01, 0.011	0.12	0.906	0.980
CinguloParietalFrontoParietal	-0.009	-0.018, 0.001	-1.86	0.063	0.570
CinguloParietalRetrosplenialTemporal	-0.003	-0.017, 0.011	-0.47	0.642	0.928
CinguloParietalSensorimotorHand	-0.003	-0.013, 0.007	-0.59	0.553	0.928
CinguloParietalSensorimotorMouth	0.002	-0.012, 0.016	0.30	0.763	0.956
CinguloParietalSalience	-0.002	-0.015, 0.012	-0.24	0.812	0.962
CinguloParietalVentralAttention	-0.004	-0.014, 0.005	-0.86	0.390	0.853
CinguloParietalVisual	0.003	-0.01, 0.015	0.46	0.644	0.928
DefaultDorsalAttention	0.004	-0.003, 0.011	1.09	0.277	0.815
DefaultFrontoParietal	0.003	-0.002, 0.009	1.20	0.229	0.782
DefaultRetrosplenialTemporal	-0.002	-0.01, 0.005	-0.61	0.543	0.928
DefaultSensorimotorHand	-0.002	-0.009, 0.004	-0.76	0.450	0.885
DefaultSensorimotorMouth	-0.001	-0.009, 0.007	-0.24	0.812	0.962
DefaultSalience	0.004	-0.005, 0.013	0.97	0.335	0.815
DefaultVentralAttention	0.001	-0.005, 0.007	0.40	0.687	0.943
DefaultVisual	-0.006	-0.013, 0.001	-1.77	0.077	0.596
DorsalAttentionFrontoParietal	-0.003	-0.008, 0.003	-0.94	0.350	0.817
DorsalAttentionRetrosplenialTemporal	0.000	-0.007, 0.008	0.09	0.926	0.991
DorsalAttentionSensorimotorHand	0.004	-0.002, 0.009	1.26	0.208	0.782

eTable 9. Differences in functional connectivity between the never-suicidal group and the caregiver-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
DorsalAttentionSensorimotorMouth	0.008	0, 0.016	2.05	0.041	0.512
DorsalAttentionSalience	-0.001	-0.009, 0.008	-0.19	0.851	0.966
DorsalAttentionVentralAttention	0.002	-0.004, 0.008	0.64	0.525	0.928
DorsalAttentionVisual	-0.002	-0.009, 0.005	-0.57	0.567	0.928
FrontoParietalRetrosplenialTemporal	0.000	-0.007, 0.007	-0.08	0.936	0.991
FrontoParietalSensorimotorHand	-0.003	-0.009, 0.003	-0.89	0.372	0.834
FrontoParietalSensorimotorMouth	0.000	-0.007, 0.008	0.08	0.934	0.991
FrontoParietalSalience	0.007	-0.001, 0.015	1.79	0.074	0.596
FrontoParietalVentralAttention	0.001	-0.004, 0.007	0.52	0.605	0.928
FrontoParietalVisual	-0.008	-0.014, -0.001	-2.43	0.016	0.458
RetrosplenialTemporalSensorimotorHand	0.001	-0.007, 0.008	0.15	0.880	0.972
RetrosplenialTemporalSensorimotorMouth	0.002	-0.009, 0.013	0.38	0.701	0.943
RetrosplenialTemporalSalience	-0.014	-0.026, -0.002	-2.21	0.028	0.458
RetrosplenialTemporalVentralAttention	0.000	-0.009, 0.008	-0.04	0.966	0.994
RetrosplenialTemporalVisual	0.000	-0.01, 0.01	-0.06	0.949	0.994
SensorimotorHandSensorimotorMouth	0.011	0, 0.023	1.95	0.052	0.559
SensorimotorHandSalience	0.002	-0.006, 0.011	0.48	0.632	0.928
SensorimotorHandVentralAttention	0.003	-0.003, 0.009	1.06	0.288	0.815
SensorimotorHandVisual	0.006	-0.001, 0.014	1.61	0.109	0.653
SensorimotorMouthSalience	-0.007	-0.018, 0.004	-1.25	0.213	0.782
SensorimotorMouthVentralAttention	-0.001	-0.008, 0.007	-0.12	0.902	0.979
SensorimotorMouthVisual	-0.002	-0.013, 0.008	-0.44	0.661	0.928
SalienceVentralAttention	0.007	-0.002, 0.016	1.56	0.119	0.653
SalienceVisual	-0.006	-0.015, 0.004	-1.13	0.261	0.807
VentralAttentionVisual	-0.003	-0.01, 0.005	-0.72	0.469	0.892

Bold denotes group differences at p<0.05 uncorrected

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
AuditoryLeftCerebellumCortex	0.020	-0.003, 0.042	1.70	0.092	0.992
AuditoryLeftThalamusProper	0.004	-0.019, 0.027	0.35	0.729	0.992
AuditoryLeftCaudate	-0.017	-0.049, 0.016	-1.00	0.319	0.992
AuditoryLeftPutamen	-0.004	-0.027, 0.019	-0.36	0.720	0.992
AuditoryLeftPallidum	-0.009	-0.027, 0.008	-1.05	0.296	0.992
AuditoryBrainStem	-0.002	-0.024, 0.02	-0.17	0.865	0.992
AuditoryLeftHippocampus	0.007	-0.005, 0.019	1.21	0.228	0.992
AuditoryLeftAmygdala	0.006	-0.019, 0.031	0.50	0.620	0.992
AuditoryLeftAccumbensArea	-0.002	-0.025, 0.02	-0.21	0.831	0.992
AuditoryLeftVentraldc	-0.003	-0.032, 0.027	-0.18	0.860	0.992

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
AuditoryRightCerebellumCortex	0.004	-0.028, 0.036	0.25	0.801	0.992
AuditoryRightThalamusProper	0.011	-0.011, 0.032	0.97	0.334	0.992
AuditoryRightCaudate	-0.002	-0.029, 0.025	-0.15	0.878	0.992
AuditoryRightPutamen	-0.002	-0.026, 0.022	-0.17	0.867	0.992
AuditoryRightPallidum	0.000	-0.024, 0.025	0.03	0.977	0.992
AuditoryRightHippocampus	0.024	-0.009, 0.057	1.44	0.153	0.992
AuditoryRightAmygdala	0.003	-0.016, 0.022	0.33	0.743	0.992
AuditoryRightAccumbensArea	0.010	-0.008, 0.028	1.09	0.278	0.992
AuditoryRightVentraldc	-0.001	-0.019, 0.018	-0.05	0.961	0.992
CinguloOpercularLeftCerebellumCortex	0.004	-0.006, 0.014	0.82	0.414	0.992
CinguloOpercularLeftThalamusProper	-0.022	-0.05, 0.006	-1.54	0.127	0.992
CinguloOpercularLeftCaudate	-0.003	-0.029, 0.023	-0.22	0.823	0.992
CinguloOpercularLeftPutamen	-0.019	-0.054, 0.016	-1.08	0.284	0.992
CinguloOpercularLeftPallidum	-0.015	-0.042, 0.012	-1.13	0.263	0.992
CinguloOpercularBrainStem	0.012	-0.009, 0.034	1.12	0.267	0.992
CinguloOpercularLeftHippocampus	0.001	-0.025, 0.026	0.05	0.958	0.992
CinguloOpercularLeftAmygdala	0.021	-0.002, 0.043	1.86	0.067	0.992
CinguloOpercularLeftAccumbensArea	-0.007	-0.029, 0.016	-0.60	0.553	0.992
CinguloOpercularLeftVentraldc	0.001	-0.03, 0.032	0.04	0.965	0.992
CinguloOpercularRightCerebellumCortex	0.001	-0.019, 0.022	0.14	0.887	0.992
CinguloOpercularRightThalamusProper	-0.004	-0.022, 0.013	-0.49	0.629	0.992
CinguloOpercularRightCaudate	-0.006	-0.024, 0.013	-0.59	0.559	0.992
CinguloOpercularRightPutamen	-0.001	-0.012, 0.01	-0.21	0.838	0.992
CinguloOpercularRightPallidum	-0.001	-0.028, 0.026	-0.08	0.940	0.992
CinguloOpercularRightHippocampus	0.010	-0.012, 0.031	0.89	0.377	0.992
CinguloOpercularRightAmygdala	0.012	-0.019, 0.042	0.77	0.446	0.992
CinguloOpercularRightAccumbensArea	0.003	-0.025, 0.031	0.20	0.842	0.992
CinguloOpercularRightVentraldc	0.011	-0.011, 0.033	0.99	0.325	0.992
CinguloParietalLeftCerebellumCortex	-0.012	-0.037, 0.014	-0.90	0.370	0.992
CinguloParietalLeftThalamusProper	0.018	-0.009, 0.044	1.33	0.186	0.992
CinguloParietalLeftCaudate	0.013	-0.009, 0.035	1.14	0.259	0.992
CinguloParietalLeftPutamen	0.002	-0.024, 0.027	0.13	0.900	0.992
CinguloParietalLeftPallidum	-0.007	-0.027, 0.013	-0.68	0.497	0.992
CinguloParietalBrainStem	0.008	-0.008, 0.024	1.02	0.311	0.992
CinguloParietalLeftHippocampus	0.016	-0.001, 0.033	1.92	0.058	0.992
CinguloParietalLeftAmygdala	-0.002	-0.012, 0.008	-0.40	0.691	0.992
CinguloParietalLeftAccumbensArea	-0.024	-0.051, 0.003	-1.74	0.085	0.992
CinguloParietalLeftVentraldc	0.000	-0.024, 0.024	0.01	0.995	0.995
CinguloParietalRightCerebellumCortex	0.001	-0.031, 0.033	0.08	0.934	0.992
CinguloParietalRightThalamusProper	0.009	-0.02, 0.038	0.63	0.530	0.992
CinguloParietalRightCaudate	0.002	-0.02, 0.024	0.20	0.842	0.992
CinguloParietalRightPutamen	-0.003	-0.028, 0.021	-0.26	0.797	0.992

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
CinguloParietalRightPallidum	0.010	-0.015, 0.034	0.77	0.443	0.992
CinguloParietalRightHippocampus	-0.002	-0.023, 0.02	-0.13	0.893	0.992
CinguloParietalRightAmygdala	-0.005	-0.033, 0.023	-0.36	0.722	0.992
CinguloParietalRightAccumbensArea	0.001	-0.018, 0.019	0.11	0.916	0.992
CinguloParietalRightVentraldc	-0.001	-0.017, 0.015	-0.12	0.904	0.992
DefaultLeftCerebellumCortex	0.003	-0.015, 0.02	0.29	0.774	0.992
DefaultLeftThalamusProper	0.009	-0.001, 0.02	1.77	0.081	0.992
DefaultLeftCaudate	0.018	-0.007, 0.043	1.45	0.150	0.992
DefaultLeftPutamen	0.001	-0.018, 0.021	0.14	0.888	0.992
DefaultLeftPallidum	-0.004	-0.03, 0.023	-0.28	0.777	0.992
DefaultBrainStem	0.011	-0.019, 0.042	0.75	0.454	0.992
DefaultLeftHippocampus	0.012	-0.007, 0.031	1.29	0.202	0.992
DefaultLeftAmygdala	-0.007	-0.03, 0.017	-0.59	0.560	0.992
DefaultLeftAccumbensArea	0.006	-0.016, 0.028	0.53	0.600	0.992
DefaultLeftVentraldc	0.005	-0.013, 0.024	0.57	0.570	0.992
DefaultRightCerebellumCortex	0.012	-0.02, 0.043	0.73	0.465	0.992
DefaultRightThalamusProper	0.011	-0.009, 0.03	1.09	0.277	0.992
DefaultRightCaudate	-0.025	-0.042, -0.008	-2.91	0.005	0.711
DefaultRightPutamen	0.002	-0.017, 0.022	0.25	0.804	0.992
DefaultRightPallidum	0.002	-0.012, 0.015	0.24	0.812	0.992
DefaultRightHippocampus	-0.032	-0.061, -0.003	-2.20	0.030	0.992
DefaultRightAmygdala	-0.004	-0.023, 0.016	-0.38	0.703	0.992
DefaultRightAccumbensArea	0.011	-0.016, 0.038	0.80	0.423	0.992
DefaultRightVentraldc	-0.007	-0.036, 0.022	-0.47	0.642	0.992
DorsalAttentionLeftCerebellumCortex	0.025	0.005, 0.044	2.50	0.014	0.871
DorsalAttentionLeftThalamusProper	-0.006	-0.03, 0.017	-0.53	0.600	0.992
DorsalAttentionLeftCaudate	0.004	-0.017, 0.025	0.37	0.711	0.992
DorsalAttentionLeftPutamen	0.000	-0.024, 0.025	0.03	0.978	0.992
DorsalAttentionLeftPallidum	-0.016	-0.049, 0.017	-0.96	0.338	0.992
DorsalAttentionBrainStem	-0.023	-0.043, -0.003	-2.24	0.028	0.992
DorsalAttentionLeftHippocampus	0.004	-0.015, 0.023	0.45	0.653	0.992
DorsalAttentionLeftAmygdala	0.000	-0.016, 0.016	0.03	0.979	0.992
DorsalAttentionLeftAccumbensArea	0.002	-0.011, 0.014	0.29	0.769	0.992
DorsalAttentionLeftVentraldc	0.000	-0.032, 0.032	-0.02	0.987	0.993
DorsalAttentionRightCerebellumCortex	0.015	-0.005, 0.035	1.48	0.142	0.992
DorsalAttentionRightThalamusProper	0.006	-0.021, 0.033	0.43	0.668	0.992
DorsalAttentionRightCaudate	-0.022	-0.051, 0.007	-1.50	0.138	0.992
DorsalAttentionRightPutamen	0.007	-0.012, 0.026	0.77	0.441	0.992
DorsalAttentionRightPallidum	-0.003	-0.023, 0.018	-0.25	0.802	0.992
DorsalAttentionRightHippocampus	-0.010	-0.032, 0.012	-0.87	0.386	0.992
DorsalAttentionRightAmygdala	-0.003	-0.023, 0.017	-0.29	0.775	0.992
DorsalAttentionRightAccumbensArea	-0.017	-0.043, 0.009	-1.28	0.204	0.992
DorsalAttentionRightVentraldc	0.001	-0.019, 0.021	0.07	0.941	0.992

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
FrontoParietalLeftCerebellumCortex	-0.011	-0.026, 0.005	-1.34	0.183	0.992
FrontoParietalLeftThalamusProper	-0.006	-0.02, 0.008	-0.84	0.401	0.992
FrontoParietalLeftCaudate	0.000	-0.013, 0.012	-0.03	0.978	0.992
FrontoParietalLeftPutamen	-0.009	-0.033, 0.015	-0.73	0.469	0.992
FrontoParietalLeftPallidum	0.000	-0.019, 0.018	-0.05	0.963	0.992
FrontoParietalBrainStem	0.004	-0.025, 0.034	0.30	0.769	0.992
FrontoParietalLeftHippocampus	-0.026	-0.053, 0.001	-1.91	0.060	0.992
FrontoParietalLeftAmygdala	0.011	-0.009, 0.03	1.10	0.273	0.992
FrontoParietalLeftAccumbensArea	0.004	-0.018, 0.027	0.38	0.705	0.992
FrontoParietalLeftVentraldc	0.002	-0.02, 0.025	0.21	0.838	0.992
FrontoParietalRightCerebellumCortex	-0.006	-0.026, 0.014	-0.61	0.541	0.992
FrontoParietalRightThalamusProper	-0.007	-0.037, 0.024	-0.43	0.671	0.992
FrontoParietalRightCaudate	0.005	-0.016, 0.026	0.45	0.650	0.992
FrontoParietalRightPutamen	-0.011	-0.031, 0.009	-1.10	0.275	0.992
FrontoParietalRightPallidum	-0.001	-0.018, 0.015	-0.16	0.872	0.992
FrontoParietalRightHippocampus	0.006	-0.006, 0.018	0.99	0.325	0.992
FrontoParietalRightAmygdala	-0.005	-0.03, 0.02	-0.42	0.675	0.992
FrontoParietalRightAccumbensArea	-0.006	-0.025, 0.013	-0.66	0.512	0.992
FrontoParietalRightVentraldc	-0.002	-0.028, 0.023	-0.17	0.862	0.992
RetrosplenialTemporalLeftCerebellumCortex	0.005	-0.019, 0.028	0.40	0.688	0.992
RetrosplenialTemporalLeftThalamusProper	-0.011	-0.032, 0.01	-1.01	0.314	0.992
RetrosplenialTemporalLeftCaudate	-0.008	-0.03, 0.013	-0.76	0.451	0.992
RetrosplenialTemporalLeftPutamen	-0.003	-0.016, 0.009	-0.53	0.598	0.992
RetrosplenialTemporalLeftPallidum	0.014	-0.012, 0.04	1.10	0.274	0.992
RetrosplenialTemporalBrainStem	-0.008	-0.031, 0.015	-0.67	0.507	0.992
RetrosplenialTemporalLeftHippocampus	0.008	-0.021, 0.037	0.55	0.584	0.992
RetrosplenialTemporalLeftAmygdala	0.004	-0.025, 0.033	0.26	0.792	0.992
RetrosplenialTemporalLeftAccumbensArea	0.003	-0.019, 0.025	0.27	0.785	0.992
RetrosplenialTemporalLeftVentraldc	0.007	-0.02, 0.034	0.53	0.595	0.992
RetrosplenialTemporalRightCerebellumCortex	0.001	-0.027, 0.029	0.09	0.926	0.992
RetrosplenialTemporalRightThalamusProper	0.005	-0.017, 0.027	0.41	0.680	0.992
RetrosplenialTemporalRightCaudate	0.008	-0.027, 0.044	0.47	0.640	0.992
RetrosplenialTemporalRightPutamen	-0.011	-0.031, 0.009	-1.06	0.292	0.992
RetrosplenialTemporalRightPallidum	0.005	-0.012, 0.021	0.56	0.578	0.992
RetrosplenialTemporalRightHippocampus	-0.005	-0.022, 0.012	-0.57	0.570	0.992
RetrosplenialTemporalRightAmygdala	0.007	-0.004, 0.017	1.27	0.207	0.992
RetrosplenialTemporalRightAccumbensArea	-0.025	-0.05, 0.001	-1.94	0.056	0.992
RetrosplenialTemporalRightVentraldc	0.006	-0.022, 0.033	0.40	0.690	0.992
SensorimotorHandLeftCerebellumCortex	-0.013	-0.043, 0.017	-0.84	0.404	0.992
SensorimotorHandLeftThalamusProper	0.001	-0.028, 0.029	0.03	0.974	0.992
SensorimotorHandLeftCaudate	0.017	-0.003, 0.037	1.65	0.102	0.992
SensorimotorHandLeftPutamen	0.005	-0.017, 0.028	0.45	0.656	0.992

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
SensorimotorHandLeftPallidum	0.010	-0.016, 0.036	0.74	0.460	0.992
SensorimotorHandBrainStem	0.008	-0.014, 0.029	0.71	0.477	0.992
SensorimotorHandLeftHippocampus	0.021	-0.012, 0.055	1.26	0.211	0.992
SensorimotorHandLeftAmygdala	-0.007	-0.028, 0.014	-0.67	0.504	0.992
SensorimotorHandLeftAccumbensArea	0.006	-0.013, 0.025	0.65	0.518	0.992
SensorimotorHandLeftVentraldc	-0.005	-0.024, 0.014	-0.57	0.573	0.992
SensorimotorHandRightCerebellumCortex	-0.008	-0.02, 0.004	-1.27	0.206	0.992
SensorimotorHandRightThalamusProper	-0.015	-0.04, 0.01	-1.20	0.235	0.992
SensorimotorHandRightCaudate	0.009	-0.014, 0.032	0.78	0.435	0.992
SensorimotorHandRightPutamen	-0.003	-0.036, 0.029	-0.20	0.838	0.992
SensorimotorHandRightPallidum	0.014	-0.011, 0.039	1.11	0.270	0.992
SensorimotorHandRightHippocampus	0.002	-0.021, 0.025	0.17	0.868	0.992
SensorimotorHandRightAmygdala	0.013	-0.012, 0.038	1.01	0.316	0.992
SensorimotorHandRightAccumbensArea	0.008	-0.017, 0.033	0.64	0.526	0.992
SensorimotorHandRightVentraldc	0.018	-0.004, 0.04	1.61	0.112	0.992
SensorimotorMouthLeftCerebellumCortex	0.008	-0.019, 0.036	0.60	0.553	0.992
SensorimotorMouthLeftThalamusProper	-0.007	-0.028, 0.015	-0.61	0.542	0.992
SensorimotorMouthLeftCaudate	0.006	-0.011, 0.021	0.68	0.497	0.992
SensorimotorMouthLeftPutamen	0.010	-0.007, 0.027	1.16	0.248	0.992
SensorimotorMouthLeftPallidum	-0.001	-0.012, 0.01	-0.19	0.847	0.992
SensorimotorMouthBrainStem	-0.009	-0.036, 0.017	-0.70	0.485	0.992
SensorimotorMouthLeftHippocampus	-0.001	-0.023, 0.022	-0.05	0.958	0.992
SensorimotorMouthLeftAmygdala	-0.008	-0.039, 0.022	-0.55	0.583	0.992
SensorimotorMouthLeftAccumbensArea	0.021	-0.008, 0.05	1.41	0.161	0.992
SensorimotorMouthLeftVentraldc	0.008	-0.012, 0.028	0.78	0.435	0.992
SensorimotorMouthRightCerebellumCortex	-0.012	-0.035, 0.011	-1.06	0.293	0.992
SensorimotorMouthRightThalamusProper	0.004	-0.02, 0.029	0.34	0.732	0.992
SensorimotorMouthRightCaudate	0.011	-0.011, 0.034	0.99	0.326	0.992
SensorimotorMouthRightPutamen	0.013	-0.013, 0.039	1.00	0.322	0.992
SensorimotorMouthRightPallidum	-0.012	-0.03, 0.006	-1.31	0.193	0.992
SensorimotorMouthRightHippocampus	0.009	-0.01, 0.027	0.94	0.350	0.992
SensorimotorMouthRightAmygdala	0.005	-0.01, 0.02	0.66	0.510	0.992
SensorimotorMouthRightAccumbensArea	-0.001	-0.012, 0.01	-0.16	0.871	0.992
SensorimotorMouthRightVentraldc	-0.003	-0.03, 0.024	-0.25	0.803	0.992
SaliencLeftCerebellumCortex	0.001	-0.02, 0.022	0.09	0.926	0.992
SaliencLeftThalamusProper	0.002	-0.022, 0.026	0.14	0.889	0.992
SaliencLeftCaudate	0.017	-0.012, 0.047	1.17	0.244	0.992
SaliencLeftPutamen	0.006	-0.012, 0.023	0.64	0.522	0.992
SaliencLeftPallidum	0.001	-0.02, 0.022	0.10	0.923	0.992
SaliencBrainStem	0.014	-0.007, 0.035	1.33	0.187	0.992
SaliencLeftHippocampus	0.006	-0.014, 0.026	0.58	0.562	0.992
SaliencLeftAmygdala	-0.036	-0.069, -0.003	-2.15	0.034	0.992
SaliencLeftAccumbensArea	-0.028	-0.05, -0.006	-2.56	0.012	0.871

eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
SaliencLeftVentraldc	0.007	-0.012, 0.026	0.71	0.478	0.992
SaliencRightCerebellumCortex	-0.006	-0.022, 0.011	-0.65	0.520	0.992
SaliencRightThalamusProper	0.003	-0.009, 0.015	0.52	0.603	0.992
SaliencRightCaudate	0.004	-0.027, 0.035	0.26	0.793	0.992
SaliencRightPutamen	0.015	-0.003, 0.033	1.64	0.104	0.992
SaliencRightPallidum	0.021	-0.005, 0.046	1.62	0.109	0.992
SaliencRightHippocampus	-0.031	-0.059, -0.003	-2.16	0.033	0.992
SaliencRightAmygdala	0.000	-0.018, 0.019	0.02	0.985	0.993
SaliencRightAccumbensArea	0.006	-0.017, 0.029	0.49	0.627	0.992
SaliencRightVentraldc	-0.005	-0.025, 0.016	-0.43	0.670	0.992
VentralAttentionLeftCerebellumCortex	-0.005	-0.027, 0.016	-0.50	0.619	0.992
VentralAttentionLeftThalamusProper	-0.021	-0.052, 0.01	-1.38	0.172	0.992
VentralAttentionLeftCaudate	-0.008	-0.029, 0.013	-0.76	0.451	0.992
VentralAttentionLeftPutamen	-0.012	-0.03, 0.005	-1.40	0.166	0.992
VentralAttentionLeftPallidum	0.000	-0.015, 0.016	0.03	0.976	0.992
VentralAttentionBrainStem	-0.006	-0.02, 0.008	-0.83	0.410	0.992
VentralAttentionLeftHippocampus	-0.018	-0.048, 0.012	-1.19	0.239	0.992
VentralAttentionLeftAmygdala	-0.001	-0.021, 0.019	-0.11	0.914	0.992
VentralAttentionLeftAccumbensArea	0.010	-0.019, 0.039	0.68	0.500	0.992
VentralAttentionLeftVentraldc	-0.004	-0.032, 0.025	-0.26	0.795	0.992
VentralAttentionRightCerebellumCortex	-0.003	-0.02, 0.015	-0.27	0.784	0.992
VentralAttentionRightThalamusProper	0.014	-0.009, 0.036	1.24	0.220	0.992
VentralAttentionRightCaudate	0.006	-0.017, 0.028	0.49	0.624	0.992
VentralAttentionRightPutamen	0.002	-0.016, 0.02	0.21	0.836	0.992
VentralAttentionRightPallidum	-0.007	-0.036, 0.022	-0.47	0.636	0.992
VentralAttentionRightHippocampus	-0.002	-0.022, 0.018	-0.23	0.816	0.992
VentralAttentionRightAmygdala	-0.013	-0.03, 0.003	-1.61	0.112	0.992
VentralAttentionRightAccumbensArea	-0.017	-0.032, -0.001	-2.16	0.033	0.992
VentralAttentionRightVentraldc	-0.005	-0.016, 0.006	-0.87	0.386	0.992
VisualLeftCerebellumCortex	0.008	-0.018, 0.034	0.60	0.551	0.992
VisualLeftThalamusProper	0.003	-0.014, 0.02	0.34	0.736	0.992
VisualLeftCaudate	0.027	0.001, 0.053	2.04	0.044	0.992
VisualLeftPutamen	0.000	-0.027, 0.027	0.01	0.990	0.993
VisualLeftPallidum	0.005	-0.012, 0.022	0.54	0.589	0.992
VisualBrainStem	0.002	-0.02, 0.023	0.14	0.892	0.992
VisualLeftHippocampus	-0.001	-0.024, 0.023	-0.04	0.968	0.992
VisualLeftAmygdala	-0.003	-0.025, 0.019	-0.28	0.779	0.992
VisualLeftAccumbensArea	0.004	-0.025, 0.033	0.25	0.805	0.992
VisualLeftVentraldc	-0.017	-0.037, 0.004	-1.61	0.110	0.992
VisualRightCerebellumCortex	-0.012	-0.027, 0.004	-1.52	0.132	0.992
VisualRightThalamusProper	-0.008	-0.025, 0.01	-0.88	0.382	0.992
VisualRightCaudate	-0.001	-0.013, 0.01	-0.22	0.823	0.992


eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
VisualRightPutamen	-0.008	-0.035, 0.019	-0.59	0.556	0.992
VisualRightPallidum	0.010	-0.012, 0.031	0.89	0.375	0.992
VisualRightHippocampus	-0.005	-0.033, 0.024	-0.33	0.743	0.992
VisualRightAmygdala	-0.020	-0.051, 0.011	-1.27	0.207	0.992
VisualRightAccumbensArea	0.011	-0.008, 0.029	1.14	0.255	0.992
VisualRightVentraldc	0.012	-0.011, 0.036	1.06	0.292	0.992
WithinAuditory	0.002	-0.016, 0.02	0.27	0.788	0.992
WithinCinguloOpercula	0.005	-0.012, 0.022	0.62	0.538	0.992
WithinCinguloParietal	0.020	-0.029, 0.068	0.81	0.421	0.992
WithinDefault	0.008	-0.007, 0.024	1.04	0.301	0.992
WithinDorsalAttention	0.007	-0.007, 0.02	0.95	0.344	0.992
WithinFrontoParietal	0.009	-0.005, 0.022	1.33	0.187	0.992
WithinRetrosplenialTemporal	0.013	-0.013, 0.039	0.97	0.335	0.992
WithinSensorimotorHand	0.008	-0.009, 0.024	0.89	0.375	0.992
WithinSensorimotorMouth	-0.007	-0.041, 0.027	-0.39	0.699	0.992
WithinSalience	-0.007	-0.04, 0.026	-0.43	0.670	0.992
WithinVentralAttention	-0.004	-0.018, 0.011	-0.51	0.614	0.992
WithinVisual	-0.009	-0.029, 0.012	-0.83	0.406	0.992
AuditoryCinguloOpercular	0.003	-0.012, 0.017	0.36	0.723	0.992
AuditoryCinguloParietal	0.011	-0.007, 0.029	1.24	0.217	0.992
AuditoryDefault	-0.001	-0.013, 0.011	-0.18	0.860	0.992
AuditoryDorsalAttention	0.001	-0.01, 0.01	0.10	0.924	0.992
AuditoryFrontoParietal	0.005	-0.006, 0.016	0.92	0.358	0.992
AuditoryRetrosplenialTemporal	-0.013	-0.027, 0.002	-1.75	0.083	0.992
AuditorySensorimotorHand	0.006	-0.007, 0.019	0.96	0.341	0.992
AuditorySensorimotorMouth	0.001	-0.02, 0.02	0.05	0.963	0.992
AuditorySalience	-0.006	-0.023, 0.011	-0.74	0.462	0.992
AuditoryVentralAttention	0.005	-0.007, 0.017	0.82	0.412	0.992
AuditoryVisual	-0.002	-0.017, 0.013	-0.27	0.788	0.992
CinguloOpercularCinguloParietal	0.005	-0.01, 0.021	0.68	0.501	0.992
CinguloOpercularDefault	-0.006	-0.019, 0.008	-0.84	0.405	0.992
CinguloOpercularDorsalAttention	0.010	0, 0.02	2.00	0.048	0.992
CinguloOpercularFrontoParietal	-0.005	-0.017, 0.006	-0.92	0.360	0.992
CinguloOpercularRetrosplenialTemporal	-0.002	-0.018, 0.014	-0.23	0.817	0.992
CinguloOpercularSensorimotorHand	0.009	-0.002, 0.02	1.71	0.090	0.992
CinguloOpercularSensorimotorMouth	0.000	-0.018, 0.018	-0.03	0.979	0.992
CinguloOpercularSalience	0.000	-0.016, 0.017	0.03	0.975	0.992
CinguloOpercularVentralAttention	0.003	-0.009, 0.015	0.50	0.616	0.992
CinguloOpercularVisual	-0.008	-0.021, 0.005	-1.24	0.219	0.992
CinguloParietalDefault	-0.003	-0.021, 0.016	-0.27	0.785	0.992
CinguloParietalDorsalAttention	0.005	-0.012, 0.021	0.55	0.585	0.992
CinguloParietalFrontoParietal	-0.016	-0.032, 0	-1.94	0.056	0.992
CinguloParietalRetrosplenialTemporal	-0.002	-0.027, 0.022	-0.18	0.859	0.992


eTable 10. Differences in functional connectivity between the never-suicidal group and the concordantly-reported suicidality group: Welch's t-test

rs-fMRI indices	Mean	95%CI	t	p (unc)	p FDR
CinguloParietalSensorimotorHand	0.005	-0.012, 0.022	0.55	0.582	0.992
CinguloParietalSensorimotorMouth	0.007	-0.018, 0.032	0.54	0.594	0.992
CinguloParietalSalience	-0.001	-0.025, 0.024	-0.04	0.966	0.992
CinguloParietalVentralAttention	-0.004	-0.019, 0.011	-0.50	0.621	0.992
CinguloParietalVisual	0.001	-0.019, 0.022	0.13	0.895	0.992
DefaultDorsalAttention	-0.006	-0.018, 0.005	-1.11	0.270	0.992
DefaultFrontoParietal	0.002	-0.007, 0.011	0.44	0.664	0.992
DefaultRetrosplenialTemporal	0.000	-0.015, 0.014	-0.05	0.962	0.992
DefaultSensorimotorHand	-0.009	-0.021, 0.002	-1.59	0.115	0.992
DefaultSensorimotorMouth	0.017	0.004, 0.03	2.63	0.010	0.871
DefaultSalience	0.002	-0.013, 0.017	0.29	0.776	0.992
DefaultVentralAttention	0.002	-0.008, 0.013	0.45	0.652	0.992
DefaultVisual	-0.003	-0.015, 0.009	-0.48	0.636	0.992
DorsalAttentionFrontoParietal	-0.003	-0.013, 0.007	-0.52	0.602	0.992
DorsalAttentionRetrosplenialTemporal	0.003	-0.012, 0.017	0.36	0.718	0.992
DorsalAttentionSensorimotorHand	0.007	-0.002, 0.015	1.59	0.116	0.992
DorsalAttentionSensorimotorMouth	0.000	-0.013, 0.013	0.05	0.961	0.992
DorsalAttentionSalience	-0.005	-0.017, 0.008	-0.71	0.483	0.992
DorsalAttentionVentralAttention	-0.001	-0.012, 0.01	-0.15	0.878	0.992
DorsalAttentionVisual	0.002	-0.01, 0.013	0.27	0.787	0.992
FrontoParietalRetrosplenialTemporal	0.001	-0.011, 0.014	0.22	0.823	0.992
FrontoParietalSensorimotorHand	-0.003	-0.013, 0.008	-0.47	0.642	0.992
FrontoParietalSensorimotorMouth	0.006	-0.007, 0.019	0.89	0.373	0.992
FrontoParietalSalience	0.010	-0.004, 0.024	1.45	0.150	0.992
FrontoParietalVentralAttention	0.004	-0.005, 0.012	0.86	0.392	0.992
FrontoParietalVisual	-0.008	-0.018, 0.003	-1.45	0.150	0.992
RetrosplenialTemporalSensorimotorHand	-0.007	-0.02, 0.007	-0.99	0.323	0.992
RetrosplenialTemporalSensorimotorMouth	-0.006	-0.025, 0.013	-0.61	0.544	0.992
RetrosplenialTemporalSalience	-0.014	-0.033, 0.006	-1.38	0.172	0.992
RetrosplenialTemporalVentralAttention	0.001	-0.014, 0.016	0.14	0.887	0.992
RetrosplenialTemporalVisual	-0.010	-0.027, 0.007	-1.13	0.260	0.992
SensorimotorHandSensorimotorMouth	0.018	-0.001, 0.037	1.87	0.064	0.992
SensorimotorHandSalience	0.006	-0.009, 0.021	0.80	0.424	0.992
SensorimotorHandVentralAttention	0.006	-0.003, 0.015	1.33	0.188	0.992
SensorimotorHandVisual	0.008	-0.006, 0.022	1.09	0.278	0.992
SensorimotorMouthSalience	-0.002	-0.02, 0.016	-0.24	0.811	0.992
SensorimotorMouthVentralAttention	0.004	-0.009, 0.017	0.56	0.576	0.992
SensorimotorMouthVisual	-0.028	-0.045, -0.011	-3.19	0.002	0.598
SalienceVentralAttention	0.004	-0.008, 0.016	0.64	0.524	0.992
SalienceVisual	-0.014	-0.029, 0.001	-1.91	0.059	0.992
VentralAttentionVisual	-0.009	-0.02, 0.002	-1.62	0.108	0.992


Bold denotes group differences at p<0.05 uncorrected




eFigure 8 (1 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 8 (2 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 8 (3 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 8 (4 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 8 (5 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 8 (6 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the child-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (1 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (2 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (3 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (4 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (5 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 9 (6 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the caregiver-reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 10 (1 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 10 (2 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 10 (3 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 10 (4 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 10 (5 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 10 (6 of 6). Results of the Equivalence test for differences in connectivity between the never-suicidal group and the concordantly -reported suicidality group. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

5. Task-Based Functional Imaging

Monetary Incentive Delay task (MID)

Anticipation of large reward vs neutral

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
------	------	-------	---	---------	-------

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.019	-0.046, 0.009	-1.32	0.187	0.600
Left.Thalamus.Proper	-0.008	-0.024, 0.009	-0.92	0.357	0.747
Left.Caudate	-0.004	-0.027, 0.02	-0.30	0.764	0.911
Left.Putamen	-0.008	-0.027, 0.011	-0.83	0.410	0.752
Left.Pallidum	0.005	-0.012, 0.022	0.61	0.544	0.834
Brain.Stem	-0.005	-0.025, 0.015	-0.49	0.624	0.848
Left.Hippocampus	0.000	-0.017, 0.017	0.02	0.984	0.996
Left.Amygdala	0.005	-0.015, 0.025	0.49	0.623	0.848
Left.Accumbens.area	0.008	-0.02, 0.036	0.58	0.565	0.838
Left.VentralDC	-0.012	-0.033, 0.008	-1.19	0.235	0.680
Right.Cerebellum.Cortex	-0.008	-0.034, 0.018	-0.61	0.545	0.834
Right.Thalamus.Proper	-0.010	-0.028, 0.008	-1.08	0.280	0.708
Right.Caudate	0.000	-0.026, 0.026	-0.01	0.990	0.996
Right.Putamen	-0.010	-0.031, 0.011	-0.94	0.347	0.745
Right.Pallidum	0.007	-0.01, 0.025	0.83	0.408	0.752
Right.Hippocampus	-0.007	-0.024, 0.009	-0.85	0.398	0.752
Right.Amygdala	-0.011	-0.035, 0.012	-0.95	0.345	0.745
Right.Accumbens.area	0.009	-0.016, 0.034	0.67	0.504	0.801
Right.VentralDC	0.002	-0.021, 0.024	0.15	0.880	0.973
Cortical regions					
lh_G_and_S_frontomargin	-0.007	-0.045, 0.032	-0.33	0.738	0.898
lh_G_and_S_occipital_inf	-0.026	-0.059, 0.007	-1.56	0.118	0.593
lh_G_and_S_paracentral	-0.022	-0.042, -0.001	-2.03	0.043	0.401
lh_G_and_S_subcentral	-0.024	-0.046, -0.001	-2.05	0.041	0.401
lh_G_and_S_transv_frontopol	-0.004	-0.041, 0.033	-0.22	0.825	0.944
lh_G_and_S_cingul.Ant	-0.002	-0.024, 0.021	-0.13	0.894	0.976
lh_G_and_S_cingul.Mid.Ant	-0.001	-0.02, 0.018	-0.09	0.925	0.982
lh_G_and_S_cingul.Mid.Post	-0.005	-0.023, 0.013	-0.54	0.591	0.848
lh_G_cingul.Post.dorsal	0.004	-0.016, 0.024	0.41	0.679	0.869
lh_G_cingul.Post.ventral	-0.029	-0.07, 0.013	-1.37	0.173	0.593
lh_G_cuneus	-0.038	-0.068, -0.008	-2.48	0.013	0.248
lh_G_front_inf.Opercular	-0.003	-0.022, 0.017	-0.28	0.782	0.927
lh_G_front_inf.Orbital	0.011	-0.021, 0.042	0.68	0.498	0.801
lh_G_front_inf.Triangul	0.001	-0.025, 0.026	0.07	0.941	0.983
lh_G_front_middle	-0.002	-0.021, 0.017	-0.24	0.812	0.944
lh_G_front_sup	-0.004	-0.019, 0.011	-0.48	0.635	0.848
lh_G_Ins_lg_and_S_cent_ins	0.002	-0.018, 0.021	0.16	0.872	0.971
lh_G_insular_short	-0.005	-0.023, 0.014	-0.48	0.630	0.848
lh_G_occipital_middle	-0.014	-0.037, 0.008	-1.25	0.211	0.653
lh_G_occipital_sup	-0.017	-0.042, 0.008	-1.33	0.186	0.600
lh_G_oc.temp_lat.fusifor	-0.022	-0.049, 0.004	-1.64	0.101	0.593
lh_G_oc.temp_med.Lingual	-0.030	-0.064, 0.004	-1.72	0.086	0.535

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_oc.temp_med.Parahip	-0.010	-0.035, 0.015	-0.76	0.450	0.790
lh_G_orbital	-0.001	-0.035, 0.034	-0.03	0.975	0.992
lh_G_pariet_inf.Angular	0.004	-0.015, 0.022	0.39	0.697	0.878
lh_G_pariet_inf.Supramar	-0.018	-0.035, -0.001	-2.05	0.041	0.401
lh_G_parietal_sup	-0.006	-0.026, 0.014	-0.62	0.535	0.834
lh_G_postcentral	-0.017	-0.037, 0.003	-1.63	0.103	0.593
lh_G_precentral	-0.008	-0.024, 0.008	-0.96	0.338	0.745
lh_G_precuneus	-0.008	-0.027, 0.011	-0.84	0.403	0.752
lh_G_rectus	-0.036	-0.104, 0.033	-1.02	0.307	0.742
lh_G_subcallosal	0.004	-0.031, 0.039	0.23	0.815	0.944
lh_G_temp_sup.G_T_transv	-0.029	-0.051, -0.007	-2.64	0.009	0.208
lh_G_temp_sup.Lateral	-0.007	-0.024, 0.011	-0.74	0.458	0.790
lh_G_temp_sup.Plan_polar	0.005	-0.012, 0.021	0.56	0.577	0.838
lh_G_temp_sup.Plan_tempo	-0.022	-0.041, -0.002	-2.22	0.027	0.401
lh_G_temporal_inf	0.004	-0.028, 0.037	0.26	0.794	0.933
lh_G_temporal_middle	-0.001	-0.024, 0.021	-0.10	0.919	0.982
lh_Lat_Fis.ant.Horizontal	0.003	-0.024, 0.03	0.21	0.831	0.944
lh_Lat_Fis.ant.Vertical	-0.001	-0.03, 0.027	-0.09	0.929	0.982
lh_Lat_Fis.post	-0.021	-0.043, 0.002	-1.77	0.077	0.497
lh_Pole_occipital	-0.024	-0.066, 0.019	-1.09	0.278	0.708
lh_Pole_temporal	0.008	-0.022, 0.037	0.51	0.611	0.848
lh_S_calcarine	-0.046	-0.077, -0.016	-3.01	0.003	0.163
lh_S_central	-0.024	-0.047, -0.001	-2.03	0.043	0.401
lh_S_cingul.Marginalis	-0.026	-0.045, -0.008	-2.79	0.005	0.183
lh_S_circular_insula_ant	0.002	-0.021, 0.024	0.14	0.888	0.976
lh_S_circular_insula_inf	-0.013	-0.031, 0.004	-1.49	0.136	0.593
lh_S_circular_insula_sup	-0.004	-0.022, 0.013	-0.49	0.626	0.848
lh_S_collat_transv_ant	-0.024	-0.057, 0.009	-1.40	0.161	0.593
lh_S_collat_transv_post	-0.006	-0.038, 0.025	-0.39	0.700	0.878
lh_S_front_inf	0.010	-0.013, 0.032	0.83	0.406	0.752
lh_S_front_middle	-0.006	-0.029, 0.018	-0.49	0.627	0.848
lh_S_front_sup	-0.002	-0.019, 0.015	-0.21	0.830	0.944
lh_S_interm_prim.Jensen	-0.001	-0.025, 0.022	-0.13	0.900	0.976
lh_S_intrapariet_and_P_trans	-0.008	-0.028, 0.013	-0.73	0.467	0.790
lh_S_oc_middle_and_Lunatus	-0.020	-0.049, 0.009	-1.35	0.177	0.593
lh_S_oc_sup_and_transversal	-0.023	-0.047, 0.001	-1.87	0.062	0.451
lh_S_occipital_ant	-0.009	-0.03, 0.012	-0.88	0.378	0.752
lh_S_oc.temp_lat	-0.008	-0.028, 0.013	-0.74	0.461	0.790
lh_S_oc.temp_med_and_Lingual	-0.011	-0.032, 0.009	-1.13	0.260	0.697
lh_S_orbital_lateral	-0.001	-0.034, 0.032	-0.04	0.971	0.992
lh_S_orbital_med.olfact	0.014	-0.041, 0.07	0.50	0.614	0.848
lh_S_orbital.H_Shaped	-0.004	-0.031, 0.022	-0.33	0.742	0.898
lh_S_parieto_occipital	-0.027	-0.048, -0.007	-2.68	0.008	0.208

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_pericallosal	-0.012	-0.035, 0.011	-1.00	0.319	0.745
lh_S_postcentral	-0.015	-0.033, 0.004	-1.58	0.114	0.593
lh_S_precentral.inf.part	0.000	-0.023, 0.023	0.00	0.998	0.998
lh_S_precentral.sup.part	-0.017	-0.035, 0.001	-1.88	0.061	0.451
lh_S_suborbital	-0.015	-0.062, 0.033	-0.60	0.549	0.834
lh_S_subparietal	0.010	-0.013, 0.033	0.88	0.381	0.752
lh_S_temporal_inf	-0.006	-0.037, 0.026	-0.35	0.729	0.895
lh_S_temporal_sup	-0.008	-0.025, 0.01	-0.84	0.403	0.752
lh_S_temporal_transverse	-0.028	-0.056, 0	-1.95	0.052	0.432
rh_G_and_S_frontomargin	-0.014	-0.055, 0.027	-0.69	0.493	0.801
rh_G_and_S_occipital_inf	-0.015	-0.047, 0.017	-0.92	0.358	0.747
rh_G_and_S_paracental	-0.021	-0.042, 0.001	-1.85	0.065	0.451
rh_G_and_S_subcentral	-0.017	-0.039, 0.005	-1.53	0.127	0.593
rh_G_and_S_transv_frontopol	-0.018	-0.05, 0.014	-1.10	0.271	0.706
rh_G_and_S_cingul.Ant	0.008	-0.015, 0.032	0.68	0.498	0.801
rh_G_and_S_cingul.Mid.Ant	-0.004	-0.023, 0.015	-0.44	0.659	0.853
rh_G_and_S_cingul.Mid.Post	-0.012	-0.03, 0.006	-1.35	0.177	0.593
rh_G_cingul.Post.dorsal	-0.005	-0.025, 0.015	-0.47	0.641	0.850
rh_G_cingul.Post.ventral	-0.033	-0.065, 0	-1.98	0.049	0.428
rh_G_cuneus	-0.022	-0.051, 0.008	-1.43	0.153	0.593
rh_G_front_inf.Opercular	-0.014	-0.033, 0.006	-1.35	0.177	0.593
rh_G_front_inf.Orbital	-0.005	-0.036, 0.026	-0.32	0.747	0.898
rh_G_front_inf.Triangul	-0.001	-0.024, 0.023	-0.05	0.957	0.992
rh_G_front_middle	-0.005	-0.025, 0.016	-0.45	0.655	0.853
rh_G_front_sup	-0.005	-0.02, 0.01	-0.67	0.503	0.801
rh_G_Ins_lg_and_S_cent_ins	-0.016	-0.036, 0.005	-1.48	0.140	0.593
rh_G_insular_short	-0.010	-0.028, 0.009	-1.04	0.301	0.739
rh_G_occipital_middle	-0.013	-0.034, 0.008	-1.23	0.219	0.654
rh_G_occipital_sup	-0.018	-0.041, 0.005	-1.56	0.118	0.593
rh_G_oc.temp_lat.fusifor	-0.021	-0.05, 0.008	-1.45	0.147	0.593
rh_G_oc.temp_med.Lingual	-0.035	-0.068, -0.002	-2.06	0.040	0.401
rh_G_oc.temp_med.Parahip	0.016	-0.012, 0.043	1.12	0.263	0.697
rh_G_orbital	-0.010	-0.045, 0.025	-0.56	0.573	0.838
rh_G_pariet_inf.Angular	-0.006	-0.026, 0.013	-0.66	0.508	0.801
rh_G_pariet_inf.Supramar	-0.025	-0.042, -0.008	-2.88	0.004	0.172
rh_G_parietal_sup	-0.009	-0.028, 0.01	-0.97	0.332	0.745
rh_G_postcentral	-0.012	-0.031, 0.008	-1.15	0.252	0.690
rh_G_precentral	-0.006	-0.021, 0.008	-0.85	0.398	0.752
rh_G_precuneus	-0.007	-0.026, 0.012	-0.68	0.496	0.801
rh_G_rectus	-0.003	-0.082, 0.076	-0.08	0.940	0.983
rh_G_subcallosal	0.019	-0.018, 0.056	1.00	0.320	0.745
rh_G_temp_sup.G_T_transv	-0.017	-0.039, 0.005	-1.51	0.131	0.593
rh_G_temp_sup.Lateral	-0.012	-0.029, 0.005	-1.39	0.165	0.593

eTable 11. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_temp_sup.Plan_polar	-0.003	-0.021, 0.015	-0.35	0.723	0.895
rh_G_temp_sup.Plan_tempo	-0.024	-0.044, -0.005	-2.48	0.013	0.248
rh_G_temporal_inf	-0.010	-0.045, 0.025	-0.57	0.567	0.838
rh_G_temporal_middle	-0.012	-0.033, 0.008	-1.19	0.236	0.680
rh_Lat_Fis.ant.Horizontal	-0.006	-0.03, 0.018	-0.46	0.647	0.851
rh_Lat_Fis.ant.Vertical	0.002	-0.029, 0.033	0.12	0.908	0.979
rh_Lat_Fis.post	-0.015	-0.039, 0.009	-1.23	0.218	0.654
rh_Pole_occipital	-0.035	-0.074, 0.004	-1.77	0.077	0.497
rh_Pole_temporal	0.015	-0.018, 0.048	0.89	0.376	0.752
rh_S_calcarine	-0.037	-0.069, -0.006	-2.34	0.020	0.326
rh_S_central	-0.024	-0.046, -0.002	-2.12	0.035	0.401
rh_S_cingul.Marginalis	-0.030	-0.049, -0.012	-3.23	0.001	0.163
rh_S_circular_insula_ant	-0.011	-0.034, 0.012	-0.94	0.348	0.745
rh_S_circular_insula_inf	-0.022	-0.041, -0.002	-2.19	0.029	0.401
rh_S_circular_insula_sup	-0.004	-0.025, 0.016	-0.41	0.681	0.869
rh_S_collat_transv_ant	-0.007	-0.042, 0.028	-0.38	0.705	0.878
rh_S_collat_transv_post	-0.024	-0.056, 0.008	-1.46	0.144	0.593
rh_S_front_inf	0.009	-0.014, 0.033	0.78	0.437	0.782
rh_S_front_middle	0.000	-0.024, 0.023	-0.03	0.973	0.992
rh_S_front_sup	-0.004	-0.022, 0.013	-0.50	0.614	0.848
rh_S_interm_prim.Jensen	-0.021	-0.043, 0.001	-1.91	0.056	0.447
rh_S_intrapariet_and_P_trans	-0.014	-0.033, 0.006	-1.36	0.175	0.593
rh_S_oc_middle_and_Lunatus	-0.024	-0.054, 0.005	-1.60	0.110	0.593
rh_S_oc_sup_and_transversal	-0.014	-0.038, 0.01	-1.18	0.240	0.680
rh_S_occipital_ant	-0.016	-0.036, 0.005	-1.49	0.137	0.593
rh_S_oc.temp_lat	-0.008	-0.031, 0.014	-0.73	0.468	0.790
rh_S_oc.temp_med_and_Lingual	-0.010	-0.03, 0.01	-0.96	0.336	0.745
rh_S_orbital_lateral	0.013	-0.018, 0.045	0.83	0.405	0.752
rh_S_orbital_med.olfact	0.042	-0.016, 0.1	1.41	0.159	0.593
rh_S_orbital.H_Shaped	-0.003	-0.031, 0.025	-0.19	0.848	0.951
rh_S_parieto_occipital	-0.032	-0.053, -0.011	-2.99	0.003	0.163
rh_S_pericallosal	-0.010	-0.037, 0.016	-0.77	0.440	0.782
rh_S_postcentral	-0.002	-0.018, 0.015	-0.20	0.845	0.951
rh_S_precentral.inf.part	-0.009	-0.031, 0.013	-0.79	0.429	0.779
rh_S_precentral.sup.part	-0.012	-0.03, 0.005	-1.36	0.174	0.593
rh_S_suborbital	0.031	-0.033, 0.095	0.96	0.338	0.745
rh_S_subparietal	-0.007	-0.028, 0.015	-0.59	0.557	0.837
rh_S_temporal_inf	-0.019	-0.053, 0.016	-1.07	0.286	0.713
rh_S_temporal_sup	-0.011	-0.028, 0.006	-1.29	0.197	0.620
rh_S_temporal_transverse	-0.017	-0.047, 0.012	-1.15	0.250	0.690

Bold denotes group differences at p<0.05 uncorrected

eTable 12. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.012	-0.017, 0.041	0.80	0.424	0.919
Left.Thalamus.Proper	0.010	-0.008, 0.028	1.08	0.280	0.919
Left.Caudate	0.019	-0.007, 0.046	1.45	0.149	0.919
Left.Putamen	0.010	-0.011, 0.031	0.89	0.371	0.919
Left.Pallidum	0.003	-0.015, 0.021	0.29	0.774	0.950
Brain.Stem	0.004	-0.017, 0.025	0.39	0.699	0.919
Left.Hippocampus	0.008	-0.01, 0.025	0.86	0.391	0.919
Left.Amygdala	0.010	-0.012, 0.032	0.91	0.362	0.919
Left.Accumbens.area	0.013	-0.017, 0.042	0.85	0.398	0.919
Left.VentralDC	-0.002	-0.025, 0.02	-0.18	0.855	0.950
Right.Cerebellum.Cortex	0.009	-0.019, 0.037	0.63	0.531	0.919
Right.Thalamus.Proper	0.005	-0.015, 0.024	0.47	0.639	0.919
Right.Caudate	0.017	-0.009, 0.044	1.28	0.202	0.919
Right.Putamen	0.016	-0.005, 0.037	1.51	0.131	0.919
Right.Pallidum	0.014	-0.005, 0.032	1.42	0.156	0.919
Right.Hippocampus	0.006	-0.012, 0.024	0.64	0.522	0.919
Right.Amygdala	0.005	-0.019, 0.029	0.40	0.688	0.919
Right.Accumbens.area	0.011	-0.016, 0.038	0.78	0.434	0.919
Right.VentralDC	-0.005	-0.028, 0.018	-0.44	0.659	0.919
Cortical regions					
lh_G_and_S_frontomargin	0.015	-0.027, 0.057	0.68	0.496	0.919
lh_G_and_S_occipital_inf	-0.008	-0.041, 0.026	-0.45	0.654	0.919
lh_G_and_S_paracentral	0.004	-0.017, 0.025	0.38	0.706	0.919
lh_G_and_S_subcentral	-0.010	-0.032, 0.013	-0.86	0.390	0.919
lh_G_and_S_transv_frontopol	0.035	-0.005, 0.074	1.74	0.083	0.919
lh_G_and_S_cingul.Ant	0.014	-0.009, 0.037	1.22	0.222	0.919
lh_G_and_S_cingul.Mid.Ant	0.005	-0.015, 0.024	0.47	0.639	0.919
lh_G_and_S_cingul.Mid.Post	0.017	-0.001, 0.035	1.84	0.067	0.919
lh_G_cingul.Post.dorsal	0.021	-0.001, 0.042	1.86	0.063	0.919
lh_G_cingul.Post.ventral	0.005	-0.038, 0.047	0.21	0.831	0.950
lh_G_cuneus	-0.020	-0.053, 0.012	-1.22	0.222	0.919
lh_G_front_inf.Opercular	0.005	-0.015, 0.026	0.51	0.612	0.919
lh_G_front_inf.Orbital	0.009	-0.024, 0.041	0.52	0.601	0.919
lh_G_front_inf.Triangul	0.009	-0.018, 0.036	0.66	0.512	0.919
lh_G_front_middle	-0.005	-0.026, 0.017	-0.43	0.670	0.919
lh_G_front_sup	0.012	-0.004, 0.028	1.46	0.145	0.919
lh_G_Ins_lg_and_S_cent_ins	-0.002	-0.023, 0.018	-0.22	0.826	0.950
lh_G_insular_short	0.009	-0.01, 0.027	0.94	0.347	0.919
lh_G_occipital_middle	-0.006	-0.029, 0.017	-0.51	0.610	0.919
lh_G_occipital_sup	-0.013	-0.039, 0.014	-0.96	0.340	0.919

eTable 12. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_oc.temp_lat.fusifor	-0.022	-0.05, 0.007	-1.51	0.132	0.919
lh_G_oc.temp_med.Lingual	-0.013	-0.05, 0.023	-0.73	0.466	0.919
lh_G_oc.temp_med.Parahip	-0.004	-0.028, 0.021	-0.29	0.772	0.950
lh_G_orbital	0.015	-0.021, 0.052	0.82	0.411	0.919
lh_G_pariet_inf.Angular	0.012	-0.008, 0.032	1.15	0.252	0.919
lh_G_pariet_inf.Supramar	-0.004	-0.022, 0.014	-0.41	0.682	0.919
lh_G_parietal_sup	0.007	-0.012, 0.027	0.74	0.461	0.919
lh_G_postcentral	0.000	-0.022, 0.021	-0.02	0.986	0.996
lh_G_precentral	0.002	-0.016, 0.019	0.19	0.851	0.950
lh_G_precuneus	0.009	-0.011, 0.028	0.85	0.396	0.919
lh_G_rectus	0.002	-0.07, 0.074	0.06	0.949	0.973
lh_G_subcallosal	0.007	-0.031, 0.045	0.38	0.705	0.919
lh_G_temp_sup.G_T_transv	-0.005	-0.028, 0.017	-0.46	0.648	0.919
lh_G_temp_sup.Lateral	0.012	-0.006, 0.031	1.31	0.192	0.919
lh_G_temp_sup.Plan_polar	0.005	-0.013, 0.023	0.54	0.587	0.919
lh_G_temp_sup.Plan_tempo	-0.015	-0.036, 0.006	-1.44	0.151	0.919
lh_G_temporal_inf	0.000	-0.034, 0.034	0.01	0.993	0.996
lh_G_temporal_middle	0.002	-0.022, 0.025	0.13	0.896	0.950
lh_Lat_Fis.ant.Horizontal	0.014	-0.014, 0.043	1.00	0.318	0.919
lh_Lat_Fis.ant.Vertical	0.004	-0.026, 0.034	0.26	0.798	0.950
lh_Lat_Fis.post	-0.017	-0.04, 0.006	-1.41	0.158	0.919
lh_Pole_occipital	0.019	-0.025, 0.064	0.85	0.394	0.919
lh_Pole_temporal	-0.007	-0.038, 0.024	-0.45	0.656	0.919
lh_S_calcarine	-0.036	-0.069, -0.003	-2.16	0.032	0.919
lh_S_central	-0.004	-0.027, 0.019	-0.34	0.735	0.924
lh_S_cingul.Marginalis	-0.007	-0.026, 0.012	-0.70	0.486	0.919
lh_S_circular_insula_ant	0.022	-0.001, 0.045	1.86	0.064	0.919
lh_S_circular_insula_inf	0.000	-0.019, 0.019	0.00	0.996	0.996
lh_S_circular_insula_sup	-0.004	-0.024, 0.016	-0.41	0.679	0.919
lh_S_collat_transv_ant	-0.012	-0.046, 0.022	-0.69	0.494	0.919
lh_S_collat_transv_post	-0.017	-0.051, 0.018	-0.96	0.337	0.919
lh_S_front_inf	0.014	-0.011, 0.04	1.08	0.279	0.919
lh_S_front_middle	0.007	-0.018, 0.031	0.53	0.598	0.919
lh_S_front_sup	0.006	-0.012, 0.024	0.61	0.542	0.919
lh_S_interm_prim.Jensen	0.016	-0.009, 0.042	1.25	0.212	0.919
lh_S_intrapariet_and_P_trans	0.001	-0.02, 0.023	0.12	0.903	0.950
lh_S_oc_middle_and_Lunatus	-0.008	-0.038, 0.023	-0.48	0.628	0.919
lh_S_oc_sup_and_transversal	-0.016	-0.041, 0.009	-1.24	0.217	0.919
lh_S_occipital_ant	0.009	-0.015, 0.032	0.73	0.468	0.919
lh_S_oc.temp_lat	-0.014	-0.035, 0.007	-1.32	0.189	0.919
lh_S_oc.temp_med_and_Lingual	-0.014	-0.034, 0.005	-1.41	0.158	0.919
lh_S_orbital_lateral	0.012	-0.021, 0.045	0.72	0.474	0.919
lh_S_orbital_med.olfact	0.004	-0.058, 0.065	0.12	0.905	0.950

eTable 12. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital.H_Shaped	0.005	-0.023, 0.033	0.35	0.723	0.924
lh_S_parieto_occipital	-0.009	-0.03, 0.012	-0.80	0.424	0.919
lh_S_pericallosal	0.010	-0.013, 0.033	0.84	0.402	0.919
lh_S_postcentral	0.010	-0.009, 0.029	1.02	0.306	0.919
lh_S_precentral.inf.part	0.009	-0.014, 0.032	0.75	0.452	0.919
lh_S_precentral.sup.part	-0.004	-0.024, 0.015	-0.43	0.668	0.919
lh_S_suborbital	-0.005	-0.052, 0.042	-0.20	0.842	0.950
lh_S_subparietal	0.001	-0.023, 0.026	0.10	0.921	0.956
lh_S_temporal_inf	0.002	-0.031, 0.036	0.15	0.885	0.950
lh_S_temporal_sup	-0.004	-0.023, 0.015	-0.43	0.668	0.919
lh_S_temporal_transverse	0.002	-0.027, 0.031	0.15	0.881	0.950
rh_G_and_S_frontomargin	-0.023	-0.067, 0.02	-1.05	0.294	0.919
rh_G_and_S_occipital_inf	-0.008	-0.041, 0.026	-0.45	0.655	0.919
rh_G_and_S_paracentral	-0.001	-0.023, 0.021	-0.08	0.934	0.963
rh_G_and_S_subcentral	-0.003	-0.028, 0.021	-0.28	0.780	0.950
rh_G_and_S_transv_frontopol	0.017	-0.016, 0.049	0.99	0.322	0.919
rh_G_and_S_cingul.Ant	0.017	-0.007, 0.041	1.38	0.169	0.919
rh_G_and_S_cingul.Mid.Ant	0.000	-0.019, 0.02	0.01	0.989	0.996
rh_G_and_S_cingul.Mid.Post	0.002	-0.016, 0.02	0.18	0.856	0.950
rh_G_cingul.Post.dorsal	0.006	-0.015, 0.026	0.56	0.576	0.919
rh_G_cingul.Post.ventral	0.010	-0.025, 0.045	0.58	0.563	0.919
rh_G_cuneus	-0.010	-0.041, 0.021	-0.62	0.534	0.919
rh_G_front_inf.Opercular	0.011	-0.009, 0.03	1.06	0.292	0.919
rh_G_front_inf.Orbital	0.014	-0.017, 0.044	0.90	0.369	0.919
rh_G_front_inf.Triangul	0.014	-0.01, 0.039	1.15	0.252	0.919
rh_G_front_middle	0.012	-0.01, 0.033	1.07	0.285	0.919
rh_G_front_sup	0.005	-0.012, 0.021	0.56	0.579	0.919
rh_G_Ins_lg_and_S_cent_ins	0.004	-0.018, 0.026	0.38	0.705	0.919
rh_G_insular_short	0.008	-0.012, 0.027	0.78	0.436	0.919
rh_G_occipital_middle	-0.004	-0.026, 0.019	-0.34	0.736	0.924
rh_G_occipital_sup	-0.007	-0.033, 0.019	-0.54	0.591	0.919
rh_G_oc.temp_lat.fusifor	-0.006	-0.035, 0.022	-0.43	0.668	0.919
rh_G_oc.temp_med.Lingual	-0.026	-0.061, 0.009	-1.46	0.145	0.919
rh_G_oc.temp_med.Parahip	0.012	-0.014, 0.038	0.91	0.361	0.919
rh_G_orbital	0.011	-0.028, 0.05	0.56	0.575	0.919
rh_G_pariet_inf.Angular	0.006	-0.014, 0.026	0.59	0.558	0.919
rh_G_pariet_inf.Supramar	0.004	-0.014, 0.023	0.45	0.653	0.919
rh_G_parietal_sup	0.006	-0.014, 0.026	0.57	0.570	0.919
rh_G_postcentral	-0.006	-0.026, 0.015	-0.53	0.596	0.919
rh_G_precentral	0.010	-0.007, 0.026	1.14	0.257	0.919
rh_G_precuneus	0.007	-0.013, 0.027	0.70	0.483	0.919
rh_G_rectus	-0.018	-0.104, 0.068	-0.40	0.687	0.919
rh_G_subcallosal	0.015	-0.02, 0.051	0.84	0.404	0.919

eTable 12. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_temp_sup.G_T_transv	0.005	-0.018, 0.028	0.46	0.646	0.919
rh_G_temp_sup.Lateral	-0.001	-0.019, 0.017	-0.13	0.899	0.950
rh_G_temp_sup.Plan_polar	0.002	-0.016, 0.02	0.22	0.826	0.950
rh_G_temp_sup.Plan_tempo	0.005	-0.016, 0.025	0.43	0.669	0.919
rh_G_temporal_inf	0.003	-0.031, 0.038	0.19	0.849	0.950
rh_G_temporal_middle	0.012	-0.009, 0.033	1.11	0.267	0.919
rh_Lat_Fis.ant.Horizontal	-0.003	-0.029, 0.023	-0.22	0.824	0.950
rh_Lat_Fis.ant.Vertical	0.018	-0.014, 0.051	1.10	0.273	0.919
rh_Lat_Fis.post	-0.001	-0.025, 0.023	-0.10	0.922	0.956
rh_Pole_occipital	0.012	-0.029, 0.052	0.56	0.578	0.919
rh_Pole_temporal	0.005	-0.03, 0.039	0.26	0.796	0.950
rh_S_calcarine	-0.034	-0.067, -0.002	-2.11	0.035	0.919
rh_S_central	-0.014	-0.037, 0.01	-1.13	0.259	0.919
rh_S_cingul.Marginalis	-0.010	-0.03, 0.009	-1.03	0.305	0.919
rh_S_circular_insula_ant	0.006	-0.017, 0.029	0.54	0.591	0.919
rh_S_circular_insula_inf	-0.004	-0.024, 0.016	-0.37	0.710	0.919
rh_S_circular_insula_sup	0.007	-0.014, 0.027	0.65	0.517	0.919
rh_S_collat_transv_ant	0.038	0, 0.076	1.95	0.052	0.919
rh_S_collat_transv_post	-0.014	-0.048, 0.019	-0.85	0.395	0.919
rh_S_front_inf	0.011	-0.014, 0.036	0.88	0.377	0.919
rh_S_front_middle	0.010	-0.015, 0.035	0.78	0.436	0.919
rh_S_front_sup	0.009	-0.009, 0.027	1.00	0.317	0.919
rh_S_interm_prim.Jensen	-0.003	-0.026, 0.02	-0.25	0.800	0.950
rh_S_intrapariet_and_P_trans	0.005	-0.015, 0.025	0.52	0.604	0.919
rh_S_oc_middle_and_Lunatus	-0.002	-0.032, 0.028	-0.14	0.889	0.950
rh_S_oc_sup_and_transversal	-0.007	-0.032, 0.017	-0.59	0.555	0.919
rh_S_occipital_ant	-0.002	-0.024, 0.02	-0.17	0.865	0.950
rh_S_oc.temp_lat	0.002	-0.02, 0.024	0.14	0.885	0.950
rh_S_oc.temp_med_and_Lingual	-0.005	-0.026, 0.016	-0.48	0.632	0.919
rh_S_orbital_lateral	0.034	0.001, 0.068	2.03	0.043	0.919
rh_S_orbital_med.olfact	0.049	-0.014, 0.111	1.53	0.126	0.919
rh_S_orbital.H_Shaped	0.019	-0.01, 0.048	1.32	0.189	0.919
rh_S_parieto_occipital	-0.014	-0.035, 0.007	-1.30	0.194	0.919
rh_S_pericallosal	0.018	-0.009, 0.044	1.29	0.196	0.919
rh_S_postcentral	0.004	-0.015, 0.022	0.39	0.694	0.919
rh_S_precentral.inf.part	0.009	-0.014, 0.031	0.73	0.466	0.919
rh_S_precentral.sup.part	0.012	-0.007, 0.03	1.23	0.220	0.919
rh_S_suborbital	0.012	-0.056, 0.08	0.34	0.733	0.924
rh_S_subparietal	0.004	-0.019, 0.027	0.33	0.744	0.927
rh_S_temporal_inf	0.029	-0.006, 0.064	1.65	0.100	0.919
rh_S_temporal_sup	0.008	-0.01, 0.025	0.87	0.384	0.919
rh_S_temporal_transverse	-0.004	-0.035, 0.028	-0.24	0.807	0.950

Bold denotes group differences at p<0.05 uncorrected

eTable 13. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.022	-0.078, 0.035	-0.76	0.450	0.974
Left.Thalamus.Proper	-0.006	-0.044, 0.032	-0.31	0.754	0.974
Left.Caudate	0.008	-0.044, 0.06	0.30	0.763	0.974
Left.Putamen	-0.009	-0.047, 0.03	-0.44	0.660	0.974
Left.Pallidum	0.008	-0.027, 0.044	0.46	0.643	0.974
Brain.Stem	-0.009	-0.045, 0.028	-0.48	0.635	0.974
Left.Hippocampus	0.000	-0.032, 0.033	0.02	0.987	0.994
Left.Amygdala	0.001	-0.04, 0.043	0.07	0.944	0.988
Left.Accumbens.area	0.038	-0.022, 0.098	1.25	0.213	0.974
Left.VentralDC	-0.019	-0.06, 0.022	-0.92	0.358	0.974
Right.Cerebellum.Cortex	0.022	-0.031, 0.076	0.82	0.414	0.974
Right.Thalamus.Proper	-0.003	-0.042, 0.036	-0.14	0.887	0.974
Right.Caudate	0.030	-0.025, 0.086	1.08	0.282	0.974
Right.Putamen	0.012	-0.029, 0.053	0.58	0.566	0.974
Right.Pallidum	0.016	-0.015, 0.048	1.02	0.309	0.974
Right.Hippocampus	-0.002	-0.037, 0.033	-0.10	0.918	0.974
Right.Amygdala	0.000	-0.052, 0.051	-0.02	0.985	0.994
Right.Accumbens.area	0.029	-0.024, 0.083	1.09	0.278	0.974
Right.VentralDC	0.006	-0.041, 0.052	0.24	0.814	0.974
Cortical regions					
lh_G_and_S_frontomargin	0.023	-0.061, 0.108	0.55	0.585	0.974
lh_G_and_S_occipital_inf	-0.038	-0.103, 0.026	-1.18	0.239	0.974
lh_G_and_S_paracentral	-0.018	-0.06, 0.023	-0.86	0.390	0.974
lh_G_and_S_subcentral	-0.024	-0.072, 0.024	-1.00	0.319	0.974
lh_G_and_S_transv_frontopol	0.023	-0.054, 0.101	0.60	0.549	0.974
lh_G_and_S_cingul.Ant	0.000	-0.046, 0.047	0.00	0.998	0.998
lh_G_and_S_cingul.Mid.Ant	0.016	-0.025, 0.057	0.76	0.448	0.974
lh_G_and_S_cingul.Mid.Post	0.002	-0.036, 0.041	0.11	0.913	0.974
lh_G_cingul.Post.dorsal	0.022	-0.02, 0.063	1.03	0.305	0.974
lh_G_cingul.Post.ventral	-0.065	-0.14, 0.01	-1.72	0.088	0.974
lh_G_cuneus	-0.064	-0.124, -0.004	-2.11	0.038	0.928
lh_G_front_inf.Opercular	-0.016	-0.059, 0.027	-0.76	0.451	0.974
lh_G_front_inf.Orbital	-0.024	-0.095, 0.047	-0.67	0.504	0.974
lh_G_front_inf.Triangul	0.017	-0.036, 0.07	0.63	0.530	0.974
lh_G_front_middle	-0.009	-0.049, 0.031	-0.46	0.648	0.974
lh_G_front_sup	-0.005	-0.037, 0.028	-0.30	0.763	0.974
lh_G_Ins_lg_and_S_cent_ins	-0.034	-0.074, 0.007	-1.63	0.107	0.974
lh_G_insular_short	0.002	-0.034, 0.038	0.10	0.921	0.974

eTable 13. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_occipital_middle	-0.017	-0.06, 0.025	-0.82	0.414	0.974
lh_G_occipital_sup	-0.016	-0.068, 0.036	-0.60	0.550	0.974
lh_G_oc.temp_lat.fusifor	-0.031	-0.086, 0.023	-1.15	0.252	0.974
lh_G_oc.temp_med.Lingual	-0.053	-0.122, 0.016	-1.52	0.131	0.974
lh_G_oc.temp_med.Parahip	-0.040	-0.088, 0.009	-1.63	0.106	0.974
lh_G_orbital	-0.004	-0.07, 0.061	-0.13	0.898	0.974
lh_G_pariet_inf.Angular	0.005	-0.029, 0.04	0.30	0.762	0.974
lh_G_pariet_inf.Supramar	-0.029	-0.065, 0.008	-1.57	0.120	0.974
lh_G_parietal_sup	0.002	-0.035, 0.04	0.13	0.896	0.974
lh_G_postcentral	-0.011	-0.052, 0.029	-0.56	0.578	0.974
lh_G_precentral	-0.011	-0.045, 0.023	-0.62	0.538	0.974
lh_G_precuneus	-0.009	-0.046, 0.028	-0.47	0.636	0.974
lh_G_rectus	-0.056	-0.204, 0.093	-0.75	0.457	0.974
lh_G_subcallosal	-0.007	-0.081, 0.068	-0.17	0.863	0.974
lh_G_temp_sup.G_T_transv	-0.042	-0.087, 0.003	-1.86	0.066	0.974
lh_G_temp_sup.Lateral	-0.008	-0.044, 0.029	-0.42	0.679	0.974
lh_G_temp_sup.Plan_polar	0.009	-0.025, 0.044	0.54	0.589	0.974
lh_G_temp_sup.Plan_tempo	-0.047	-0.087, -0.006	-2.30	0.024	0.916
lh_G_temporal_inf	-0.014	-0.084, 0.057	-0.39	0.701	0.974
lh_G_temporal_middle	-0.021	-0.069, 0.028	-0.85	0.399	0.974
lh_Lat_Fis.ant.Horizontal	-0.005	-0.063, 0.053	-0.18	0.856	0.974
lh_Lat_Fis.ant.Vertical	-0.029	-0.092, 0.034	-0.91	0.366	0.974
lh_Lat_Fis.post	-0.013	-0.058, 0.031	-0.59	0.560	0.974
lh_Pole_occipital	0.024	-0.06, 0.109	0.57	0.570	0.974
lh_Pole_temporal	-0.003	-0.061, 0.055	-0.11	0.916	0.974
lh_S_calcarine	-0.103	-0.166, -0.04	-3.24	0.002	0.267
lh_S_central	-0.038	-0.088, 0.011	-1.55	0.125	0.974
lh_S_cingul.Marginalis	-0.032	-0.069, 0.005	-1.71	0.091	0.974
lh_S_circular_insula_ant	0.008	-0.041, 0.058	0.33	0.743	0.974
lh_S_circular_insula_inf	-0.027	-0.063, 0.009	-1.50	0.136	0.974
lh_S_circular_insula_sup	-0.002	-0.041, 0.037	-0.11	0.910	0.974
lh_S_collat_transv_ant	-0.007	-0.072, 0.059	-0.20	0.843	0.974
lh_S_collat_transv_post	-0.017	-0.083, 0.049	-0.52	0.601	0.974
lh_S_front_inf	0.001	-0.049, 0.05	0.03	0.977	0.994
lh_S_front_middle	-0.001	-0.051, 0.049	-0.03	0.976	0.994
lh_S_front_sup	-0.012	-0.049, 0.024	-0.67	0.504	0.974
lh_S_interm_prim.Jensen	-0.026	-0.077, 0.025	-1.00	0.322	0.974
lh_S_intrapariet_and_P_trans	0.004	-0.034, 0.043	0.21	0.833	0.974
lh_S_oc_middle_and_Lunatus	-0.028	-0.083, 0.027	-1.02	0.311	0.974
lh_S_oc_sup_and_transversal	-0.028	-0.076, 0.02	-1.16	0.249	0.974
lh_S_occipital_ant	-0.014	-0.058, 0.029	-0.64	0.522	0.974
lh_S_oc.temp_lat	-0.022	-0.061, 0.017	-1.12	0.268	0.974
lh_S_oc.temp_med_and_Lingual	-0.025	-0.064, 0.013	-1.31	0.192	0.974

eTable 13. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital_lateral	-0.016	-0.081, 0.049	-0.49	0.627	0.974
lh_S_orbital_med.olfact	0.020	-0.105, 0.144	0.31	0.755	0.974
lh_S_orbital.H_Shaped	-0.019	-0.077, 0.038	-0.67	0.507	0.974
lh_S_parieto_occipital	-0.045	-0.082, -0.007	-2.36	0.020	0.916
lh_S_pericallosal	-0.007	-0.055, 0.04	-0.32	0.753	0.974
lh_S_postcentral	0.004	-0.035, 0.043	0.19	0.851	0.974
lh_S_precentral.inf.part	0.003	-0.042, 0.049	0.15	0.880	0.974
lh_S_precentral.sup.part	-0.018	-0.058, 0.022	-0.89	0.378	0.974
lh_S_suborbital	-0.031	-0.13, 0.068	-0.62	0.537	0.974
lh_S_subparietal	-0.004	-0.049, 0.041	-0.18	0.856	0.974
lh_S_temporal_inf	-0.035	-0.098, 0.029	-1.09	0.280	0.974
lh_S_temporal_sup	-0.024	-0.062, 0.014	-1.26	0.209	0.974
lh_S_temporal_transverse	-0.053	-0.108, 0.002	-1.91	0.060	0.974
rh_G_and_S_frontomargin	-0.022	-0.108, 0.065	-0.50	0.621	0.974
rh_G_and_S_occipital_inf	-0.031	-0.09, 0.029	-1.02	0.311	0.974
rh_G_and_S_paracentral	-0.018	-0.06, 0.024	-0.86	0.393	0.974
rh_G_and_S_subcentral	-0.030	-0.076, 0.016	-1.30	0.197	0.974
rh_G_and_S_transv_frontopol	-0.040	-0.104, 0.024	-1.25	0.214	0.974
rh_G_and_S_cingul.Ant	0.030	-0.019, 0.08	1.22	0.227	0.974
rh_G_and_S_cingul.Mid.Ant	0.008	-0.031, 0.048	0.42	0.679	0.974
rh_G_and_S_cingul.Mid.Post	-0.013	-0.05, 0.023	-0.74	0.462	0.974
rh_G_cingul.Post.dorsal	0.016	-0.02, 0.053	0.88	0.379	0.974
rh_G_cingul.Post.ventral	-0.031	-0.09, 0.029	-1.02	0.311	0.974
rh_G_cuneus	-0.050	-0.107, 0.008	-1.70	0.092	0.974
rh_G_front_inf.Opercular	0.002	-0.037, 0.041	0.11	0.913	0.974
rh_G_front_inf.Orbital	-0.007	-0.067, 0.053	-0.23	0.816	0.974
rh_G_front_inf.Triangul	-0.001	-0.043, 0.04	-0.06	0.954	0.989
rh_G_front_middle	0.014	-0.027, 0.054	0.66	0.509	0.974
rh_G_front_sup	-0.007	-0.039, 0.025	-0.44	0.659	0.974
rh_G_Ins_lg_and_S_cent_ins	-0.015	-0.058, 0.027	-0.71	0.480	0.974
rh_G_insular_short	-0.005	-0.045, 0.035	-0.27	0.788	0.974
rh_G_occipital_middle	-0.003	-0.041, 0.035	-0.16	0.870	0.974
rh_G_occipital_sup	-0.032	-0.076, 0.012	-1.43	0.157	0.974
rh_G_oc.temp_lat.fusifor	-0.041	-0.094, 0.013	-1.52	0.133	0.974
rh_G_oc.temp_med.Lingual	-0.075	-0.141, -0.008	-2.24	0.027	0.916
rh_G_oc.temp_med.Parahip	0.006	-0.044, 0.057	0.25	0.803	0.974
rh_G_orbital	0.024	-0.052, 0.1	0.63	0.532	0.974
rh_G_pariet_inf.Angular	0.003	-0.031, 0.037	0.17	0.864	0.974
rh_G_pariet_inf.Supramar	-0.008	-0.046, 0.029	-0.44	0.659	0.974
rh_G_parietal_sup	0.000	-0.039, 0.038	-0.01	0.988	0.994
rh_G_postcentral	-0.015	-0.054, 0.023	-0.79	0.434	0.974
rh_G_precentral	-0.003	-0.034, 0.028	-0.18	0.858	0.974
rh_G_precuneus	0.003	-0.03, 0.036	0.18	0.855	0.974


eTable 13. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_rectus	0.015	-0.138, 0.167	0.19	0.849	0.974
rh_G_subcallosal	0.018	-0.058, 0.095	0.47	0.638	0.974
rh_G_temp_sup.G_T_transv	-0.027	-0.073, 0.019	-1.18	0.243	0.974
rh_G_temp_sup.Lateral	-0.017	-0.051, 0.018	-0.95	0.343	0.974
rh_G_temp_sup.Plan_polar	0.012	-0.021, 0.045	0.70	0.486	0.974
rh_G_temp_sup.Plan_tempo	-0.018	-0.056, 0.019	-0.96	0.339	0.974
rh_G_temporal_inf	-0.004	-0.066, 0.059	-0.11	0.911	0.974
rh_G_temporal_middle	0.009	-0.032, 0.049	0.42	0.676	0.974
rh_Lat_Fis.ant.Horizontal	-0.012	-0.058, 0.033	-0.54	0.591	0.974
rh_Lat_Fis.ant.Vertical	-0.019	-0.079, 0.041	-0.63	0.532	0.974
rh_Lat_Fis.post	-0.005	-0.051, 0.042	-0.20	0.845	0.974
rh_Pole_occipital	0.013	-0.066, 0.092	0.32	0.748	0.974
rh_Pole_temporal	0.039	-0.025, 0.102	1.21	0.228	0.974
rh_S_calcarine	-0.086	-0.143, -0.029	-3.00	0.003	0.284
rh_S_central	-0.043	-0.09, 0.005	-1.77	0.079	0.974
rh_S_cingul.Marginalis	-0.023	-0.06, 0.013	-1.29	0.201	0.974
rh_S_circular_insula_ant	-0.005	-0.052, 0.041	-0.23	0.815	0.974
rh_S_circular_insula_inf	-0.002	-0.042, 0.037	-0.12	0.907	0.974
rh_S_circular_insula_sup	0.006	-0.033, 0.046	0.32	0.752	0.974
rh_S_collat_transv_ant	0.008	-0.069, 0.085	0.21	0.837	0.974
rh_S_collat_transv_post	-0.046	-0.109, 0.017	-1.46	0.147	0.974
rh_S_front_inf	0.024	-0.023, 0.071	1.02	0.310	0.974
rh_S_front_middle	0.010	-0.036, 0.056	0.41	0.680	0.974
rh_S_front_sup	0.006	-0.029, 0.041	0.34	0.733	0.974
rh_S_interm_prim.Jensen	-0.025	-0.068, 0.019	-1.13	0.262	0.974
rh_S_intrapariet_and_P_trans	0.002	-0.032, 0.036	0.12	0.906	0.974
rh_S_oc_middle_and_Lunatus	-0.014	-0.072, 0.045	-0.46	0.643	0.974
rh_S_oc_sup_and_transversal	-0.044	-0.09, 0.003	-1.87	0.064	0.974
rh_S_occipital_ant	-0.003	-0.041, 0.035	-0.16	0.873	0.974
rh_S_oc.temp_lat	-0.021	-0.064, 0.021	-1.01	0.317	0.974
rh_S_oc.temp_med_and_Lingual	-0.022	-0.062, 0.017	-1.11	0.269	0.974
rh_S_orbital_lateral	0.028	-0.032, 0.089	0.94	0.350	0.974
rh_S_orbital_med.olfact	0.109	-0.012, 0.229	1.79	0.076	0.974
rh_S_orbital.H_Shaped	0.021	-0.034, 0.076	0.76	0.447	0.974
rh_S_parieto_occipital	-0.041	-0.081, -0.002	-2.09	0.039	0.928
rh_S_pericallosal	0.016	-0.039, 0.071	0.58	0.563	0.974
rh_S_postcentral	0.007	-0.024, 0.038	0.44	0.662	0.974
rh_S_precentral.inf.part	0.003	-0.04, 0.046	0.13	0.894	0.974
rh_S_precentral.sup.part	-0.003	-0.042, 0.036	-0.14	0.886	0.974
rh_S_suborbital	0.047	-0.082, 0.176	0.73	0.468	0.974
rh_S_subparietal	-0.009	-0.051, 0.033	-0.42	0.674	0.974
rh_S_temporal_inf	-0.002	-0.074, 0.069	-0.07	0.946	0.988
rh_S_temporal_sup	-0.009	-0.042, 0.024	-0.55	0.584	0.974


eTable 13. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_transverse	-0.027	-0.082, 0.028	-0.97	0.336	0.974


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 12. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 13 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 13 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 13 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 14. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 15 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 15 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 15 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 16. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 17 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 17 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 17 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of reward. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Anticipation of large loss vs neutral

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.016	-0.044, 0.013	-1.09	0.278	0.979
Left.Thalamus.Proper	-0.001	-0.017, 0.014	-0.14	0.888	0.979
Left.Caudate	0.009	-0.013, 0.032	0.83	0.408	0.979
Left.Putamen	0.001	-0.016, 0.019	0.17	0.864	0.979
Left.Pallidum	0.008	-0.01, 0.025	0.87	0.387	0.979
Brain.Stem	0.005	-0.014, 0.023	0.50	0.615	0.979
Left.Hippocampus	0.005	-0.011, 0.022	0.66	0.509	0.979
Left.Amygdala	0.014	-0.006, 0.035	1.37	0.172	0.979
Left.Accumbens.area	0.021	-0.005, 0.047	1.55	0.121	0.979
Left.VentralDC	0.004	-0.015, 0.023	0.43	0.665	0.979
Right.Cerebellum.Cortex	0.008	-0.02, 0.035	0.57	0.569	0.979
Right.Thalamus.Proper	-0.001	-0.018, 0.015	-0.16	0.869	0.979
Right.Caudate	0.003	-0.021, 0.027	0.24	0.808	0.979
Right.Putamen	0.005	-0.013, 0.024	0.58	0.562	0.979
Right.Pallidum	0.009	-0.008, 0.026	1.00	0.316	0.979
Right.Hippocampus	0.008	-0.008, 0.024	0.95	0.341	0.979
Right.Amygdala	0.008	-0.014, 0.03	0.73	0.466	0.979
Right.Accumbens.area	0.016	-0.009, 0.042	1.24	0.214	0.979
Right.VentralDC	0.002	-0.018, 0.023	0.21	0.831	0.979
Cortical regions					
lh_G_and_S_frontomargin	-0.018	-0.051, 0.014	-1.10	0.271	0.979
lh_G_and_S_occipital_inf	-0.006	-0.026, 0.014	-0.56	0.573	0.979
lh_G_and_S_paracentral	-0.016	-0.037, 0.004	-1.56	0.120	0.979
lh_G_and_S_subcentral	0.012	-0.024, 0.048	0.65	0.518	0.979
lh_G_and_S_transv_frontopol	0.004	-0.017, 0.025	0.39	0.693	0.979
lh_G_and_S_cingul.Ant	0.003	-0.015, 0.021	0.31	0.755	0.979
lh_G_and_S_cingul.Mid.Ant	-0.001	-0.017, 0.016	-0.10	0.918	0.979
lh_G_and_S_cingul.Mid.Post	0.003	-0.016, 0.022	0.33	0.740	0.979
lh_G_cingul.Post.dorsal	-0.005	-0.044, 0.035	-0.24	0.811	0.979
lh_G_cingul.Post.ventral	-0.027	-0.059, 0.004	-1.70	0.091	0.979
lh_G_cuneus	-0.008	-0.027, 0.011	-0.86	0.391	0.979
lh_G_front_inf.Opercular	0.032	0.002, 0.062	2.10	0.036	0.979
lh_G_front_inf.Orbital	0.009	-0.016, 0.034	0.74	0.460	0.979
lh_G_front_inf.Triangul	-0.004	-0.023, 0.016	-0.38	0.705	0.979
lh_G_front_middle	-0.001	-0.015, 0.013	-0.10	0.920	0.979
lh_G_front_sup	-0.011	-0.03, 0.009	-1.08	0.281	0.979
lh_G_Ins_lg_and_S_cent_ins	-0.003	-0.02, 0.014	-0.36	0.716	0.979
lh_G_insular_short	-0.004	-0.024, 0.017	-0.35	0.724	0.979

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_occipital_middle	-0.001	-0.025, 0.023	-0.11	0.913	0.979
lh_G_occipital_sup	-0.008	-0.033, 0.017	-0.62	0.539	0.979
lh_G_oc.temp_lat.fusifor	-0.015	-0.05, 0.019	-0.87	0.384	0.979
lh_G_oc.temp_med.Lingual	0.002	-0.022, 0.025	0.14	0.888	0.979
lh_G_oc.temp_med.Parahip	0.018	-0.016, 0.052	1.03	0.304	0.979
lh_G_orbital	0.002	-0.017, 0.021	0.18	0.860	0.979
lh_G_pariet_inf.Angular	-0.010	-0.027, 0.006	-1.25	0.213	0.979
lh_G_pariet_inf.Supramar	-0.008	-0.026, 0.01	-0.90	0.368	0.979
lh_G_parietal_sup	-0.008	-0.028, 0.011	-0.84	0.403	0.979
lh_G_postcentral	-0.008	-0.024, 0.007	-1.09	0.277	0.979
lh_G_precentral	-0.012	-0.03, 0.006	-1.33	0.184	0.979
lh_G_precuneus	0.025	-0.041, 0.09	0.73	0.463	0.979
lh_G_rectus	0.016	-0.017, 0.049	0.96	0.336	0.979
lh_G_subcallosal	-0.020	-0.041, 0.001	-1.87	0.062	0.979
lh_G_temp_sup.G_T_transv	-0.005	-0.022, 0.011	-0.66	0.512	0.979
lh_G_temp_sup.Lateral	0.008	-0.009, 0.026	0.96	0.339	0.979
lh_G_temp_sup.Plan_polar	-0.008	-0.026, 0.01	-0.90	0.371	0.979
lh_G_temp_sup.Plan_tempo	0.010	-0.019, 0.039	0.68	0.500	0.979
lh_G_temporal_inf	-0.009	-0.031, 0.012	-0.86	0.389	0.979
lh_G_temporal_middle	0.022	-0.003, 0.047	1.72	0.086	0.979
lh_Lat_Fis.ant.Horizontal	0.019	-0.009, 0.046	1.33	0.183	0.979
lh_Lat_Fis.ant.Vertical	-0.008	-0.029, 0.013	-0.74	0.462	0.979
lh_Lat_Fis.post	-0.009	-0.051, 0.033	-0.43	0.669	0.979
lh_Pole_occipital	0.022	-0.005, 0.049	1.62	0.106	0.979
lh_Pole_temporal	-0.039	-0.07, -0.007	-2.39	0.017	0.979
lh_S_calcarine	-0.013	-0.033, 0.008	-1.21	0.226	0.979
lh_S_central	-0.011	-0.028, 0.007	-1.17	0.243	0.979
lh_S_cingul.Marginalis	0.019	-0.003, 0.04	1.71	0.087	0.979
lh_S_circular_insula_ant	-0.004	-0.021, 0.013	-0.43	0.666	0.979
lh_S_circular_insula_inf	-0.004	-0.023, 0.014	-0.44	0.662	0.979
lh_S_circular_insula_sup	-0.007	-0.038, 0.024	-0.44	0.662	0.979
lh_S_collat_transv_ant	0.017	-0.018, 0.051	0.96	0.338	0.979
lh_S_collat_transv_post	0.010	-0.012, 0.032	0.87	0.385	0.979
lh_S_front_inf	0.003	-0.018, 0.025	0.31	0.757	0.979
lh_S_front_middle	0.001	-0.016, 0.017	0.10	0.919	0.979
lh_S_front_sup	0.000	-0.025, 0.024	-0.04	0.971	0.990
lh_S_interm_prim.Jensen	-0.003	-0.023, 0.017	-0.30	0.764	0.979
lh_S_intrapariet_and_P_trans	-0.017	-0.047, 0.013	-1.13	0.259	0.979
lh_S_oc_middle_and_Lunatus	-0.003	-0.026, 0.019	-0.29	0.770	0.979
lh_S_oc_sup_and_transversal	0.004	-0.016, 0.025	0.42	0.675	0.979
lh_S_occipital_ant	-0.007	-0.026, 0.013	-0.69	0.492	0.979
lh_S_oc.temp_lat	0.003	-0.017, 0.022	0.27	0.788	0.979
lh_S_oc.temp_med_and_Lingual	0.028	-0.004, 0.061	1.70	0.090	0.979

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital_lateral	0.006	-0.052, 0.064	0.22	0.828	0.979
lh_S_orbital_med.olfact	0.008	-0.019, 0.034	0.55	0.580	0.979
lh_S_orbital.H_Shaped	-0.002	-0.022, 0.018	-0.20	0.840	0.979
lh_S_parieto_occipital	0.005	-0.018, 0.028	0.44	0.658	0.979
lh_S_pericallosal	0.002	-0.015, 0.019	0.21	0.834	0.979
lh_S_postcentral	0.002	-0.02, 0.023	0.14	0.886	0.979
lh_S_precentral.inf.part	-0.007	-0.024, 0.01	-0.82	0.413	0.979
lh_S_precentral.sup.part	0.047	0, 0.094	1.99	0.048	0.979
lh_S_suborbital	0.008	-0.015, 0.03	0.66	0.512	0.979
lh_S_subparietal	-0.004	-0.035, 0.028	-0.23	0.815	0.979
lh_S_temporal_inf	0.001	-0.015, 0.018	0.13	0.899	0.979
lh_S_temporal_sup	-0.021	-0.047, 0.005	-1.62	0.106	0.979
lh_S_temporal_transverse	0.017	-0.023, 0.057	0.83	0.408	0.979
rh_G_and_S_frontomargin	0.001	-0.031, 0.033	0.06	0.951	0.990
rh_G_and_S_occipital_inf	-0.015	-0.036, 0.006	-1.40	0.163	0.979
rh_G_and_S_paracentral	-0.019	-0.04, 0.001	-1.84	0.066	0.979
rh_G_and_S_subcentral	-0.002	-0.036, 0.031	-0.14	0.885	0.979
rh_G_and_S_transv_frontopol	0.016	-0.005, 0.037	1.46	0.146	0.979
rh_G_and_S_cingul.Ant	-0.008	-0.026, 0.009	-0.93	0.350	0.979
rh_G_and_S_cingul.Mid.Ant	-0.005	-0.021, 0.011	-0.62	0.533	0.979
rh_G_and_S_cingul.Mid.Post	0.002	-0.017, 0.021	0.24	0.813	0.979
rh_G_cingul.Post.dorsal	-0.009	-0.041, 0.023	-0.55	0.582	0.979
rh_G_cingul.Post.ventral	-0.019	-0.05, 0.012	-1.21	0.226	0.979
rh_G_cuneus	-0.007	-0.024, 0.011	-0.73	0.469	0.979
rh_G_front_inf.Opercular	-0.012	-0.04, 0.017	-0.81	0.419	0.979
rh_G_front_inf.Orbital	0.002	-0.02, 0.024	0.18	0.861	0.979
rh_G_front_inf.Triangul	0.002	-0.019, 0.022	0.15	0.882	0.979
rh_G_front_middle	-0.005	-0.019, 0.009	-0.70	0.481	0.979
rh_G_front_sup	0.006	-0.015, 0.028	0.59	0.555	0.979
rh_G_Ins_lg_and_S_cent_ins	-0.005	-0.022, 0.013	-0.55	0.583	0.979
rh_G_insular_short	-0.006	-0.027, 0.015	-0.52	0.602	0.979
rh_G_occipital_middle	-0.004	-0.027, 0.019	-0.38	0.706	0.979
rh_G_occipital_sup	-0.010	-0.038, 0.017	-0.74	0.461	0.979
rh_G_oc.temp_lat.fusifor	-0.011	-0.046, 0.023	-0.64	0.524	0.979
rh_G_oc.temp_med.Lingual	0.004	-0.023, 0.03	0.28	0.780	0.979
rh_G_oc.temp_med.Parahip	0.001	-0.034, 0.035	0.03	0.977	0.990
rh_G_orbital	0.004	-0.015, 0.023	0.41	0.684	0.979
rh_G_pariet_inf.Angular	-0.015	-0.032, 0.002	-1.79	0.075	0.979
rh_G_pariet_inf.Supramar	-0.008	-0.026, 0.01	-0.84	0.400	0.979
rh_G_parietal_sup	-0.012	-0.032, 0.007	-1.26	0.208	0.979
rh_G_postcentral	-0.003	-0.018, 0.011	-0.42	0.674	0.979
rh_G_precentral	0.000	-0.019, 0.018	-0.05	0.961	0.990
rh_G_precuneus	-0.005	-0.082, 0.073	-0.12	0.904	0.979

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_rectus	-0.006	-0.039, 0.027	-0.34	0.734	0.979
rh_G_subcallosal	0.005	-0.018, 0.027	0.39	0.696	0.979
rh_G_temp_sup.G_T_transv	-0.004	-0.02, 0.012	-0.50	0.617	0.979
rh_G_temp_sup.Lateral	0.009	-0.007, 0.026	1.12	0.265	0.979
rh_G_temp_sup.Plan_polar	-0.015	-0.034, 0.004	-1.51	0.132	0.979
rh_G_temp_sup.Plan_tempo	0.010	-0.023, 0.043	0.59	0.555	0.979
rh_G_temporal_inf	-0.001	-0.021, 0.019	-0.10	0.918	0.979
rh_G_temporal_middle	-0.004	-0.028, 0.021	-0.29	0.771	0.979
rh_Lat_Fis.ant.Horizontal	0.011	-0.018, 0.04	0.74	0.460	0.979
rh_Lat_Fis.ant.Vertical	-0.002	-0.023, 0.019	-0.18	0.860	0.979
rh_Lat_Fis.post	-0.011	-0.051, 0.029	-0.53	0.595	0.979
rh_Pole_occipital	0.010	-0.024, 0.043	0.58	0.563	0.979
rh_Pole_temporal	-0.032	-0.065, 0	-1.97	0.050	0.979
rh_S_calcarine	-0.006	-0.028, 0.016	-0.52	0.600	0.979
rh_S_central	-0.012	-0.03, 0.005	-1.37	0.173	0.979
rh_S_cingul.Marginalis	-0.004	-0.024, 0.016	-0.38	0.701	0.979
rh_S_circular_insula_ant	-0.003	-0.021, 0.015	-0.34	0.734	0.979
rh_S_circular_insula_inf	0.011	-0.009, 0.03	1.10	0.272	0.979
rh_S_circular_insula_sup	0.001	-0.031, 0.033	0.07	0.943	0.990
rh_S_collat_transv_ant	-0.011	-0.044, 0.022	-0.63	0.527	0.979
rh_S_collat_transv_post	0.000	-0.022, 0.022	0.02	0.987	0.993
rh_S_front_inf	-0.007	-0.03, 0.017	-0.56	0.574	0.979
rh_S_front_middle	-0.003	-0.019, 0.014	-0.31	0.760	0.979
rh_S_front_sup	0.000	-0.022, 0.022	0.00	0.999	0.999
rh_S_interm_prim.Jensen	-0.001	-0.02, 0.019	-0.08	0.938	0.990
rh_S_intrapariet_and_P_trans	-0.016	-0.044, 0.013	-1.08	0.279	0.979
rh_S_oc_middle_and_Lunatus	-0.008	-0.031, 0.016	-0.65	0.514	0.979
rh_S_oc_sup_and_transversal	0.003	-0.016, 0.022	0.32	0.748	0.979
rh_S_occipital_ant	0.006	-0.015, 0.026	0.54	0.593	0.979
rh_S_oc.temp_lat	0.009	-0.011, 0.03	0.91	0.364	0.979
rh_S_oc.temp_med_and_Lingual	0.000	-0.03, 0.031	0.03	0.978	0.990
rh_S_orbital_lateral	0.010	-0.048, 0.068	0.33	0.745	0.979
rh_S_orbital_med.olfact	-0.011	-0.036, 0.015	-0.82	0.415	0.979
rh_S_orbital.H_Shaped	-0.002	-0.023, 0.019	-0.18	0.856	0.979
rh_S_parieto_occipital	-0.005	-0.03, 0.02	-0.39	0.695	0.979
rh_S_pericallosal	-0.001	-0.018, 0.016	-0.14	0.888	0.979
rh_S_postcentral	-0.005	-0.025, 0.015	-0.47	0.640	0.979
rh_S_precentral.inf.part	-0.012	-0.028, 0.005	-1.38	0.169	0.979
rh_S_precentral.sup.part	0.039	-0.023, 0.102	1.23	0.218	0.979
rh_S_suborbital	0.004	-0.017, 0.026	0.42	0.678	0.979
rh_S_subparietal	-0.005	-0.038, 0.028	-0.30	0.762	0.979
rh_S_temporal_inf	0.000	-0.017, 0.017	0.04	0.971	0.990
rh_S_temporal_sup	0.002	-0.026, 0.03	0.16	0.877	0.979

eTable 14. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_transverse	0.013	-0.024, 0.051	0.70	0.484	0.979

Bold denotes group differences at $p < 0.05$ uncorrected

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.008	-0.036, 0.02	-0.55	0.582	0.987
Left.Thalamus.Proper	-0.005	-0.021, 0.011	-0.65	0.514	0.987
Left.Caudate	-0.004	-0.028, 0.019	-0.36	0.717	0.987
Left.Putamen	-0.002	-0.02, 0.016	-0.21	0.834	0.987
Left.Pallidum	-0.001	-0.018, 0.016	-0.16	0.872	0.987
Brain.Stem	-0.015	-0.033, 0.003	-1.61	0.108	0.845
Left.Hippocampus	-0.007	-0.024, 0.01	-0.79	0.429	0.987
Left.Amygdala	0.006	-0.015, 0.027	0.52	0.600	0.987
Left.Accumbens.area	0.005	-0.023, 0.032	0.34	0.733	0.987
Left.VentralDC	-0.015	-0.035, 0.005	-1.49	0.136	0.845
Right.Cerebellum.Cortex	-0.019	-0.047, 0.01	-1.30	0.193	0.845
Right.Thalamus.Proper	-0.010	-0.027, 0.006	-1.22	0.223	0.885
Right.Caudate	-0.004	-0.029, 0.021	-0.32	0.750	0.987
Right.Putamen	0.004	-0.015, 0.022	0.39	0.700	0.987
Right.Pallidum	0.012	-0.005, 0.029	1.38	0.168	0.845
Right.Hippocampus	0.003	-0.013, 0.02	0.40	0.687	0.987
Right.Amygdala	-0.012	-0.033, 0.009	-1.09	0.274	0.899
Right.Accumbens.area	0.003	-0.023, 0.028	0.22	0.828	0.987
Right.VentralDC	-0.015	-0.035, 0.005	-1.51	0.133	0.845
Cortical regions					
lh_G_and_S_frontomargin	-0.042	-0.073, -0.011	-2.66	0.008	0.478
lh_G_and_S_occipital_inf	-0.005	-0.025, 0.015	-0.49	0.621	0.987
lh_G_and_S_paracentral	-0.029	-0.05, -0.007	-2.62	0.009	0.478
lh_G_and_S_subcentral	-0.026	-0.066, 0.014	-1.27	0.206	0.860
lh_G_and_S_transv_frontopol	-0.001	-0.022, 0.02	-0.12	0.903	0.987
lh_G_and_S_cingul.Ant	0.002	-0.016, 0.021	0.26	0.796	0.987
lh_G_and_S_cingul.Mid.Ant	-0.004	-0.02, 0.013	-0.43	0.668	0.987
lh_G_and_S_cingul.Mid.Post	-0.009	-0.03, 0.012	-0.84	0.403	0.987
lh_G_cingul.Post.dorsal	-0.029	-0.069, 0.01	-1.45	0.147	0.845
lh_G_cingul.Post.ventral	-0.026	-0.059, 0.007	-1.57	0.118	0.845
lh_G_cuneus	-0.009	-0.028, 0.011	-0.87	0.387	0.987
lh_G_front_inf.Opercular	0.012	-0.018, 0.042	0.80	0.426	0.987
lh_G_front_inf.Orbital	-0.007	-0.034, 0.019	-0.56	0.576	0.987

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_front_inf.Triangul	-0.006	-0.027, 0.015	-0.56	0.576	0.987
lh_G_front_middle	0.001	-0.013, 0.016	0.19	0.846	0.987
lh_G_front_sup	-0.018	-0.039, 0.002	-1.77	0.078	0.845
lh_G_Ins_lg_and_S_cent_ins	0.001	-0.017, 0.018	0.08	0.940	0.987
lh_G_insular_short	-0.017	-0.038, 0.005	-1.52	0.130	0.845
lh_G_occipital_middle	-0.003	-0.027, 0.022	-0.21	0.833	0.987
lh_G_occipital_sup	-0.031	-0.058, -0.004	-2.24	0.026	0.478
lh_G_oc.temp_lat.fusifor	-0.032	-0.066, 0.002	-1.86	0.064	0.845
lh_G_oc.temp_med.Lingual	-0.027	-0.05, -0.004	-2.32	0.021	0.478
lh_G_oc.temp_med.Parahip	-0.032	-0.068, 0.004	-1.73	0.085	0.845
lh_G_orbital	0.001	-0.019, 0.021	0.10	0.923	0.987
lh_G_pariet_inf.Angular	0.002	-0.015, 0.02	0.27	0.789	0.987
lh_G_pariet_inf.Supramar	-0.015	-0.034, 0.004	-1.58	0.115	0.845
lh_G_parietal_sup	-0.002	-0.022, 0.018	-0.19	0.849	0.987
lh_G_postcentral	-0.006	-0.022, 0.01	-0.74	0.458	0.987
lh_G_precentral	-0.016	-0.035, 0.003	-1.66	0.098	0.845
lh_G_precuneus	-0.017	-0.085, 0.051	-0.49	0.623	0.987
lh_G_rectus	-0.005	-0.039, 0.029	-0.32	0.751	0.987
lh_G_subcallosal	-0.014	-0.037, 0.009	-1.18	0.237	0.899
lh_G_temp_sup.G_T_transv	-0.002	-0.018, 0.014	-0.22	0.830	0.987
lh_G_temp_sup.Lateral	0.000	-0.018, 0.018	-0.03	0.974	0.987
lh_G_temp_sup.Plan_polar	-0.025	-0.044, -0.006	-2.60	0.010	0.478
lh_G_temp_sup.Plan_tempo	0.001	-0.032, 0.034	0.04	0.968	0.987
lh_G_temporal_inf	-0.007	-0.029, 0.015	-0.63	0.526	0.987
lh_G_temporal_middle	0.015	-0.012, 0.043	1.11	0.267	0.899
lh_Lat_Fis.ant.Horizontal	-0.001	-0.031, 0.029	-0.08	0.939	0.987
lh_Lat_Fis.ant.Vertical	-0.013	-0.036, 0.009	-1.17	0.243	0.899
lh_Lat_Fis.post	0.014	-0.026, 0.054	0.69	0.490	0.987
lh_Pole_occipital	0.010	-0.02, 0.039	0.63	0.532	0.987
lh_Pole_temporal	-0.033	-0.066, -0.001	-2.04	0.042	0.641
lh_S_calcarine	-0.012	-0.033, 0.009	-1.12	0.263	0.899
lh_S_central	-0.022	-0.04, -0.004	-2.37	0.018	0.478
lh_S_cingul.Marginalis	0.004	-0.018, 0.026	0.37	0.712	0.987
lh_S_circular_insula_ant	-0.012	-0.03, 0.006	-1.33	0.183	0.845
lh_S_circular_insula_inf	-0.004	-0.024, 0.016	-0.39	0.700	0.987
lh_S_circular_insula_sup	-0.010	-0.042, 0.021	-0.65	0.517	0.987
lh_S_collat_transv_ant	-0.024	-0.059, 0.011	-1.36	0.173	0.845
lh_S_collat_transv_post	0.010	-0.014, 0.033	0.80	0.424	0.987
lh_S_front_inf	0.001	-0.023, 0.024	0.06	0.954	0.987
lh_S_front_middle	0.000	-0.017, 0.017	-0.04	0.971	0.987
lh_S_front_sup	0.006	-0.019, 0.03	0.45	0.651	0.987
lh_S_interm_prim.Jensen	-0.001	-0.022, 0.02	-0.08	0.932	0.987
lh_S_intrapariet_and_P_trans	-0.018	-0.047, 0.012	-1.18	0.238	0.899

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_oc_middle_and_Lunatus	-0.014	-0.038, 0.01	-1.13	0.258	0.899
lh_S_oc_sup_and_transversal	0.001	-0.02, 0.022	0.07	0.941	0.987
lh_S_occipital_ant	-0.010	-0.03, 0.009	-1.02	0.310	0.954
lh_S_oc.temp_lat	-0.026	-0.046, -0.005	-2.44	0.015	0.478
lh_S_oc.temp_med_and_Lingual	0.015	-0.018, 0.048	0.88	0.382	0.987
lh_S_orbital_lateral	0.008	-0.048, 0.065	0.29	0.771	0.987
lh_S_orbital_med.olfact	-0.008	-0.037, 0.02	-0.58	0.559	0.987
lh_S_orbital.H_Shaped	-0.001	-0.023, 0.02	-0.10	0.917	0.987
lh_S_parieto_occipital	0.012	-0.009, 0.034	1.11	0.270	0.899
lh_S_pericallosal	-0.002	-0.021, 0.016	-0.27	0.790	0.987
lh_S_postcentral	0.003	-0.02, 0.025	0.24	0.810	0.987
lh_S_precentral.inf.part	-0.003	-0.02, 0.014	-0.35	0.727	0.987
lh_S_precentral.sup.part	0.015	-0.032, 0.062	0.62	0.533	0.987
lh_S_suborbital	-0.002	-0.027, 0.023	-0.17	0.864	0.987
lh_S_subparietal	-0.015	-0.045, 0.015	-1.00	0.320	0.954
lh_S_temporal_inf	-0.002	-0.021, 0.017	-0.22	0.827	0.987
lh_S_temporal_sup	-0.023	-0.051, 0.006	-1.57	0.116	0.845
lh_S_temporal_transverse	-0.028	-0.07, 0.014	-1.30	0.194	0.845
rh_G_and_S_frontomargin	-0.022	-0.053, 0.008	-1.42	0.155	0.845
rh_G_and_S_occipital_inf	-0.009	-0.03, 0.012	-0.82	0.414	0.987
rh_G_and_S_paracentral	-0.013	-0.034, 0.009	-1.16	0.248	0.899
rh_G_and_S_subcentral	-0.009	-0.043, 0.024	-0.56	0.575	0.987
rh_G_and_S_transv_frontopol	0.003	-0.018, 0.025	0.30	0.765	0.987
rh_G_and_S_cingul.Ant	0.000	-0.017, 0.018	0.02	0.982	0.988
rh_G_and_S_cingul.Mid.Ant	-0.003	-0.02, 0.013	-0.42	0.674	0.987
rh_G_and_S_cingul.Mid.Post	0.001	-0.019, 0.02	0.07	0.941	0.987
rh_G_cingul.Post.dorsal	-0.011	-0.046, 0.023	-0.65	0.519	0.987
rh_G_cingul.Post.ventral	-0.009	-0.042, 0.023	-0.57	0.569	0.987
rh_G_cuneus	0.002	-0.017, 0.022	0.24	0.812	0.987
rh_G_front_inf.Opercular	0.021	-0.01, 0.052	1.36	0.175	0.845
rh_G_front_inf.Orbital	0.011	-0.012, 0.035	0.93	0.350	0.987
rh_G_front_inf.Triangul	-0.006	-0.026, 0.015	-0.52	0.601	0.987
rh_G_front_middle	-0.005	-0.02, 0.01	-0.60	0.550	0.987
rh_G_front_sup	-0.012	-0.034, 0.011	-1.02	0.309	0.954
rh_G_Ins_lg_and_S_cent_ins	-0.002	-0.02, 0.017	-0.17	0.862	0.987
rh_G_insular_short	-0.015	-0.036, 0.006	-1.39	0.166	0.845
rh_G_occipital_middle	-0.012	-0.035, 0.011	-1.00	0.316	0.954
rh_G_occipital_sup	-0.021	-0.048, 0.005	-1.58	0.114	0.845
rh_G_oc.temp_lat.fusifor	-0.028	-0.061, 0.006	-1.61	0.109	0.845
rh_G_oc.temp_med.Lingual	-0.006	-0.033, 0.02	-0.45	0.653	0.987
rh_G_oc.temp_med.Parahip	-0.008	-0.048, 0.032	-0.40	0.691	0.987
rh_G_orbital	0.002	-0.018, 0.022	0.21	0.832	0.987
rh_G_pariet_inf.Angular	-0.001	-0.019, 0.016	-0.14	0.890	0.987

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_pariet_inf.Supramar	-0.004	-0.023, 0.015	-0.43	0.668	0.987
rh_G_parietal_sup	-0.014	-0.035, 0.007	-1.29	0.197	0.845
rh_G_postcentral	-0.001	-0.016, 0.014	-0.16	0.870	0.987
rh_G_precentral	-0.009	-0.028, 0.01	-0.89	0.373	0.987
rh_G_precuneus	0.008	-0.069, 0.086	0.21	0.831	0.987
rh_G_rectus	-0.040	-0.073, -0.006	-2.34	0.020	0.478
rh_G_subcallosal	0.003	-0.019, 0.025	0.26	0.796	0.987
rh_G_temp_sup.G_T_transv	-0.006	-0.023, 0.012	-0.65	0.518	0.987
rh_G_temp_sup.Lateral	0.009	-0.009, 0.028	1.00	0.320	0.954
rh_G_temp_sup.Plan_polar	-0.006	-0.026, 0.014	-0.58	0.565	0.987
rh_G_temp_sup.Plan_tempo	0.006	-0.027, 0.039	0.36	0.722	0.987
rh_G_temporal_inf	0.002	-0.019, 0.024	0.22	0.824	0.987
rh_G_temporal_middle	0.001	-0.024, 0.026	0.06	0.952	0.987
rh_Lat_Fis.ant.Horizontal	0.002	-0.029, 0.032	0.10	0.922	0.987
rh_Lat_Fis.ant.Vertical	-0.009	-0.033, 0.015	-0.74	0.459	0.987
rh_Lat_Fis.post	-0.016	-0.055, 0.023	-0.81	0.419	0.987
rh_Pole_occipital	-0.014	-0.048, 0.02	-0.80	0.423	0.987
rh_Pole_temporal	-0.036	-0.068, -0.004	-2.24	0.025	0.478
rh_S_calcarine	-0.015	-0.037, 0.006	-1.39	0.164	0.845
rh_S_central	-0.013	-0.032, 0.005	-1.43	0.153	0.845
rh_S_cingul.Marginalis	-0.001	-0.022, 0.02	-0.09	0.926	0.987
rh_S_circular_insula_ant	0.003	-0.017, 0.024	0.31	0.757	0.987
rh_S_circular_insula_inf	0.010	-0.011, 0.03	0.92	0.356	0.987
rh_S_circular_insula_sup	-0.009	-0.042, 0.025	-0.51	0.613	0.987
rh_S_collat_transv_ant	-0.022	-0.055, 0.011	-1.32	0.187	0.845
rh_S_collat_transv_post	0.001	-0.022, 0.024	0.09	0.930	0.987
rh_S_front_inf	0.001	-0.022, 0.024	0.10	0.917	0.987
rh_S_front_middle	-0.001	-0.019, 0.016	-0.15	0.881	0.987
rh_S_front_sup	0.002	-0.021, 0.025	0.19	0.850	0.987
rh_S_interm_prim.Jensen	0.009	-0.012, 0.029	0.85	0.397	0.987
rh_S_intrapariet_and_P_trans	-0.029	-0.057, -0.001	-2.04	0.042	0.641
rh_S_oc_middle_and_Lunatus	-0.008	-0.03, 0.014	-0.71	0.480	0.987
rh_S_oc_sup_and_transversal	-0.002	-0.022, 0.018	-0.16	0.876	0.987
rh_S_occipital_ant	0.008	-0.013, 0.029	0.79	0.432	0.987
rh_S_oc.temp_lat	-0.006	-0.027, 0.016	-0.53	0.599	0.987
rh_S_oc.temp_med_and_Lingual	0.022	-0.011, 0.055	1.31	0.192	0.845
rh_S_orbital_lateral	0.000	-0.06, 0.061	0.01	0.990	0.990
rh_S_orbital_med.olfact	-0.013	-0.041, 0.015	-0.90	0.367	0.987
rh_S_orbital.H_Shaped	0.003	-0.018, 0.025	0.31	0.755	0.987
rh_S_parieto_occipital	0.001	-0.025, 0.028	0.11	0.911	0.987
rh_S_pericallosal	-0.006	-0.023, 0.011	-0.70	0.484	0.987
rh_S_postcentral	0.005	-0.017, 0.027	0.47	0.636	0.987
rh_S_precentral.inf.part	-0.006	-0.023, 0.012	-0.65	0.515	0.987

eTable 15. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_precentral.sup.part	-0.041	-0.108, 0.025	-1.23	0.218	0.885
rh_S_suborbital	-0.002	-0.025, 0.02	-0.19	0.846	0.987
rh_S_subparietal	-0.001	-0.033, 0.032	-0.03	0.975	0.987
rh_S_temporal_inf	0.008	-0.01, 0.025	0.87	0.386	0.987
rh_S_temporal_sup	-0.006	-0.034, 0.022	-0.41	0.684	0.987
rh_S_temporal_transverse	0.032	-0.008, 0.072	1.55	0.122	0.845

Bold denotes group differences at p<0.05 uncorrected

eTable 16. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.007	-0.061, 0.046	-0.27	0.787	0.977
Left.Thalamus.Proper	-0.009	-0.039, 0.022	-0.58	0.566	0.977
Left.Caudate	-0.017	-0.057, 0.024	-0.83	0.410	0.977
Left.Putamen	-0.001	-0.034, 0.031	-0.07	0.945	0.977
Left.Pallidum	0.019	-0.016, 0.054	1.05	0.295	0.977
Brain.Stem	-0.001	-0.035, 0.033	-0.07	0.944	0.977
Left.Hippocampus	0.004	-0.03, 0.038	0.22	0.827	0.977
Left.Amygdala	0.012	-0.03, 0.053	0.55	0.584	0.977
Left.Accumbens.area	0.025	-0.029, 0.078	0.92	0.361	0.977
Left.VentralDC	-0.006	-0.044, 0.031	-0.34	0.734	0.977
Right.Cerebellum.Cortex	0.003	-0.05, 0.057	0.13	0.897	0.977
Right.Thalamus.Proper	-0.004	-0.039, 0.031	-0.25	0.804	0.977
Right.Caudate	-0.006	-0.051, 0.039	-0.26	0.799	0.977
Right.Putamen	0.010	-0.025, 0.045	0.56	0.575	0.977
Right.Pallidum	0.018	-0.018, 0.055	1.00	0.319	0.977
Right.Hippocampus	0.017	-0.015, 0.05	1.06	0.290	0.977
Right.Amygdala	0.011	-0.028, 0.05	0.57	0.567	0.977
Right.Accumbens.area	0.038	-0.009, 0.086	1.60	0.112	0.850
Right.VentralDC	0.004	-0.037, 0.045	0.21	0.832	0.977
Cortical regions					
lh_G_and_S_frontomargin	-0.009	-0.067, 0.05	-0.29	0.770	0.977
lh_G_and_S_occipital_inf	-0.019	-0.059, 0.021	-0.94	0.352	0.977
lh_G_and_S_paracentral	-0.059	-0.1, -0.019	-2.90	0.005	0.394
lh_G_and_S_subcentral	0.039	-0.039, 0.118	1.00	0.320	0.977

eTable 16. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_and_S_transv_frontopol	-0.015	-0.054, 0.023	-0.79	0.433	0.977
lh_G_and_S_cingul.Ant	-0.016	-0.051, 0.019	-0.92	0.360	0.977
lh_G_and_S_cingul.Mid.Ant	-0.022	-0.054, 0.01	-1.36	0.177	0.942
lh_G_and_S_cingul.Mid.Post	-0.002	-0.039, 0.036	-0.10	0.919	0.977
lh_G_cingul.Post.dorsal	-0.045	-0.115, 0.024	-1.29	0.201	0.958
lh_G_cingul.Post.ventral	-0.021	-0.082, 0.041	-0.67	0.505	0.977
lh_G_cuneus	-0.036	-0.071, -0.001	-2.06	0.042	0.782
lh_G_front_inf.Opercular	0.012	-0.047, 0.071	0.41	0.681	0.977
lh_G_front_inf.Orbital	-0.014	-0.065, 0.037	-0.53	0.597	0.977
lh_G_front_inf.Triangul	-0.014	-0.053, 0.025	-0.70	0.487	0.977
lh_G_front_middle	-0.001	-0.027, 0.025	-0.11	0.912	0.977
lh_G_front_sup	-0.056	-0.094, -0.017	-2.89	0.005	0.394
lh_G_Ins_lg_and_S_cent_ins	0.002	-0.03, 0.033	0.10	0.919	0.977
lh_G_insular_short	-0.011	-0.05, 0.028	-0.54	0.588	0.977
lh_G_occipital_middle	0.033	-0.012, 0.078	1.45	0.150	0.920
lh_G_occipital_sup	-0.009	-0.061, 0.044	-0.33	0.742	0.977
lh_G_oc.temp_lat.fusifor	-0.010	-0.076, 0.056	-0.31	0.757	0.977
lh_G_oc.temp_med.Lingual	-0.036	-0.082, 0.01	-1.54	0.126	0.856
lh_G_oc.temp_med.Parahip	-0.007	-0.069, 0.056	-0.21	0.834	0.977
lh_G_orbital	-0.010	-0.045, 0.026	-0.53	0.595	0.977
lh_G_pariet_inf.Angular	-0.037	-0.066, -0.007	-2.46	0.016	0.646
lh_G_pariet_inf.Supramar	-0.023	-0.056, 0.01	-1.39	0.168	0.942
lh_G_parietal_sup	-0.009	-0.046, 0.028	-0.50	0.620	0.977
lh_G_postcentral	-0.008	-0.037, 0.02	-0.57	0.568	0.977
lh_G_precentral	-0.032	-0.065, 0.002	-1.85	0.067	0.791
lh_G_precuneus	0.015	-0.115, 0.146	0.23	0.816	0.977
lh_G_rectus	0.007	-0.056, 0.07	0.22	0.827	0.977
lh_G_subcallosal	-0.042	-0.083, -0.002	-2.07	0.041	0.782
lh_G_temp_sup.G_T_transv	-0.009	-0.038, 0.02	-0.60	0.547	0.977
lh_G_temp_sup.Lateral	0.010	-0.022, 0.042	0.63	0.531	0.977
lh_G_temp_sup.Plan_polar	-0.037	-0.072, -0.002	-2.09	0.039	0.782
lh_G_temp_sup.Plan_tempo	-0.006	-0.069, 0.057	-0.19	0.848	0.977
lh_G_temporal_inf	-0.019	-0.059, 0.021	-0.93	0.356	0.977
lh_G_temporal_middle	0.026	-0.027, 0.08	0.98	0.332	0.977
lh_Lat_Fis.ant.Horizontal	-0.036	-0.095, 0.024	-1.19	0.238	0.977
lh_Lat_Fis.ant.Vertical	-0.029	-0.072, 0.014	-1.35	0.181	0.942
lh_Lat_Fis.post	0.038	-0.038, 0.113	0.99	0.323	0.977
lh_Pole_occipital	0.032	-0.022, 0.085	1.17	0.244	0.977
lh_Pole_temporal	-0.048	-0.107, 0.011	-1.62	0.109	0.850
lh_S_calcarine	-0.031	-0.068, 0.006	-1.67	0.098	0.850
lh_S_central	-0.047	-0.083, -0.011	-2.62	0.010	0.572
lh_S_cingul.Marginalis	0.007	-0.032, 0.046	0.34	0.737	0.977
lh_S_circular_insula_ant	-0.016	-0.048, 0.017	-0.96	0.338	0.977

eTable 16. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_circular_insula_inf	-0.029	-0.067, 0.009	-1.53	0.128	0.856
lh_S_circular_insula_sup	0.004	-0.055, 0.063	0.13	0.893	0.977
lh_S_collat_transv_ant	0.026	-0.037, 0.089	0.82	0.416	0.977
lh_S_collat_transv_post	0.000	-0.043, 0.042	-0.01	0.989	0.995
lh_S_front_inf	0.001	-0.046, 0.047	0.03	0.974	0.986
lh_S_front_middle	-0.008	-0.039, 0.023	-0.51	0.610	0.977
lh_S_front_sup	-0.018	-0.07, 0.033	-0.71	0.477	0.977
lh_S_interm_prim.Jensen	-0.014	-0.051, 0.024	-0.72	0.476	0.977
lh_S_intrapariet_and_P_trans	-0.017	-0.072, 0.037	-0.63	0.530	0.977
lh_S_oc_middle_and_Lunatus	-0.013	-0.059, 0.032	-0.58	0.560	0.977
lh_S_oc_sup_and_transversal	-0.012	-0.056, 0.032	-0.55	0.584	0.977
lh_S_occipital_ant	-0.010	-0.048, 0.027	-0.54	0.589	0.977
lh_S_oc.temp_lat	-0.015	-0.053, 0.024	-0.75	0.454	0.977
lh_S_oc.temp_med_and_Lingual	0.062	0.005, 0.118	2.17	0.032	0.782
lh_S_orbital_lateral	0.022	-0.099, 0.144	0.37	0.714	0.977
lh_S_orbital_med.olfact	-0.026	-0.077, 0.025	-1.03	0.308	0.977
lh_S_orbital.H_Shaped	-0.006	-0.044, 0.031	-0.35	0.731	0.977
lh_S_parieto_occipital	0.016	-0.022, 0.053	0.82	0.416	0.977
lh_S_pericallosal	-0.013	-0.048, 0.023	-0.70	0.485	0.977
lh_S_postcentral	-0.015	-0.055, 0.025	-0.74	0.459	0.977
lh_S_precentral.inf.part	-0.013	-0.046, 0.02	-0.78	0.438	0.977
lh_S_precentral.sup.part	0.021	-0.077, 0.12	0.43	0.667	0.977
lh_S_suborbital	-0.011	-0.057, 0.036	-0.46	0.646	0.977
lh_S_subparietal	0.021	-0.035, 0.077	0.74	0.458	0.977
lh_S_temporal_inf	-0.014	-0.047, 0.02	-0.82	0.411	0.977
lh_S_temporal_sup	-0.040	-0.091, 0.01	-1.58	0.118	0.856
lh_S_temporal_transverse	0.039	-0.033, 0.111	1.06	0.290	0.977
rh_G_and_S_frontomargin	-0.005	-0.062, 0.052	-0.16	0.870	0.977
rh_G_and_S_occipital_inf	-0.030	-0.072, 0.011	-1.44	0.154	0.920
rh_G_and_S_paracentral	-0.046	-0.085, -0.008	-2.38	0.019	0.646
rh_G_and_S_subcentral	0.015	-0.045, 0.075	0.49	0.622	0.977
rh_G_and_S_transv_frontopol	0.016	-0.026, 0.059	0.77	0.444	0.977
rh_G_and_S_cingul.Ant	-0.004	-0.039, 0.03	-0.24	0.807	0.977
rh_G_and_S_cingul.Mid.Ant	-0.023	-0.054, 0.008	-1.46	0.147	0.920
rh_G_and_S_cingul.Mid.Post	0.012	-0.023, 0.046	0.67	0.505	0.977
rh_G_cingul.Post.dorsal	-0.016	-0.08, 0.049	-0.48	0.632	0.977
rh_G_cingul.Post.ventral	-0.023	-0.085, 0.038	-0.75	0.455	0.977
rh_G_cuneus	-0.004	-0.037, 0.03	-0.21	0.831	0.977
rh_G_front_inf.Opercular	0.016	-0.039, 0.071	0.58	0.561	0.977
rh_G_front_inf.Orbital	0.006	-0.032, 0.043	0.30	0.762	0.977
rh_G_front_inf.Triangul	0.013	-0.026, 0.051	0.64	0.525	0.977
rh_G_front_middle	0.006	-0.019, 0.031	0.46	0.644	0.977
rh_G_front_sup	-0.015	-0.056, 0.025	-0.74	0.459	0.977


eTable 16. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_Ins_lg_and_S_cent_ins	-0.008	-0.039, 0.023	-0.52	0.606	0.977
rh_G_insular_short	-0.003	-0.04, 0.035	-0.13	0.894	0.977
rh_G_occipital_middle	-0.003	-0.048, 0.041	-0.15	0.884	0.977
rh_G_occipital_sup	-0.020	-0.071, 0.032	-0.76	0.448	0.977
rh_G_oc.temp_lat.fusifor	-0.040	-0.101, 0.021	-1.30	0.198	0.958
rh_G_oc.temp_med.Lingual	0.001	-0.047, 0.049	0.04	0.967	0.985
rh_G_oc.temp_med.Parahip	0.060	-0.014, 0.133	1.60	0.112	0.850
rh_G_orbital	0.004	-0.029, 0.037	0.24	0.809	0.977
rh_G_pariet_inf.Angular	-0.021	-0.054, 0.011	-1.29	0.200	0.958
rh_G_pariet_inf.Supramar	-0.015	-0.049, 0.02	-0.84	0.401	0.977
rh_G_parietal_sup	-0.021	-0.058, 0.016	-1.13	0.262	0.977
rh_G_postcentral	-0.004	-0.031, 0.023	-0.31	0.757	0.977
rh_G_precentral	-0.016	-0.052, 0.02	-0.88	0.379	0.977
rh_G_precuneus	-0.010	-0.141, 0.121	-0.15	0.884	0.977
rh_G_rectus	-0.036	-0.1, 0.028	-1.12	0.265	0.977
rh_G_subcallosal	0.005	-0.037, 0.047	0.23	0.817	0.977
rh_G_temp_sup.G_T_transv	0.006	-0.024, 0.037	0.42	0.677	0.977
rh_G_temp_sup.Lateral	0.030	-0.001, 0.06	1.93	0.057	0.791
rh_G_temp_sup.Plan_polar	-0.035	-0.071, 0.001	-1.90	0.060	0.791
rh_G_temp_sup.Plan_tempo	0.013	-0.045, 0.071	0.45	0.653	0.977
rh_G_temporal_inf	0.006	-0.034, 0.045	0.29	0.773	0.977
rh_G_temporal_middle	-0.006	-0.055, 0.043	-0.24	0.813	0.977
rh_Lat_Fis.ant.Horizontal	-0.009	-0.058, 0.039	-0.37	0.711	0.977
rh_Lat_Fis.ant.Vertical	-0.034	-0.076, 0.008	-1.63	0.107	0.850
rh_Lat_Fis.post	0.006	-0.065, 0.078	0.17	0.865	0.977
rh_Pole_occipital	0.060	-0.003, 0.124	1.88	0.064	0.791
rh_Pole_temporal	-0.041	-0.101, 0.019	-1.35	0.179	0.942
rh_S_calcarine	-0.039	-0.079, 0.002	-1.88	0.063	0.791
rh_S_central	-0.032	-0.068, 0.005	-1.72	0.088	0.850
rh_S_cingul.Marginalis	0.003	-0.037, 0.043	0.13	0.893	0.977
rh_S_circular_insula_ant	0.014	-0.022, 0.05	0.76	0.449	0.977
rh_S_circular_insula_inf	-0.006	-0.046, 0.034	-0.31	0.754	0.977
rh_S_circular_insula_sup	0.002	-0.06, 0.064	0.06	0.954	0.977
rh_S_collat_transv_ant	-0.002	-0.064, 0.059	-0.07	0.942	0.977
rh_S_collat_transv_post	0.002	-0.04, 0.044	0.09	0.928	0.977
rh_S_front_inf	0.005	-0.037, 0.047	0.22	0.830	0.977
rh_S_front_middle	0.002	-0.028, 0.033	0.15	0.881	0.977
rh_S_front_sup	-0.002	-0.044, 0.04	-0.08	0.936	0.977
rh_S_interm_prim.Jensen	0.000	-0.036, 0.036	0.00	0.997	0.997
rh_S_intrapariet_and_P_trans	-0.013	-0.066, 0.04	-0.48	0.629	0.977
rh_S_oc_middle_and_Lunatus	-0.004	-0.047, 0.04	-0.16	0.870	0.977
rh_S_oc_sup_and_transversal	0.017	-0.021, 0.055	0.87	0.385	0.977
rh_S_occipital_ant	-0.005	-0.041, 0.031	-0.27	0.786	0.977


eTable 16. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_oc.temp_lat	-0.022	-0.064, 0.02	-1.05	0.297	0.977
rh_S_oc.temp_med_and_Lingual	0.050	-0.007, 0.108	1.75	0.084	0.850
rh_S_orbital_lateral	0.007	-0.107, 0.122	0.13	0.897	0.977
rh_S_orbital_med.olfact	0.013	-0.031, 0.057	0.60	0.552	0.977
rh_S_orbital.H_Shaped	-0.013	-0.053, 0.028	-0.63	0.530	0.977
rh_S_parieto_occipital	0.005	-0.044, 0.053	0.19	0.852	0.977
rh_S_pericallosal	-0.016	-0.049, 0.017	-0.97	0.335	0.977
rh_S_postcentral	-0.004	-0.04, 0.032	-0.23	0.820	0.977
rh_S_precentral.inf.part	-0.016	-0.048, 0.017	-0.96	0.339	0.977
rh_S_precentral.sup.part	0.004	-0.119, 0.127	0.06	0.951	0.977
rh_S_suborbital	0.003	-0.037, 0.043	0.16	0.877	0.977
rh_S_subparietal	0.005	-0.051, 0.061	0.17	0.862	0.977
rh_S_temporal_inf	-0.004	-0.038, 0.03	-0.26	0.798	0.977
rh_S_temporal_sup	-0.013	-0.065, 0.038	-0.51	0.613	0.977
rh_S_temporal_transverse	0.064	-0.006, 0.133	1.83	0.071	0.791


Bold denotes group differences at p<0.05 uncorrected




eFigure 18. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 19 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 19 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 19 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 20. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 21 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 21 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 21 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 22. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 23 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 23 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 23 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for anticipation of loss. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Positive vs negative feedback of reward

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.011	-0.042, 0.02	-0.71	0.479	0.857
Left.Thalamus.Proper	-0.013	-0.03, 0.003	-1.59	0.114	0.805
Left.Caudate	-0.013	-0.036, 0.011	-1.06	0.290	0.835
Left.Putamen	-0.013	-0.033, 0.007	-1.31	0.190	0.805
Left.Pallidum	0.003	-0.014, 0.02	0.35	0.730	0.951
Brain.Stem	0.001	-0.019, 0.021	0.08	0.934	0.981
Left.Hippocampus	-0.006	-0.022, 0.01	-0.70	0.487	0.857
Left.Amygdala	-0.014	-0.035, 0.007	-1.35	0.178	0.805
Left.Accumbens.area	-0.011	-0.037, 0.015	-0.81	0.419	0.837
Left.VentralDC	-0.003	-0.024, 0.019	-0.24	0.812	0.958
Right.Cerebellum.Cortex	-0.025	-0.055, 0.004	-1.68	0.094	0.805
Right.Thalamus.Proper	-0.003	-0.021, 0.015	-0.28	0.780	0.951
Right.Caudate	-0.007	-0.031, 0.018	-0.54	0.588	0.903
Right.Putamen	0.000	-0.021, 0.02	-0.01	0.994	0.994
Right.Pallidum	0.003	-0.014, 0.02	0.32	0.747	0.951
Right.Hippocampus	-0.008	-0.025, 0.01	-0.87	0.387	0.837
Right.Amygdala	-0.009	-0.031, 0.014	-0.78	0.437	0.837
Right.Accumbens.area	0.000	-0.025, 0.025	0.01	0.992	0.994
Right.VentralDC	-0.016	-0.037, 0.005	-1.52	0.129	0.805
Cortical regions					
lh_G_and_S_frontomargin	-0.001	-0.046, 0.044	-0.05	0.962	0.986
lh_G_and_S_occipital_inf	0.000	-0.032, 0.033	0.02	0.986	0.994
lh_G_and_S_paracentral	-0.008	-0.031, 0.015	-0.70	0.483	0.857
lh_G_and_S_subcentral	-0.015	-0.037, 0.008	-1.27	0.205	0.805
lh_G_and_S_transv_frontopol	-0.043	-0.087, 0	-1.95	0.052	0.805
lh_G_and_S_cingul.Ant	-0.018	-0.04, 0.004	-1.63	0.105	0.805
lh_G_and_S_cingul.Mid.Ant	-0.010	-0.029, 0.01	-0.99	0.324	0.837
lh_G_and_S_cingul.Mid.Post	-0.020	-0.038, -0.001	-2.11	0.035	0.805
lh_G_cingul.Post.dorsal	-0.008	-0.029, 0.013	-0.75	0.452	0.838
lh_G_cingul.Post.ventral	0.002	-0.041, 0.045	0.09	0.928	0.980
lh_G_cuneus	-0.018	-0.05, 0.015	-1.08	0.280	0.835
lh_G_front_inf.Opercular	-0.017	-0.038, 0.004	-1.59	0.113	0.805
lh_G_front_inf.Orbital	-0.010	-0.042, 0.023	-0.58	0.559	0.899
lh_G_front_inf.Triangul	-0.007	-0.032, 0.018	-0.54	0.590	0.903
lh_G_front_middle	-0.012	-0.032, 0.009	-1.13	0.258	0.833
lh_G_front_sup	-0.007	-0.023, 0.009	-0.89	0.374	0.837
lh_G_Ins_lg_and_S_cent_ins	-0.010	-0.03, 0.01	-0.95	0.342	0.837
lh_G_insular_short	-0.003	-0.02, 0.014	-0.31	0.753	0.951

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_occipital_middle	-0.004	-0.028, 0.019	-0.38	0.706	0.951
lh_G_occipital_sup	-0.017	-0.042, 0.009	-1.28	0.203	0.805
lh_G_oc.temp_lat.fusifor	-0.021	-0.049, 0.007	-1.47	0.144	0.805
lh_G_oc.temp_med.Lingual	-0.023	-0.059, 0.013	-1.25	0.212	0.805
lh_G_oc.temp_med.Parahip	-0.008	-0.034, 0.017	-0.64	0.519	0.885
lh_G_orbital	-0.016	-0.054, 0.023	-0.80	0.424	0.837
lh_G_pariet_inf.Angular	-0.006	-0.027, 0.014	-0.61	0.543	0.899
lh_G_pariet_inf.Supramar	-0.019	-0.037, -0.001	-2.07	0.039	0.805
lh_G_parietal_sup	-0.011	-0.031, 0.01	-1.05	0.295	0.836
lh_G_postcentral	-0.021	-0.044, 0.003	-1.71	0.088	0.805
lh_G_precentral	-0.015	-0.032, 0.002	-1.76	0.079	0.805
lh_G_precuneus	-0.014	-0.034, 0.007	-1.34	0.181	0.805
lh_G_rectus	-0.005	-0.075, 0.066	-0.13	0.899	0.968
lh_G_subcallosal	-0.024	-0.061, 0.012	-1.31	0.189	0.805
lh_G_temp_sup.G_T_transv	-0.003	-0.025, 0.019	-0.28	0.778	0.951
lh_G_temp_sup.Lateral	-0.013	-0.03, 0.005	-1.40	0.164	0.805
lh_G_temp_sup.Plan_polar	-0.015	-0.032, 0.002	-1.70	0.090	0.805
lh_G_temp_sup.Plan_tempo	-0.015	-0.036, 0.006	-1.39	0.164	0.805
lh_G_temporal_inf	0.003	-0.031, 0.036	0.16	0.874	0.966
lh_G_temporal_middle	-0.005	-0.03, 0.019	-0.41	0.685	0.951
lh_Lat_Fis.ant.Horizontal	-0.018	-0.046, 0.01	-1.25	0.212	0.805
lh_Lat_Fis.ant.Vertical	-0.008	-0.039, 0.024	-0.48	0.629	0.936
lh_Lat_Fis.post	-0.016	-0.04, 0.008	-1.30	0.194	0.805
lh_Pole_occipital	-0.026	-0.068, 0.017	-1.18	0.237	0.826
lh_Pole_temporal	-0.020	-0.051, 0.011	-1.29	0.198	0.805
lh_S_calcarine	-0.005	-0.036, 0.026	-0.30	0.761	0.951
lh_S_central	-0.022	-0.048, 0.003	-1.72	0.086	0.805
lh_S_cingul.Marginalis	-0.013	-0.032, 0.006	-1.34	0.182	0.805
lh_S_circular_insula_ant	-0.010	-0.032, 0.013	-0.86	0.393	0.837
lh_S_circular_insula_inf	-0.005	-0.024, 0.014	-0.48	0.633	0.936
lh_S_circular_insula_sup	-0.017	-0.036, 0.002	-1.80	0.073	0.805
lh_S_collat_transv_ant	-0.010	-0.042, 0.022	-0.61	0.539	0.899
lh_S_collat_transv_post	-0.014	-0.044, 0.016	-0.91	0.362	0.837
lh_S_front_inf	-0.012	-0.035, 0.012	-0.97	0.334	0.837
lh_S_front_middle	-0.014	-0.038, 0.01	-1.13	0.260	0.833
lh_S_front_sup	-0.004	-0.022, 0.014	-0.46	0.646	0.946
lh_S_interm_prim.Jensen	-0.024	-0.048, 0.001	-1.92	0.055	0.805
lh_S_intrapariet_and_P_trans	-0.008	-0.029, 0.012	-0.79	0.432	0.837
lh_S_oc_middle_and_Lunatus	-0.018	-0.048, 0.012	-1.17	0.242	0.826
lh_S_oc_sup_and_transversal	-0.005	-0.028, 0.018	-0.43	0.670	0.951
lh_S_occipital_ant	-0.002	-0.022, 0.019	-0.16	0.870	0.966
lh_S_oc.temp_lat	-0.007	-0.028, 0.015	-0.61	0.545	0.899
lh_S_oc.temp_med_and_Lingual	-0.013	-0.033, 0.007	-1.28	0.202	0.805

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital_lateral	0.004	-0.033, 0.041	0.20	0.845	0.960
lh_S_orbital_med.olfact	-0.007	-0.067, 0.053	-0.24	0.813	0.958
lh_S_orbital.H_Shaped	-0.009	-0.038, 0.019	-0.65	0.517	0.885
lh_S_parieto_occipital	-0.012	-0.033, 0.009	-1.11	0.269	0.833
lh_S_pericallosal	-0.004	-0.028, 0.021	-0.29	0.775	0.951
lh_S_postcentral	-0.012	-0.031, 0.007	-1.26	0.209	0.805
lh_S_precentral.inf.part	-0.004	-0.026, 0.018	-0.35	0.726	0.951
lh_S_precentral.sup.part	0.000	-0.019, 0.018	-0.04	0.970	0.988
lh_S_suborbital	-0.009	-0.055, 0.037	-0.37	0.710	0.951
lh_S_subparietal	-0.018	-0.042, 0.005	-1.52	0.129	0.805
lh_S_temporal_inf	-0.013	-0.046, 0.021	-0.74	0.457	0.839
lh_S_temporal_sup	-0.005	-0.023, 0.013	-0.53	0.597	0.907
lh_S_temporal_transverse	0.002	-0.028, 0.032	0.15	0.880	0.966
rh_G_and_S_frontomargin	-0.051	-0.1, -0.002	-2.03	0.043	0.805
rh_G_and_S_occipital_inf	-0.028	-0.06, 0.003	-1.77	0.078	0.805
rh_G_and_S_paracentral	-0.014	-0.038, 0.009	-1.18	0.237	0.826
rh_G_and_S_subcentral	-0.012	-0.035, 0.011	-1.04	0.300	0.836
rh_G_and_S_transv_frontopol	-0.007	-0.045, 0.031	-0.38	0.704	0.951
rh_G_and_S_cingul.Ant	-0.009	-0.032, 0.014	-0.76	0.446	0.837
rh_G_and_S_cingul.Mid.Ant	-0.006	-0.024, 0.013	-0.60	0.552	0.899
rh_G_and_S_cingul.Mid.Post	-0.010	-0.028, 0.007	-1.14	0.256	0.833
rh_G_cingul.Post.dorsal	-0.003	-0.024, 0.019	-0.25	0.802	0.958
rh_G_cingul.Post.ventral	-0.004	-0.038, 0.031	-0.21	0.836	0.958
rh_G_cuneus	-0.022	-0.053, 0.009	-1.37	0.172	0.805
rh_G_front_inf.Opercular	-0.008	-0.03, 0.013	-0.77	0.442	0.837
rh_G_front_inf.Orbital	0.001	-0.032, 0.034	0.05	0.962	0.986
rh_G_front_inf.Triangul	-0.003	-0.027, 0.022	-0.22	0.830	0.958
rh_G_front_middle	-0.010	-0.032, 0.012	-0.89	0.374	0.837
rh_G_front_sup	-0.009	-0.025, 0.007	-1.11	0.266	0.833
rh_G_Ins_lg_and_S_cent_ins	-0.010	-0.031, 0.011	-0.91	0.362	0.837
rh_G_insular_short	-0.004	-0.022, 0.015	-0.39	0.696	0.951
rh_G_occipital_middle	-0.015	-0.038, 0.007	-1.35	0.179	0.805
rh_G_occipital_sup	-0.024	-0.049, 0.001	-1.86	0.064	0.805
rh_G_oc.temp_lat.fusifor	-0.002	-0.03, 0.027	-0.12	0.905	0.969
rh_G_oc.temp_med.Lingual	-0.004	-0.038, 0.031	-0.20	0.838	0.958
rh_G_oc.temp_med.Parahip	-0.005	-0.031, 0.022	-0.34	0.734	0.951
rh_G_orbital	-0.029	-0.067, 0.01	-1.48	0.141	0.805
rh_G_pariet_inf.Angular	-0.009	-0.03, 0.011	-0.89	0.373	0.837
rh_G_pariet_inf.Supramar	-0.002	-0.021, 0.016	-0.25	0.805	0.958
rh_G_parietal_sup	-0.009	-0.029, 0.01	-0.95	0.342	0.837
rh_G_postcentral	-0.011	-0.033, 0.011	-0.96	0.337	0.837
rh_G_precentral	-0.007	-0.023, 0.01	-0.78	0.438	0.837
rh_G_precuneus	-0.009	-0.03, 0.012	-0.86	0.389	0.837

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_rectus	0.029	-0.056, 0.114	0.67	0.503	0.875
rh_G_subcallosal	0.016	-0.022, 0.053	0.82	0.411	0.837
rh_G_temp_sup.G_T_transv	-0.018	-0.041, 0.005	-1.57	0.117	0.805
rh_G_temp_sup.Lateral	-0.005	-0.023, 0.013	-0.54	0.587	0.903
rh_G_temp_sup.Plan_polar	0.003	-0.013, 0.02	0.37	0.710	0.951
rh_G_temp_sup.Plan_tempo	-0.006	-0.027, 0.015	-0.58	0.560	0.899
rh_G_temporal_inf	0.022	-0.015, 0.059	1.18	0.238	0.826
rh_G_temporal_middle	-0.012	-0.034, 0.01	-1.06	0.290	0.835
rh_Lat_Fis.ant.Horizontal	-0.019	-0.046, 0.008	-1.35	0.177	0.805
rh_Lat_Fis.ant.Vertical	-0.003	-0.038, 0.032	-0.14	0.887	0.967
rh_Lat_Fis.post	-0.015	-0.039, 0.009	-1.26	0.208	0.805
rh_Pole_occipital	-0.036	-0.076, 0.003	-1.81	0.071	0.805
rh_Pole_temporal	-0.003	-0.038, 0.031	-0.18	0.856	0.966
rh_S_calcarine	-0.003	-0.035, 0.03	-0.16	0.876	0.966
rh_S_central	-0.016	-0.041, 0.009	-1.26	0.207	0.805
rh_S_cingul.Marginalis	-0.008	-0.027, 0.012	-0.79	0.433	0.837
rh_S_circular_insula_ant	0.002	-0.02, 0.023	0.14	0.891	0.967
rh_S_circular_insula_inf	-0.002	-0.022, 0.017	-0.23	0.816	0.958
rh_S_circular_insula_sup	-0.013	-0.034, 0.008	-1.23	0.221	0.818
rh_S_collat_transv_ant	-0.016	-0.051, 0.018	-0.94	0.347	0.837
rh_S_collat_transv_post	-0.013	-0.041, 0.015	-0.90	0.366	0.837
rh_S_front_inf	0.003	-0.023, 0.028	0.20	0.838	0.958
rh_S_front_middle	-0.011	-0.035, 0.014	-0.84	0.401	0.837
rh_S_front_sup	-0.004	-0.021, 0.014	-0.44	0.662	0.951
rh_S_interm_prim.Jensen	0.001	-0.022, 0.025	0.10	0.917	0.975
rh_S_intrapariet_and_P_trans	-0.001	-0.021, 0.02	-0.07	0.941	0.981
rh_S_oc_middle_and_Lunatus	-0.015	-0.042, 0.012	-1.08	0.283	0.835
rh_S_oc_sup_and_transversal	-0.004	-0.027, 0.019	-0.35	0.727	0.951
rh_S_occipital_ant	-0.005	-0.026, 0.016	-0.49	0.623	0.936
rh_S_oc.temp_lat	-0.004	-0.025, 0.016	-0.40	0.692	0.951
rh_S_oc.temp_med_and_Lingual	-0.007	-0.026, 0.012	-0.71	0.477	0.857
rh_S_orbital_lateral	0.013	-0.019, 0.046	0.81	0.420	0.837
rh_S_orbital_med.olfact	-0.025	-0.085, 0.036	-0.81	0.419	0.837
rh_S_orbital.H_Shaped	0.004	-0.023, 0.031	0.29	0.768	0.951
rh_S_parieto_occipital	-0.011	-0.032, 0.011	-0.97	0.332	0.837
rh_S_pericallosal	0.001	-0.028, 0.03	0.07	0.946	0.981
rh_S_postcentral	-0.012	-0.03, 0.006	-1.36	0.176	0.805
rh_S_precentral.inf.part	0.004	-0.019, 0.027	0.32	0.748	0.951
rh_S_precentral.sup.part	-0.003	-0.021, 0.015	-0.30	0.766	0.951
rh_S_suborbital	-0.019	-0.088, 0.049	-0.56	0.575	0.903
rh_S_subparietal	-0.007	-0.03, 0.016	-0.56	0.575	0.903
rh_S_temporal_inf	-0.039	-0.074, -0.005	-2.26	0.025	0.805
rh_S_temporal_sup	-0.004	-0.021, 0.014	-0.41	0.684	0.951

eTable 17. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_transverse	-0.014	-0.045, 0.016	-0.92	0.359	0.837

Bold denotes group differences at p<0.05 uncorrected

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.017	-0.014, 0.049	1.09	0.275	0.935
Left.Thalamus.Proper	0.001	-0.017, 0.019	0.13	0.899	0.956
Left.Caudate	-0.008	-0.031, 0.015	-0.67	0.503	0.943
Left.Putamen	-0.009	-0.032, 0.013	-0.82	0.412	0.935
Left.Pallidum	0.003	-0.015, 0.022	0.38	0.707	0.956
Brain.Stem	-0.002	-0.022, 0.019	-0.16	0.872	0.956
Left.Hippocampus	0.015	-0.003, 0.032	1.66	0.098	0.854
Left.Amygdala	0.000	-0.02, 0.021	0.04	0.967	0.978
Left.Accumbens.area	0.010	-0.016, 0.036	0.73	0.464	0.935
Left.VentralDC	0.007	-0.015, 0.029	0.65	0.514	0.943
Right.Cerebellum.Cortex	-0.002	-0.036, 0.031	-0.15	0.883	0.956
Right.Thalamus.Proper	-0.005	-0.024, 0.014	-0.53	0.594	0.946
Right.Caudate	-0.009	-0.034, 0.016	-0.72	0.470	0.935
Right.Putamen	-0.001	-0.023, 0.02	-0.13	0.900	0.956
Right.Pallidum	0.007	-0.011, 0.025	0.76	0.450	0.935
Right.Hippocampus	0.006	-0.011, 0.024	0.69	0.492	0.935
Right.Amygdala	0.014	-0.009, 0.038	1.21	0.227	0.935
Right.Accumbens.area	0.018	-0.006, 0.043	1.46	0.146	0.924
Right.VentralDC	0.019	-0.003, 0.04	1.72	0.086	0.854
Cortical regions					
lh_G_and_S_frontomargin	0.017	-0.03, 0.064	0.71	0.479	0.935
lh_G_and_S_occipital_inf	-0.002	-0.033, 0.03	-0.11	0.916	0.956
lh_G_and_S_paracentral	-0.004	-0.028, 0.02	-0.32	0.747	0.956
lh_G_and_S_subcentral	-0.021	-0.045, 0.004	-1.68	0.094	0.854
lh_G_and_S_transv_frontopol	-0.014	-0.06, 0.032	-0.61	0.540	0.946
lh_G_and_S_cingul.Ant	0.013	-0.009, 0.036	1.15	0.252	0.935
lh_G_and_S_cingul.Mid.Ant	-0.014	-0.034, 0.007	-1.29	0.198	0.935
lh_G_and_S_cingul.Mid.Post	-0.018	-0.038, 0.002	-1.74	0.082	0.854
lh_G_cingul.Post.dorsal	0.006	-0.016, 0.028	0.50	0.618	0.946
lh_G_cingul.Post.ventral	0.007	-0.038, 0.052	0.30	0.764	0.956

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_cuneus	-0.012	-0.045, 0.021	-0.74	0.460	0.935
lh_G_front_inf.Opercular	-0.012	-0.034, 0.009	-1.14	0.256	0.935
lh_G_front_inf.Orbital	-0.010	-0.042, 0.022	-0.62	0.535	0.946
lh_G_front_inf.Triangul	-0.002	-0.029, 0.025	-0.16	0.876	0.956
lh_G_front_middle	0.007	-0.015, 0.03	0.66	0.512	0.943
lh_G_front_sup	-0.005	-0.022, 0.011	-0.63	0.527	0.946
lh_G_Ins_lg_and_S_cent_ins	-0.002	-0.023, 0.02	-0.15	0.879	0.956
lh_G_insular_short	0.003	-0.016, 0.021	0.27	0.788	0.956
lh_G_occipital_middle	-0.001	-0.025, 0.022	-0.12	0.901	0.956
lh_G_occipital_sup	-0.016	-0.041, 0.01	-1.21	0.227	0.935
lh_G_oc.temp_lat.fusifor	-0.010	-0.038, 0.018	-0.70	0.487	0.935
lh_G_oc.temp_med.Lingual	-0.001	-0.037, 0.036	-0.05	0.963	0.978
lh_G_oc.temp_med.Parahip	-0.012	-0.037, 0.012	-0.98	0.329	0.935
lh_G_orbital	0.016	-0.025, 0.056	0.76	0.449	0.935
lh_G_pariet_inf.Angular	0.011	-0.011, 0.033	0.99	0.321	0.935
lh_G_pariet_inf.Supramar	-0.012	-0.031, 0.007	-1.26	0.210	0.935
lh_G_parietal_sup	-0.006	-0.026, 0.014	-0.56	0.578	0.946
lh_G_postcentral	-0.029	-0.053, -0.004	-2.32	0.021	0.698
lh_G_precentral	-0.010	-0.028, 0.009	-1.04	0.301	0.935
lh_G_precuneus	-0.001	-0.022, 0.019	-0.13	0.898	0.956
lh_G_rectus	0.016	-0.057, 0.09	0.43	0.668	0.956
lh_G_subcallosal	0.027	-0.01, 0.063	1.43	0.155	0.924
lh_G_temp_sup.G_T_transv	-0.009	-0.034, 0.016	-0.74	0.459	0.935
lh_G_temp_sup.Lateral	-0.013	-0.031, 0.006	-1.34	0.180	0.935
lh_G_temp_sup.Plan_polar	-0.007	-0.024, 0.01	-0.77	0.441	0.935
lh_G_temp_sup.Plan_tempo	-0.019	-0.042, 0.003	-1.66	0.097	0.854
lh_G_temporal_inf	-0.009	-0.043, 0.026	-0.49	0.623	0.946
lh_G_temporal_middle	-0.008	-0.031, 0.014	-0.71	0.477	0.935
lh_Lat_Fis.ant.Horizontal	-0.002	-0.03, 0.026	-0.14	0.890	0.956
lh_Lat_Fis.ant.Vertical	-0.002	-0.031, 0.028	-0.13	0.900	0.956
lh_Lat_Fis.post	-0.012	-0.038, 0.013	-0.97	0.334	0.935
lh_Pole_occipital	-0.015	-0.057, 0.027	-0.71	0.476	0.935
lh_Pole_temporal	0.015	-0.017, 0.046	0.92	0.359	0.935
lh_S_calcarine	-0.003	-0.035, 0.028	-0.21	0.834	0.956
lh_S_central	-0.036	-0.064, -0.008	-2.52	0.012	0.698
lh_S_cingul.Marginalis	-0.019	-0.04, 0.002	-1.74	0.083	0.854
lh_S_circular_insula_ant	0.014	-0.011, 0.038	1.11	0.270	0.935
lh_S_circular_insula_inf	-0.014	-0.034, 0.007	-1.32	0.187	0.935
lh_S_circular_insula_sup	-0.021	-0.041, -0.001	-2.03	0.043	0.854
lh_S_collat_transv_ant	-0.003	-0.035, 0.03	-0.15	0.878	0.956
lh_S_collat_transv_post	-0.005	-0.034, 0.025	-0.31	0.756	0.956
lh_S_front_inf	0.001	-0.024, 0.027	0.11	0.914	0.956
lh_S_front_middle	-0.021	-0.047, 0.005	-1.56	0.119	0.885

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_front_sup	0.001	-0.017, 0.02	0.15	0.882	0.956
lh_S_interm_prim.Jensen	0.006	-0.021, 0.033	0.44	0.659	0.956
lh_S_intrapariet_and_P_trans	0.008	-0.014, 0.029	0.69	0.492	0.935
lh_S_oc_middle_and_Lunatus	-0.011	-0.04, 0.019	-0.71	0.475	0.935
lh_S_oc_sup_and_transversal	0.002	-0.023, 0.027	0.13	0.900	0.956
lh_S_occipital_ant	-0.002	-0.024, 0.02	-0.18	0.856	0.956
lh_S_oc.temp_lat	-0.001	-0.022, 0.019	-0.12	0.908	0.956
lh_S_oc.temp_med_and_Lingual	-0.006	-0.026, 0.014	-0.56	0.573	0.946
lh_S_orbital_lateral	0.004	-0.034, 0.042	0.21	0.837	0.956
lh_S_orbital_med.olfact	0.016	-0.046, 0.079	0.52	0.604	0.946
lh_S_orbital.H_Shaped	0.006	-0.025, 0.036	0.36	0.716	0.956
lh_S_parieto_occipital	-0.007	-0.029, 0.015	-0.59	0.555	0.946
lh_S_pericallosal	0.010	-0.015, 0.035	0.77	0.440	0.935
lh_S_postcentral	-0.003	-0.024, 0.017	-0.30	0.761	0.956
lh_S_precentral.inf.part	0.009	-0.015, 0.033	0.75	0.454	0.935
lh_S_precentral.sup.part	-0.003	-0.022, 0.015	-0.36	0.722	0.956
lh_S_suborbital	0.003	-0.046, 0.052	0.11	0.915	0.956
lh_S_subparietal	0.007	-0.017, 0.031	0.61	0.545	0.946
lh_S_temporal_inf	-0.011	-0.044, 0.022	-0.64	0.523	0.946
lh_S_temporal_sup	-0.012	-0.03, 0.007	-1.25	0.212	0.935
lh_S_temporal_transverse	-0.012	-0.043, 0.018	-0.81	0.420	0.935
rh_G_and_S_frontomargin	-0.001	-0.052, 0.051	-0.02	0.984	0.984
rh_G_and_S_occipital_inf	0.016	-0.016, 0.047	0.98	0.329	0.935
rh_G_and_S_paracentral	-0.006	-0.03, 0.019	-0.45	0.653	0.956
rh_G_and_S_subcentral	-0.028	-0.052, -0.003	-2.20	0.028	0.783
rh_G_and_S_transv_frontopol	0.000	-0.036, 0.035	-0.03	0.979	0.984
rh_G_and_S_cingul.Ant	0.001	-0.023, 0.025	0.09	0.931	0.960
rh_G_and_S_cingul.Mid.Ant	-0.010	-0.03, 0.01	-0.97	0.335	0.935
rh_G_and_S_cingul.Mid.Post	-0.015	-0.034, 0.004	-1.54	0.125	0.885
rh_G_cingul.Post.dorsal	0.005	-0.015, 0.026	0.52	0.606	0.946
rh_G_cingul.Post.ventral	-0.007	-0.041, 0.027	-0.41	0.682	0.956
rh_G_cuneus	-0.023	-0.054, 0.009	-1.43	0.153	0.924
rh_G_front_inf.Opercular	-0.009	-0.031, 0.014	-0.75	0.455	0.935
rh_G_front_inf.Orbital	-0.003	-0.037, 0.031	-0.16	0.875	0.956
rh_G_front_inf.Triangul	-0.005	-0.03, 0.021	-0.37	0.713	0.956
rh_G_front_middle	0.002	-0.022, 0.025	0.13	0.897	0.956
rh_G_front_sup	-0.003	-0.019, 0.014	-0.30	0.766	0.956
rh_G_Ins_lg_and_S_cent_ins	-0.010	-0.034, 0.013	-0.89	0.375	0.935
rh_G_insular_short	0.002	-0.017, 0.022	0.23	0.820	0.956
rh_G_occipital_middle	0.012	-0.011, 0.034	1.02	0.308	0.935
rh_G_occipital_sup	-0.025	-0.051, 0.001	-1.91	0.057	0.854
rh_G_oc.temp_lat.fusifor	0.005	-0.024, 0.033	0.33	0.739	0.956
rh_G_oc.temp_med.Lingual	-0.026	-0.062, 0.009	-1.46	0.146	0.924

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_oc.temp_med.Parahip	-0.001	-0.03, 0.028	-0.04	0.966	0.978
rh_G_orbital	0.030	-0.009, 0.07	1.53	0.127	0.885
rh_G_pariet_inf.Angular	0.008	-0.012, 0.029	0.79	0.429	0.935
rh_G_pariet_inf.Supramar	-0.010	-0.028, 0.008	-1.07	0.286	0.935
rh_G_parietal_sup	-0.005	-0.025, 0.015	-0.53	0.598	0.946
rh_G_postcentral	-0.022	-0.045, 0.001	-1.89	0.059	0.854
rh_G_precentral	-0.006	-0.024, 0.011	-0.72	0.471	0.935
rh_G_precuneus	0.003	-0.018, 0.024	0.27	0.785	0.956
rh_G_rectus	0.061	-0.027, 0.149	1.37	0.173	0.935
rh_G_subcallosal	0.023	-0.013, 0.059	1.26	0.209	0.935
rh_G_temp_sup.G_T_transv	-0.023	-0.049, 0.002	-1.84	0.067	0.854
rh_G_temp_sup.Lateral	-0.005	-0.023, 0.014	-0.50	0.615	0.946
rh_G_temp_sup.Plan_polar	0.008	-0.01, 0.025	0.84	0.403	0.935
rh_G_temp_sup.Plan_tempo	-0.014	-0.038, 0.01	-1.12	0.265	0.935
rh_G_temporal_inf	0.021	-0.016, 0.058	1.10	0.273	0.935
rh_G_temporal_middle	-0.002	-0.025, 0.022	-0.15	0.883	0.956
rh_Lat_Fis.ant.Horizontal	0.005	-0.021, 0.032	0.40	0.686	0.956
rh_Lat_Fis.ant.Vertical	0.020	-0.014, 0.054	1.14	0.255	0.935
rh_Lat_Fis.post	-0.038	-0.064, -0.013	-2.95	0.003	0.566
rh_Pole_occipital	0.004	-0.035, 0.044	0.21	0.833	0.956
rh_Pole_temporal	-0.010	-0.047, 0.028	-0.51	0.610	0.946
rh_S_calcarine	-0.012	-0.044, 0.02	-0.76	0.449	0.935
rh_S_central	-0.033	-0.06, -0.006	-2.44	0.015	0.698
rh_S_cingul.Marginalis	-0.019	-0.038, 0	-1.96	0.051	0.854
rh_S_circular_insula_ant	0.019	-0.003, 0.041	1.67	0.095	0.854
rh_S_circular_insula_inf	-0.017	-0.038, 0.004	-1.64	0.102	0.854
rh_S_circular_insula_sup	-0.026	-0.047, -0.005	-2.40	0.017	0.698
rh_S_collat_transv_ant	-0.002	-0.038, 0.034	-0.09	0.925	0.960
rh_S_collat_transv_post	-0.006	-0.035, 0.024	-0.38	0.705	0.956
rh_S_front_inf	0.007	-0.018, 0.032	0.56	0.575	0.946
rh_S_front_middle	-0.004	-0.03, 0.022	-0.34	0.736	0.956
rh_S_front_sup	0.009	-0.01, 0.027	0.91	0.365	0.935
rh_S_interm_prim.Jensen	0.006	-0.018, 0.03	0.47	0.637	0.956
rh_S_intrapariet_and_P_trans	0.011	-0.01, 0.032	1.04	0.301	0.935
rh_S_oc_middle_and_Lunatus	0.002	-0.025, 0.03	0.18	0.860	0.956
rh_S_oc_sup_and_transversal	0.003	-0.021, 0.027	0.24	0.812	0.956
rh_S_occipital_ant	-0.009	-0.031, 0.012	-0.87	0.385	0.935
rh_S_oc.temp_lat	0.020	-0.001, 0.04	1.84	0.067	0.854
rh_S_oc.temp_med_and_Lingual	-0.012	-0.032, 0.008	-1.17	0.243	0.935
rh_S_orbital_lateral	0.003	-0.033, 0.038	0.15	0.877	0.956
rh_S_orbital_med.olfact	0.044	-0.018, 0.105	1.40	0.162	0.932
rh_S_orbital.H_Shaped	0.024	-0.006, 0.054	1.58	0.114	0.885
rh_S_parieto_occipital	-0.013	-0.035, 0.008	-1.25	0.213	0.935

eTable 18. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_pericallosal	-0.007	-0.036, 0.022	-0.46	0.646	0.956
rh_S_postcentral	-0.010	-0.029, 0.009	-1.05	0.293	0.935
rh_S_precentral.inf.part	0.006	-0.017, 0.03	0.53	0.598	0.946
rh_S_precentral.sup.part	-0.010	-0.028, 0.009	-1.05	0.296	0.935
rh_S_suborbital	-0.021	-0.095, 0.052	-0.58	0.565	0.946
rh_S_subparietal	0.008	-0.015, 0.031	0.69	0.492	0.935
rh_S_temporal_inf	0.016	-0.019, 0.051	0.89	0.375	0.935
rh_S_temporal_sup	-0.004	-0.022, 0.014	-0.43	0.669	0.956
rh_S_temporal_transverse	-0.018	-0.051, 0.014	-1.13	0.260	0.935

Bold denotes group differences at p<0.05 uncorrected

eTable 19. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.008	-0.051, 0.067	0.27	0.789	0.845
Left.Thalamus.Proper	-0.027	-0.058, 0.004	-1.70	0.093	0.310
Left.Caudate	-0.033	-0.076, 0.01	-1.54	0.126	0.335
Left.Putamen	-0.037	-0.078, 0.004	-1.80	0.075	0.280
Left.Pallidum	0.000	-0.033, 0.034	0.03	0.977	0.983
Brain.Stem	-0.021	-0.059, 0.017	-1.10	0.274	0.503
Left.Hippocampus	-0.011	-0.041, 0.019	-0.75	0.458	0.621
Left.Amygdala	-0.040	-0.074, -0.007	-2.42	0.018	0.193
Left.Accumbens.area	-0.001	-0.05, 0.047	-0.05	0.958	0.969
Left.VentralDC	-0.011	-0.055, 0.032	-0.52	0.602	0.715
Right.Cerebellum.Cortex	-0.042	-0.104, 0.02	-1.35	0.179	0.410
Right.Thalamus.Proper	-0.022	-0.055, 0.012	-1.29	0.202	0.416
Right.Caudate	-0.031	-0.078, 0.017	-1.27	0.206	0.418
Right.Putamen	-0.010	-0.048, 0.029	-0.49	0.623	0.719
Right.Pallidum	0.033	0, 0.066	1.96	0.053	0.248
Right.Hippocampus	-0.043	-0.074, -0.012	-2.72	0.008	0.174
Right.Amygdala	-0.043	-0.085, -0.002	-2.06	0.042	0.233
Right.Accumbens.area	0.018	-0.029, 0.064	0.75	0.454	0.621
Right.VentralDC	-0.018	-0.056, 0.021	-0.91	0.367	0.572
Cortical regions					
lh_G_and_S_frontomargin	-0.006	-0.097, 0.084	-0.14	0.893	0.932
lh_G_and_S_occipital_inf	-0.016	-0.08, 0.048	-0.49	0.629	0.719
lh_G_and_S_paracentral	-0.017	-0.061, 0.027	-0.76	0.448	0.621
lh_G_and_S_subcentral	-0.049	-0.091, -0.006	-2.26	0.026	0.206
lh_G_and_S_transv_frontopol	-0.122	-0.205, -0.039	-2.93	0.004	0.174

eTable 19. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_and_S_cingul.Ant	-0.007	-0.047, 0.033	-0.35	0.727	0.804
lh_G_and_S_cingul.Mid.Ant	-0.039	-0.077, -0.002	-2.08	0.040	0.233
lh_G_and_S_cingul.Mid.Post	-0.048	-0.083, -0.014	-2.76	0.007	0.174
lh_G_cingul.Post.dorsal	-0.017	-0.056, 0.023	-0.84	0.402	0.589
lh_G_cingul.Post.ventral	-0.038	-0.114, 0.037	-1.01	0.317	0.540
lh_G_cuneus	-0.077	-0.139, -0.015	-2.47	0.015	0.182
lh_G_front_inf.Opercular	-0.038	-0.079, 0.003	-1.86	0.066	0.271
lh_G_front_inf.Orbital	-0.021	-0.078, 0.036	-0.73	0.465	0.626
lh_G_front_inf.Triangul	-0.026	-0.07, 0.019	-1.15	0.252	0.490
lh_G_front_middle	-0.024	-0.062, 0.014	-1.27	0.208	0.418
lh_G_front_sup	-0.035	-0.065, -0.005	-2.35	0.021	0.193
lh_G_Ins_lg_and_S_cent_ins	-0.047	-0.087, -0.007	-2.32	0.022	0.193
lh_G_insular_short	-0.001	-0.038, 0.036	-0.06	0.950	0.968
lh_G_occipital_middle	-0.028	-0.074, 0.018	-1.21	0.231	0.453
lh_G_occipital_sup	-0.025	-0.072, 0.022	-1.05	0.298	0.533
lh_G_oc.temp_lat.fusifor	-0.038	-0.094, 0.019	-1.32	0.190	0.412
lh_G_oc.temp_med.Lingual	-0.034	-0.103, 0.034	-1.00	0.322	0.543
lh_G_oc.temp_med.Parahip	-0.026	-0.071, 0.019	-1.15	0.255	0.490
lh_G_orbital	-0.020	-0.097, 0.056	-0.53	0.599	0.715
lh_G_pariet_inf.Angular	-0.017	-0.056, 0.023	-0.84	0.406	0.589
lh_G_pariet_inf.Supramar	-0.040	-0.074, -0.006	-2.34	0.022	0.193
lh_G_parietal_sup	-0.053	-0.094, -0.013	-2.60	0.011	0.174
lh_G_postcentral	-0.062	-0.108, -0.017	-2.71	0.008	0.174
lh_G_precentral	-0.039	-0.073, -0.005	-2.31	0.023	0.193
lh_G_precuneus	-0.049	-0.087, -0.011	-2.58	0.011	0.174
lh_G_rectus	0.008	-0.121, 0.138	0.13	0.898	0.932
lh_G_subcallosal	0.007	-0.057, 0.071	0.22	0.827	0.880
lh_G_temp_sup.G_T_transv	-0.048	-0.096, -0.001	-2.03	0.045	0.233
lh_G_temp_sup.Lateral	-0.048	-0.084, -0.012	-2.63	0.010	0.174
lh_G_temp_sup.Plan_polar	-0.004	-0.035, 0.027	-0.28	0.780	0.845
lh_G_temp_sup.Plan_tempo	-0.042	-0.084, 0.001	-1.94	0.055	0.248
lh_G_temporal_inf	0.020	-0.044, 0.083	0.62	0.536	0.689
lh_G_temporal_middle	-0.037	-0.086, 0.011	-1.53	0.130	0.335
lh_Lat_Fis.ant.Horizontal	-0.044	-0.097, 0.009	-1.66	0.101	0.316
lh_Lat_Fis.ant.Vertical	-0.042	-0.101, 0.016	-1.43	0.156	0.379
lh_Lat_Fis.post	-0.041	-0.086, 0.003	-1.83	0.070	0.276
lh_Pole_occipital	-0.047	-0.121, 0.027	-1.26	0.212	0.421
lh_Pole_temporal	-0.004	-0.063, 0.054	-0.15	0.881	0.925
lh_S_calcarine	-0.031	-0.09, 0.028	-1.04	0.300	0.533
lh_S_central	-0.076	-0.126, -0.027	-3.06	0.003	0.174
lh_S_cingul.Marginalis	-0.033	-0.069, 0.003	-1.82	0.071	0.276
lh_S_circular_insula_ant	0.011	-0.034, 0.057	0.49	0.626	0.719
lh_S_circular_insula_inf	-0.018	-0.055, 0.02	-0.93	0.354	0.557

eTable 19. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_circular_insula_sup	-0.039	-0.075, -0.004	-2.19	0.031	0.208
lh_S_collat_transv_ant	0.000	-0.065, 0.065	0.00	0.998	0.998
lh_S_collat_transv_post	-0.006	-0.067, 0.056	-0.18	0.857	0.906
lh_S_front_inf	-0.038	-0.085, 0.009	-1.61	0.112	0.316
lh_S_front_middle	-0.035	-0.079, 0.008	-1.62	0.109	0.316
lh_S_front_sup	-0.015	-0.049, 0.019	-0.88	0.382	0.586
lh_S_interm_prim.Jensen	-0.013	-0.064, 0.038	-0.50	0.617	0.719
lh_S_intrapariet_and_P_trans	-0.030	-0.071, 0.011	-1.45	0.151	0.377
lh_S_oc_middle_and_Lunatus	-0.054	-0.105, -0.003	-2.11	0.038	0.233
lh_S_oc_sup_and_transversal	-0.034	-0.082, 0.013	-1.43	0.157	0.379
lh_S_occipital_ant	-0.026	-0.066, 0.014	-1.30	0.199	0.415
lh_S_oc.temp_lat	-0.016	-0.057, 0.025	-0.77	0.444	0.621
lh_S_oc.temp_med_and_Lingual	-0.015	-0.054, 0.024	-0.78	0.437	0.621
lh_S_orbital_lateral	-0.024	-0.095, 0.047	-0.66	0.509	0.664
lh_S_orbital_med.olfact	0.043	-0.069, 0.155	0.76	0.448	0.621
lh_S_orbital.H_Shaped	-0.025	-0.079, 0.03	-0.90	0.371	0.573
lh_S_parieto_occipital	-0.030	-0.068, 0.009	-1.52	0.131	0.335
lh_S_pericallosal	-0.002	-0.049, 0.046	-0.07	0.942	0.965
lh_S_postcentral	-0.043	-0.086, -0.001	-2.03	0.046	0.233
lh_S_precentral.inf.part	-0.014	-0.059, 0.031	-0.62	0.536	0.689
lh_S_precentral.sup.part	-0.013	-0.048, 0.023	-0.70	0.483	0.640
lh_S_suborbital	0.042	-0.043, 0.128	0.98	0.328	0.543
lh_S_subparietal	-0.050	-0.095, -0.005	-2.20	0.030	0.208
lh_S_temporal_inf	-0.010	-0.073, 0.053	-0.31	0.756	0.825
lh_S_temporal_sup	-0.032	-0.067, 0.002	-1.86	0.066	0.271
lh_S_temporal_transverse	-0.027	-0.088, 0.035	-0.86	0.390	0.587
rh_G_and_S_frontomargin	-0.047	-0.147, 0.052	-0.94	0.348	0.553
rh_G_and_S_occipital_inf	-0.049	-0.106, 0.008	-1.71	0.090	0.310
rh_G_and_S_paracentral	-0.026	-0.072, 0.02	-1.13	0.259	0.492
rh_G_and_S_subcentral	-0.057	-0.101, -0.014	-2.61	0.011	0.174
rh_G_and_S_transv_frontopol	-0.038	-0.106, 0.03	-1.10	0.273	0.503
rh_G_and_S_cingul.Ant	-0.010	-0.049, 0.029	-0.52	0.604	0.715
rh_G_and_S_cingul.Mid.Ant	-0.030	-0.067, 0.007	-1.63	0.107	0.316
rh_G_and_S_cingul.Mid.Post	-0.040	-0.076, -0.004	-2.22	0.029	0.208
rh_G_cingul.Post.dorsal	-0.010	-0.051, 0.03	-0.51	0.612	0.719
rh_G_cingul.Post.ventral	-0.052	-0.11, 0.005	-1.81	0.073	0.276
rh_G_cuneus	-0.053	-0.113, 0.007	-1.75	0.084	0.298
rh_G_front_inf.Opercular	-0.034	-0.073, 0.004	-1.76	0.081	0.293
rh_G_front_inf.Orbital	-0.018	-0.077, 0.042	-0.59	0.557	0.689
rh_G_front_inf.Triangul	-0.036	-0.082, 0.01	-1.54	0.127	0.335
rh_G_front_middle	-0.028	-0.07, 0.014	-1.31	0.192	0.412
rh_G_front_sup	-0.029	-0.058, -0.001	-2.02	0.046	0.233
rh_G_Ins_lg_and_S_cent_ins	-0.048	-0.091, -0.005	-2.24	0.028	0.208


eTable 19. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_insular_short	-0.007	-0.044, 0.03	-0.36	0.717	0.798
rh_G_occipital_middle	-0.021	-0.065, 0.023	-0.96	0.338	0.553
rh_G_occipital_sup	-0.060	-0.108, -0.013	-2.51	0.014	0.177
rh_G_oc.temp_lat.fusifor	-0.054	-0.109, 0.001	-1.96	0.053	0.248
rh_G_oc.temp_med.Lingual	-0.054	-0.121, 0.012	-1.62	0.110	0.316
rh_G_oc.temp_med.Parahip	-0.037	-0.092, 0.019	-1.31	0.194	0.412
rh_G_orbital	-0.026	-0.093, 0.041	-0.76	0.450	0.621
rh_G_pariet_inf.Angular	-0.042	-0.078, -0.006	-2.32	0.022	0.193
rh_G_pariet_inf.Supramar	-0.028	-0.064, 0.008	-1.56	0.122	0.335
rh_G_parietal_sup	-0.042	-0.08, -0.003	-2.16	0.033	0.213
rh_G_postcentral	-0.034	-0.077, 0.009	-1.55	0.124	0.335
rh_G_precentral	-0.014	-0.047, 0.019	-0.86	0.394	0.588
rh_G_precuneus	-0.039	-0.08, 0.002	-1.87	0.065	0.271
rh_G_rectus	0.109	-0.054, 0.273	1.33	0.186	0.412
rh_G_subcallosal	0.010	-0.062, 0.081	0.27	0.789	0.845
rh_G_temp_sup.G_T_transv	-0.034	-0.078, 0.01	-1.52	0.132	0.335
rh_G_temp_sup.Lateral	-0.045	-0.079, -0.011	-2.65	0.009	0.174
rh_G_temp_sup.Plan_polar	-0.008	-0.039, 0.023	-0.53	0.595	0.715
rh_G_temp_sup.Plan_tempo	-0.023	-0.064, 0.018	-1.12	0.266	0.499
rh_G_temporal_inf	0.028	-0.046, 0.102	0.76	0.451	0.621
rh_G_temporal_middle	-0.023	-0.069, 0.022	-1.01	0.315	0.540
rh_Lat_Fis.ant.Horizontal	-0.041	-0.092, 0.01	-1.61	0.112	0.316
rh_Lat_Fis.ant.Vertical	-0.063	-0.129, 0.004	-1.86	0.066	0.271
rh_Lat_Fis.post	-0.039	-0.086, 0.009	-1.62	0.108	0.316
rh_Pole_occipital	-0.033	-0.109, 0.044	-0.85	0.400	0.589
rh_Pole_temporal	-0.016	-0.086, 0.054	-0.45	0.656	0.740
rh_S_calcarine	-0.029	-0.088, 0.031	-0.95	0.347	0.553
rh_S_central	-0.047	-0.093, -0.001	-2.04	0.044	0.233
rh_S_cingul.Marginalis	-0.048	-0.085, -0.011	-2.55	0.012	0.174
rh_S_circular_insula_ant	0.012	-0.03, 0.054	0.59	0.557	0.689
rh_S_circular_insula_inf	-0.026	-0.064, 0.012	-1.37	0.173	0.402
rh_S_circular_insula_sup	-0.059	-0.1, -0.019	-2.92	0.004	0.174
rh_S_collat_transv_ant	0.013	-0.055, 0.081	0.38	0.707	0.792
rh_S_collat_transv_post	-0.012	-0.067, 0.043	-0.45	0.655	0.740
rh_S_front_inf	-0.014	-0.063, 0.034	-0.60	0.552	0.689
rh_S_front_middle	-0.025	-0.072, 0.023	-1.03	0.308	0.540
rh_S_front_sup	-0.023	-0.056, 0.009	-1.42	0.159	0.379
rh_S_interm_prim.Jensen	-0.031	-0.076, 0.013	-1.41	0.163	0.384
rh_S_intrapariet_and_P_trans	-0.021	-0.063, 0.021	-0.99	0.326	0.543
rh_S_oc_middle_and_Lunatus	-0.019	-0.072, 0.033	-0.73	0.470	0.628
rh_S_oc_sup_and_transversal	-0.014	-0.058, 0.031	-0.61	0.545	0.689
rh_S_occipital_ant	-0.022	-0.066, 0.021	-1.01	0.317	0.540
rh_S_oc.temp_lat	-0.012	-0.056, 0.031	-0.57	0.568	0.697


eTable 19. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for reward feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_oc.temp.med.and.Lingual	-0.019	-0.054, 0.017	-1.05	0.298	0.533
rh_S_orbital_lateral	-0.023	-0.088, 0.043	-0.69	0.493	0.649
rh_S_orbital.med.olfact	-0.020	-0.138, 0.099	-0.33	0.741	0.814
rh_S_orbital.H_Shaped	0.003	-0.046, 0.052	0.11	0.914	0.942
rh_S_parieto_occipital	-0.035	-0.077, 0.007	-1.66	0.099	0.316
rh_S_pericallosal	-0.025	-0.083, 0.033	-0.87	0.387	0.587
rh_S_postcentral	-0.034	-0.068, 0.001	-1.95	0.054	0.248
rh_S_precentral.inf.part	-0.030	-0.076, 0.015	-1.34	0.184	0.412
rh_S_precentral.sup.part	-0.023	-0.057, 0.012	-1.30	0.195	0.412
rh_S_suborbital	0.039	-0.099, 0.177	0.56	0.578	0.705
rh_S_subparietal	-0.036	-0.079, 0.006	-1.71	0.091	0.310
rh_S_temporal_inf	-0.020	-0.085, 0.045	-0.61	0.542	0.689
rh_S_temporal_sup	-0.025	-0.056, 0.006	-1.63	0.107	0.316
rh_S_temporal_transverse	-0.028	-0.085, 0.03	-0.96	0.342	0.553


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 24. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 25 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 25 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 25 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 26. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 27 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 27 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 27 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 28. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 29 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 29 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 29 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for reward feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Positive vs negative feedback of loss

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.007	-0.024, 0.037	0.43	0.669	0.996
Left.Thalamus.Proper	-0.005	-0.02, 0.011	-0.58	0.564	0.996
Left.Caudate	-0.014	-0.037, 0.009	-1.20	0.232	0.996
Left.Putamen	-0.012	-0.031, 0.007	-1.24	0.214	0.996
Left.Pallidum	-0.008	-0.025, 0.009	-0.95	0.342	0.996
Brain.Stem	0.001	-0.018, 0.02	0.08	0.937	0.996
Left.Hippocampus	0.014	-0.003, 0.03	1.64	0.101	0.996
Left.Amygdala	0.000	-0.021, 0.02	-0.02	0.984	0.996
Left.Accumbens.area	-0.002	-0.028, 0.024	-0.15	0.879	0.996
Left.VentralDC	0.002	-0.018, 0.021	0.18	0.854	0.996
Right.Cerebellum.Cortex	-0.005	-0.034, 0.024	-0.36	0.721	0.996
Right.Thalamus.Proper	-0.001	-0.017, 0.015	-0.12	0.907	0.996
Right.Caudate	-0.006	-0.029, 0.018	-0.48	0.630	0.996
Right.Putamen	-0.011	-0.031, 0.009	-1.07	0.284	0.996
Right.Pallidum	0.001	-0.016, 0.018	0.13	0.900	0.996
Right.Hippocampus	0.007	-0.01, 0.024	0.78	0.434	0.996
Right.Amygdala	0.004	-0.018, 0.025	0.35	0.726	0.996
Right.Accumbens.area	0.001	-0.023, 0.025	0.08	0.937	0.996
Right.VentralDC	0.007	-0.024, 0.037	0.43	0.669	0.996
Cortical regions					
lh_G_and_S_frontomargin	-0.002	-0.022, 0.017	-0.24	0.811	0.996
lh_G_and_S_occipital_inf	-0.033	-0.074, 0.008	-1.56	0.119	0.996
lh_G_and_S_paracentral	0.012	-0.018, 0.041	0.77	0.442	0.996
lh_G_and_S_subcentral	-0.016	-0.037, 0.004	-1.55	0.121	0.996
lh_G_and_S_transv_frontopol	-0.016	-0.038, 0.006	-1.45	0.149	0.996
lh_G_and_S_cingul.Ant	0.025	-0.017, 0.067	1.18	0.240	0.996
lh_G_and_S_cingul.Mid.Ant	0.017	-0.004, 0.039	1.59	0.113	0.996
lh_G_and_S_cingul.Mid.Post	0.004	-0.014, 0.022	0.45	0.655	0.996
lh_G_cingul.Post.dorsal	0.002	-0.015, 0.018	0.19	0.849	0.996
lh_G_cingul.Post.ventral	-0.003	-0.024, 0.018	-0.33	0.744	0.996
lh_G_cuneus	-0.013	-0.055, 0.029	-0.63	0.531	0.996
lh_G_front_inf.Opercular	-0.005	-0.036, 0.025	-0.33	0.738	0.996
lh_G_front_inf.Orbital	-0.009	-0.029, 0.01	-0.94	0.346	0.996
lh_G_front_inf.Triangul	-0.021	-0.054, 0.012	-1.26	0.207	0.996
lh_G_front_middle	0.000	-0.026, 0.027	0.01	0.989	0.996
lh_G_front_sup	-0.002	-0.022, 0.019	-0.17	0.861	0.996
lh_G_Ins_lg_and_S_cent_ins	0.000	-0.015, 0.015	0.00	0.996	0.996
lh_G_insular_short	-0.004	-0.023, 0.015	-0.44	0.659	0.996

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_occipital_middle	0.006	-0.01, 0.023	0.75	0.456	0.996
lh_G_occipital_sup	-0.007	-0.029, 0.015	-0.63	0.529	0.996
lh_G_oc.temp_lat.fusifor	-0.015	-0.04, 0.009	-1.26	0.210	0.996
lh_G_oc.temp_med.Lingual	0.006	-0.018, 0.03	0.52	0.600	0.996
lh_G_oc.temp_med.Parahip	-0.009	-0.042, 0.024	-0.52	0.607	0.996
lh_G_orbital	0.011	-0.014, 0.036	0.87	0.383	0.996
lh_G_pariet_inf.Angular	-0.016	-0.052, 0.021	-0.85	0.394	0.996
lh_G_pariet_inf.Supramar	0.012	-0.009, 0.033	1.16	0.246	0.996
lh_G_parietal_sup	-0.002	-0.02, 0.015	-0.28	0.780	0.996
lh_G_postcentral	0.007	-0.012, 0.025	0.72	0.474	0.996
lh_G_precentral	-0.009	-0.029, 0.012	-0.83	0.406	0.996
lh_G_precuneus	0.000	-0.016, 0.015	-0.03	0.972	0.996
lh_G_rectus	-0.001	-0.02, 0.019	-0.09	0.930	0.996
lh_G_subcallosal	-0.003	-0.072, 0.066	-0.09	0.926	0.996
lh_G_temp_sup.G_T_transv	-0.004	-0.041, 0.032	-0.23	0.819	0.996
lh_G_temp_sup.Lateral	-0.007	-0.028, 0.014	-0.63	0.530	0.996
lh_G_temp_sup.Plan_polar	-0.005	-0.022, 0.011	-0.62	0.534	0.996
lh_G_temp_sup.Plan_tempo	-0.008	-0.025, 0.009	-0.98	0.330	0.996
lh_G_temporal_inf	-0.014	-0.035, 0.007	-1.33	0.185	0.996
lh_G_temporal_middle	0.007	-0.024, 0.039	0.45	0.654	0.996
lh_Lat_Fis.ant.Horizontal	0.021	-0.001, 0.044	1.84	0.066	0.996
lh_Lat_Fis.ant.Vertical	-0.012	-0.039, 0.015	-0.86	0.391	0.996
lh_Lat_Fis.post	-0.001	-0.032, 0.03	-0.06	0.951	0.996
lh_Pole_occipital	-0.002	-0.024, 0.019	-0.21	0.837	0.996
lh_Pole_temporal	-0.004	-0.045, 0.038	-0.17	0.863	0.996
lh_S_calcarine	0.000	-0.029, 0.028	-0.01	0.996	0.996
lh_S_central	-0.007	-0.035, 0.022	-0.45	0.653	0.996
lh_S_cingul.Marginalis	-0.012	-0.035, 0.011	-1.03	0.304	0.996
lh_S_circular_insula_ant	0.003	-0.015, 0.021	0.30	0.766	0.996
lh_S_circular_insula_inf	0.005	-0.017, 0.028	0.46	0.647	0.996
lh_S_circular_insula_sup	0.011	-0.006, 0.028	1.30	0.195	0.996
lh_S_collat_transv_ant	-0.005	-0.024, 0.015	-0.48	0.632	0.996
lh_S_collat_transv_post	-0.017	-0.049, 0.016	-1.00	0.317	0.996
lh_S_front_inf	-0.011	-0.037, 0.015	-0.82	0.414	0.996
lh_S_front_middle	-0.003	-0.026, 0.02	-0.26	0.795	0.996
lh_S_front_sup	-0.006	-0.029, 0.018	-0.47	0.637	0.996
lh_S_interm_prim.Jensen	-0.001	-0.018, 0.016	-0.09	0.928	0.996
lh_S_intrapariet_and_P_trans	0.007	-0.018, 0.032	0.55	0.586	0.996
lh_S_oc_middle_and_Lunatus	0.008	-0.012, 0.028	0.81	0.416	0.996
lh_S_oc_sup_and_transversal	-0.004	-0.03, 0.023	-0.26	0.796	0.996
lh_S_occipital_ant	0.005	-0.018, 0.027	0.43	0.669	0.996
lh_S_oc.temp_lat	0.004	-0.016, 0.023	0.35	0.730	0.996
lh_S_oc.temp_med_and_Lingual	0.004	-0.015, 0.022	0.39	0.699	0.996

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital_lateral	0.003	-0.016, 0.022	0.30	0.762	0.996
lh_S_orbital_med.olfact	-0.012	-0.048, 0.025	-0.62	0.535	0.996
lh_S_orbital.H_Shaped	0.030	-0.031, 0.09	0.96	0.339	0.996
lh_S_parieto_occipital	-0.004	-0.032, 0.025	-0.27	0.791	0.996
lh_S_pericallosal	-0.008	-0.028, 0.012	-0.82	0.414	0.996
lh_S_postcentral	-0.009	-0.033, 0.014	-0.80	0.425	0.996
lh_S_precentral.inf.part	-0.014	-0.031, 0.003	-1.58	0.115	0.996
lh_S_precentral.sup.part	0.004	-0.018, 0.026	0.35	0.728	0.996
lh_S_suborbital	-0.002	-0.019, 0.015	-0.21	0.830	0.996
lh_S_subparietal	0.036	-0.013, 0.084	1.43	0.152	0.996
lh_S_temporal_inf	0.006	-0.017, 0.029	0.50	0.615	0.996
lh_S_temporal_sup	0.008	-0.023, 0.04	0.53	0.596	0.996
lh_S_temporal_transverse	0.004	-0.014, 0.021	0.43	0.671	0.996
rh_G_and_S_frontomargin	-0.034	-0.063, -0.004	-2.25	0.025	0.996
rh_G_and_S_occipital_inf	-0.035	-0.083, 0.013	-1.43	0.152	0.996
rh_G_and_S_paracentral	0.014	-0.016, 0.044	0.91	0.362	0.996
rh_G_and_S_subcentral	-0.001	-0.024, 0.022	-0.06	0.949	0.996
rh_G_and_S_transv_frontopol	-0.013	-0.035, 0.01	-1.08	0.279	0.996
rh_G_and_S_cingul.Ant	-0.003	-0.036, 0.03	-0.19	0.852	0.996
rh_G_and_S_cingul.Mid.Ant	0.006	-0.016, 0.028	0.52	0.605	0.996
rh_G_and_S_cingul.Mid.Post	-0.006	-0.025, 0.012	-0.68	0.495	0.996
rh_G_cingul.Post.dorsal	0.005	-0.011, 0.021	0.60	0.551	0.996
rh_G_cingul.Post.ventral	-0.002	-0.022, 0.018	-0.21	0.833	0.996
rh_G_cuneus	-0.003	-0.037, 0.031	-0.18	0.858	0.996
rh_G_front_inf.Opercular	-0.011	-0.04, 0.018	-0.71	0.475	0.996
rh_G_front_inf.Orbital	-0.001	-0.021, 0.019	-0.10	0.922	0.996
rh_G_front_inf.Triangul	-0.024	-0.055, 0.007	-1.50	0.134	0.996
rh_G_front_middle	-0.011	-0.035, 0.013	-0.93	0.353	0.996
rh_G_front_sup	-0.002	-0.024, 0.02	-0.18	0.854	0.996
rh_G_Ins_lg_and_S_cent_ins	0.001	-0.015, 0.016	0.09	0.926	0.996
rh_G_insular_short	0.007	-0.014, 0.029	0.66	0.508	0.996
rh_G_occipital_middle	0.003	-0.015, 0.021	0.32	0.748	0.996
rh_G_occipital_sup	0.012	-0.009, 0.034	1.12	0.263	0.996
rh_G_oc.temp_lat.fusifor	0.000	-0.024, 0.023	-0.04	0.968	0.996
rh_G_oc.temp_med.Lingual	0.006	-0.019, 0.032	0.49	0.627	0.996
rh_G_oc.temp_med.Parahip	-0.007	-0.037, 0.024	-0.42	0.674	0.996
rh_G_orbital	-0.011	-0.038, 0.016	-0.81	0.421	0.996
rh_G_pariet_inf.Angular	-0.021	-0.06, 0.017	-1.11	0.268	0.996
rh_G_pariet_inf.Supramar	0.002	-0.017, 0.021	0.19	0.848	0.996
rh_G_parietal_sup	-0.009	-0.026, 0.009	-0.98	0.329	0.996
rh_G_postcentral	0.003	-0.015, 0.022	0.34	0.737	0.996
rh_G_precentral	0.000	-0.021, 0.021	0.03	0.979	0.996
rh_G_precuneus	-0.011	-0.026, 0.004	-1.44	0.150	0.996

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_rectus	0.003	-0.018, 0.023	0.25	0.805	0.996
rh_G_subcallosal	0.064	-0.021, 0.149	1.49	0.138	0.996
rh_G_temp_sup.G_T_transv	0.006	-0.028, 0.04	0.35	0.729	0.996
rh_G_temp_sup.Lateral	-0.004	-0.026, 0.017	-0.41	0.682	0.996
rh_G_temp_sup.Plan_polar	-0.004	-0.02, 0.012	-0.53	0.599	0.996
rh_G_temp_sup.Plan_tempo	-0.005	-0.02, 0.011	-0.58	0.563	0.996
rh_G_temporal_inf	-0.003	-0.024, 0.019	-0.24	0.814	0.996
rh_G_temporal_middle	-0.012	-0.047, 0.022	-0.71	0.475	0.996
rh_Lat_Fis.ant.Horizontal	0.000	-0.021, 0.022	0.02	0.981	0.996
rh_Lat_Fis.ant.Vertical	-0.011	-0.036, 0.015	-0.81	0.418	0.996
rh_Lat_Fis.post	0.006	-0.028, 0.04	0.36	0.722	0.996
rh_Pole_occipital	-0.001	-0.024, 0.023	-0.06	0.948	0.996
rh_Pole_temporal	0.008	-0.033, 0.048	0.37	0.714	0.996
rh_S_calcarine	-0.028	-0.061, 0.005	-1.66	0.098	0.996
rh_S_central	-0.011	-0.041, 0.019	-0.72	0.472	0.996
rh_S_cingul.Marginalis	-0.007	-0.031, 0.017	-0.56	0.573	0.996
rh_S_circular_insula_ant	0.001	-0.017, 0.018	0.07	0.942	0.996
rh_S_circular_insula_inf	0.006	-0.017, 0.029	0.51	0.613	0.996
rh_S_circular_insula_sup	0.009	-0.009, 0.027	0.94	0.347	0.996
rh_S_collat_transv_ant	-0.006	-0.026, 0.015	-0.54	0.591	0.996
rh_S_collat_transv_post	-0.022	-0.057, 0.012	-1.26	0.207	0.996
rh_S_front_inf	0.008	-0.018, 0.034	0.59	0.555	0.996
rh_S_front_middle	0.005	-0.019, 0.028	0.40	0.689	0.996
rh_S_front_sup	-0.001	-0.025, 0.022	-0.09	0.929	0.996
rh_S_interm_prim.Jensen	0.006	-0.012, 0.023	0.61	0.543	0.996
rh_S_intrapariet_and_P_trans	0.007	-0.016, 0.031	0.63	0.531	0.996
rh_S_oc_middle_and_Lunatus	-0.002	-0.02, 0.017	-0.20	0.844	0.996
rh_S_oc_sup_and_transversal	-0.001	-0.027, 0.025	-0.09	0.928	0.996
rh_S_occipital_ant	0.003	-0.018, 0.024	0.30	0.765	0.996
rh_S_oc.temp_lat	-0.011	-0.031, 0.008	-1.12	0.263	0.996
rh_S_oc.temp_med_and_Lingual	0.005	-0.015, 0.024	0.47	0.636	0.996
rh_S_orbital_lateral	0.013	-0.006, 0.032	1.35	0.179	0.996
rh_S_orbital_med.olfact	-0.008	-0.04, 0.023	-0.52	0.606	0.996
rh_S_orbital.H_Shaped	-0.014	-0.08, 0.052	-0.41	0.681	0.996
rh_S_parieto_occipital	-0.008	-0.037, 0.022	-0.52	0.601	0.996
rh_S_pericallosal	0.007	-0.012, 0.027	0.73	0.465	0.996
rh_S_postcentral	0.002	-0.024, 0.027	0.13	0.899	0.996
rh_S_precentral.inf.part	-0.010	-0.027, 0.007	-1.10	0.273	0.996
rh_S_precentral.sup.part	0.001	-0.021, 0.023	0.06	0.948	0.996
rh_S_suborbital	-0.002	-0.02, 0.015	-0.28	0.780	0.996
rh_S_subparietal	0.049	-0.02, 0.118	1.41	0.161	0.996
rh_S_temporal_inf	-0.002	-0.025, 0.02	-0.21	0.831	0.996
rh_S_temporal_sup	0.001	-0.032, 0.034	0.07	0.948	0.996

eTable 20. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_transverse	0.009	-0.008, 0.026	1.01	0.312	0.996

Bold denotes group differences at $p < 0.05$ uncorrected

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.021	-0.011, 0.053	1.27	0.206	0.840
Left.Thalamus.Proper	0.007	-0.01, 0.024	0.83	0.407	0.840
Left.Caudate	0.013	-0.009, 0.035	1.14	0.257	0.840
Left.Putamen	0.006	-0.014, 0.026	0.59	0.558	0.840
Left.Pallidum	-0.001	-0.019, 0.016	-0.14	0.886	0.954
Brain.Stem	0.006	-0.014, 0.027	0.60	0.550	0.840
Left.Hippocampus	0.012	-0.005, 0.029	1.37	0.172	0.840
Left.Amygdala	0.008	-0.014, 0.031	0.75	0.455	0.840
Left.Accumbens.area	0.011	-0.013, 0.035	0.88	0.377	0.840
Left.VentralDC	0.009	-0.012, 0.029	0.83	0.409	0.840
Right.Cerebellum.Cortex	0.010	-0.02, 0.041	0.66	0.508	0.840
Right.Thalamus.Proper	0.016	-0.001, 0.033	1.83	0.068	0.840
Right.Caudate	0.015	-0.007, 0.037	1.36	0.174	0.840
Right.Putamen	0.010	-0.011, 0.03	0.93	0.353	0.840
Right.Pallidum	-0.002	-0.019, 0.015	-0.20	0.843	0.931
Right.Hippocampus	0.014	-0.004, 0.032	1.56	0.120	0.840
Right.Amygdala	0.019	-0.004, 0.042	1.61	0.107	0.840
Right.Accumbens.area	0.019	-0.007, 0.044	1.42	0.157	0.840
Right.VentralDC	0.002	-0.019, 0.022	0.18	0.854	0.932
Cortical regions					
lh_G_and_S_frontomargin	0.006	-0.04, 0.051	0.25	0.804	0.911
lh_G_and_S_occipital_inf	0.008	-0.024, 0.039	0.48	0.628	0.850
lh_G_and_S_paracentral	-0.005	-0.026, 0.016	-0.46	0.645	0.853
lh_G_and_S_subcentral	-0.010	-0.033, 0.014	-0.83	0.408	0.840
lh_G_and_S_transv_frontopol	0.022	-0.023, 0.067	0.96	0.336	0.840
lh_G_and_S_cingul.Ant	0.011	-0.01, 0.032	1.00	0.318	0.840
lh_G_and_S_cingul.Mid.Ant	0.009	-0.01, 0.028	0.89	0.374	0.840
lh_G_and_S_cingul.Mid.Post	0.001	-0.017, 0.019	0.11	0.914	0.954
lh_G_cingul.Post.dorsal	0.002	-0.017, 0.022	0.24	0.808	0.911
lh_G_cingul.Post.ventral	0.006	-0.038, 0.049	0.26	0.797	0.911
lh_G_cuneus	-0.013	-0.041, 0.015	-0.93	0.351	0.840
lh_G_front_inf.Opercular	0.007	-0.014, 0.027	0.64	0.521	0.840

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_front_inf.Orbital	0.028	-0.005, 0.061	1.67	0.096	0.840
lh_G_front_inf.Triangul	0.033	0.006, 0.059	2.41	0.017	0.554
lh_G_front_middle	-0.003	-0.023, 0.018	-0.29	0.775	0.911
lh_G_front_sup	0.005	-0.011, 0.02	0.59	0.555	0.840
lh_G_Ins_lg_and_S_cent_ins	0.006	-0.014, 0.026	0.57	0.568	0.840
lh_G_insular_short	0.014	-0.004, 0.031	1.55	0.122	0.840
lh_G_occipital_middle	-0.008	-0.028, 0.013	-0.72	0.473	0.840
lh_G_occipital_sup	-0.014	-0.037, 0.008	-1.27	0.206	0.840
lh_G_oc.temp_lat.fusifor	-0.001	-0.027, 0.024	-0.09	0.930	0.954
lh_G_oc.temp_med.Lingual	-0.006	-0.038, 0.026	-0.37	0.709	0.903
lh_G_oc.temp_med.Parahip	0.004	-0.021, 0.028	0.31	0.760	0.911
lh_G_orbital	0.019	-0.019, 0.057	0.97	0.333	0.840
lh_G_pariet_inf.Angular	0.016	-0.004, 0.036	1.53	0.126	0.840
lh_G_pariet_inf.Supramar	0.002	-0.015, 0.02	0.25	0.800	0.911
lh_G_parietal_sup	0.005	-0.014, 0.025	0.54	0.588	0.850
lh_G_postcentral	0.009	-0.012, 0.03	0.87	0.386	0.840
lh_G_precentral	0.004	-0.012, 0.02	0.53	0.598	0.850
lh_G_precuneus	-0.006	-0.024, 0.013	-0.62	0.538	0.840
lh_G_rectus	-0.014	-0.083, 0.055	-0.40	0.689	0.886
lh_G_subcallosal	0.041	0.004, 0.078	2.20	0.028	0.673
lh_G_temp_sup.G_T_transv	-0.015	-0.039, 0.009	-1.23	0.218	0.840
lh_G_temp_sup.Lateral	0.004	-0.012, 0.021	0.50	0.615	0.850
lh_G_temp_sup.Plan_polar	-0.009	-0.028, 0.01	-0.95	0.344	0.840
lh_G_temp_sup.Plan_tempo	0.001	-0.02, 0.022	0.09	0.927	0.954
lh_G_temporal_inf	0.004	-0.029, 0.038	0.25	0.805	0.911
lh_G_temporal_middle	0.040	0.017, 0.062	3.45	0.001	0.104
lh_Lat_Fis.ant.Horizontal	0.021	-0.007, 0.049	1.48	0.139	0.840
lh_Lat_Fis.ant.Vertical	-0.005	-0.038, 0.027	-0.32	0.752	0.911
lh_Lat_Fis.post	-0.002	-0.026, 0.023	-0.13	0.900	0.954
lh_Pole_occipital	0.002	-0.041, 0.044	0.08	0.936	0.954
lh_Pole_temporal	0.010	-0.021, 0.041	0.64	0.524	0.840
lh_S_calcarine	-0.011	-0.04, 0.018	-0.74	0.462	0.840
lh_S_central	-0.019	-0.044, 0.006	-1.47	0.142	0.840
lh_S_cingul.Marginalis	-0.008	-0.028, 0.011	-0.87	0.384	0.840
lh_S_circular_insula_ant	0.005	-0.018, 0.028	0.46	0.649	0.853
lh_S_circular_insula_inf	-0.001	-0.02, 0.018	-0.13	0.900	0.954
lh_S_circular_insula_sup	-0.006	-0.027, 0.015	-0.58	0.565	0.840
lh_S_collat_transv_ant	0.012	-0.023, 0.047	0.67	0.501	0.840
lh_S_collat_transv_post	-0.007	-0.033, 0.019	-0.51	0.613	0.850
lh_S_front_inf	0.010	-0.014, 0.033	0.79	0.428	0.840
lh_S_front_middle	-0.006	-0.03, 0.018	-0.48	0.629	0.850
lh_S_front_sup	0.005	-0.013, 0.022	0.53	0.594	0.850
lh_S_interm_prim.Jensen	0.033	0.008, 0.058	2.62	0.009	0.385

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_intrapariet_and_P_trans	0.003	-0.018, 0.023	0.27	0.787	0.911
lh_S_oc_middle_and_Lunatus	-0.001	-0.026, 0.024	-0.08	0.937	0.954
lh_S_oc_sup_and_transversal	-0.001	-0.023, 0.021	-0.11	0.911	0.954
lh_S_occipital_ant	0.007	-0.013, 0.028	0.69	0.488	0.840
lh_S_oc.temp_lat	0.012	-0.007, 0.031	1.20	0.231	0.840
lh_S_oc.temp_med_and_Lingual	-0.006	-0.025, 0.012	-0.68	0.494	0.840
lh_S_orbital_lateral	0.023	-0.015, 0.06	1.19	0.233	0.840
lh_S_orbital_med.olfact	0.047	-0.011, 0.105	1.59	0.114	0.840
lh_S_orbital.H_Shaped	0.006	-0.023, 0.034	0.41	0.685	0.886
lh_S_parieto_occipital	-0.006	-0.026, 0.013	-0.64	0.520	0.840
lh_S_pericallosal	-0.004	-0.027, 0.019	-0.32	0.747	0.911
lh_S_postcentral	0.007	-0.012, 0.025	0.71	0.479	0.840
lh_S_precentral.inf.part	0.013	-0.009, 0.034	1.15	0.251	0.840
lh_S_precentral.sup.part	0.005	-0.012, 0.023	0.58	0.562	0.840
lh_S_suborbital	0.012	-0.038, 0.062	0.46	0.643	0.853
lh_S_subparietal	0.008	-0.015, 0.032	0.72	0.474	0.840
lh_S_temporal_inf	0.001	-0.03, 0.032	0.06	0.952	0.963
lh_S_temporal_sup	0.025	0.006, 0.043	2.63	0.009	0.385
lh_S_temporal_transverse	-0.017	-0.048, 0.014	-1.07	0.285	0.840
rh_G_and_S_frontomargin	-0.012	-0.058, 0.034	-0.51	0.613	0.850
rh_G_and_S_occipital_inf	-0.001	-0.03, 0.029	-0.05	0.962	0.968
rh_G_and_S_paracentral	0.006	-0.017, 0.029	0.48	0.630	0.850
rh_G_and_S_subcentral	0.000	-0.024, 0.024	0.01	0.996	0.996
rh_G_and_S_transv_frontopol	-0.006	-0.04, 0.028	-0.34	0.733	0.911
rh_G_and_S_cingul.Ant	0.009	-0.014, 0.032	0.78	0.436	0.840
rh_G_and_S_cingul.Mid.Ant	-0.003	-0.022, 0.016	-0.28	0.778	0.911
rh_G_and_S_cingul.Mid.Post	0.004	-0.013, 0.022	0.48	0.631	0.850
rh_G_cingul.Post.dorsal	0.014	-0.005, 0.032	1.46	0.144	0.840
rh_G_cingul.Post.ventral	0.012	-0.022, 0.047	0.70	0.484	0.840
rh_G_cuneus	-0.021	-0.049, 0.007	-1.49	0.138	0.840
rh_G_front_inf.Opercular	0.009	-0.012, 0.03	0.82	0.412	0.840
rh_G_front_inf.Orbital	0.016	-0.015, 0.048	1.04	0.299	0.840
rh_G_front_inf.Triangul	0.001	-0.022, 0.024	0.08	0.937	0.954
rh_G_front_middle	0.017	-0.004, 0.038	1.61	0.108	0.840
rh_G_front_sup	0.001	-0.014, 0.016	0.14	0.886	0.954
rh_G_Ins_lg_and_S_cent_ins	0.004	-0.019, 0.026	0.34	0.735	0.911
rh_G_insular_short	0.007	-0.012, 0.026	0.76	0.446	0.840
rh_G_occipital_middle	0.010	-0.011, 0.031	0.94	0.347	0.840
rh_G_occipital_sup	-0.014	-0.036, 0.008	-1.23	0.221	0.840
rh_G_oc.temp_lat.fusifor	-0.007	-0.034, 0.02	-0.52	0.604	0.850
rh_G_oc.temp_med.Lingual	-0.015	-0.045, 0.016	-0.94	0.347	0.840
rh_G_oc.temp_med.Parahip	0.014	-0.013, 0.041	0.99	0.324	0.840
rh_G_orbital	0.024	-0.013, 0.06	1.27	0.203	0.840

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_pariet_inf.Angular	0.008	-0.012, 0.027	0.79	0.431	0.840
rh_G_pariet_inf.Supramar	0.005	-0.012, 0.023	0.61	0.544	0.840
rh_G_parietal_sup	0.006	-0.012, 0.025	0.69	0.493	0.840
rh_G_postcentral	0.009	-0.011, 0.029	0.88	0.377	0.840
rh_G_precentral	-0.002	-0.018, 0.013	-0.27	0.785	0.911
rh_G_precuneus	-0.006	-0.024, 0.013	-0.60	0.548	0.840
rh_G_rectus	0.027	-0.061, 0.115	0.61	0.541	0.840
rh_G_subcallosal	0.011	-0.023, 0.045	0.63	0.530	0.840
rh_G_temp_sup.G_T_transv	-0.008	-0.031, 0.015	-0.69	0.491	0.840
rh_G_temp_sup.Lateral	0.006	-0.01, 0.023	0.76	0.450	0.840
rh_G_temp_sup.Plan_polar	-0.008	-0.026, 0.009	-0.92	0.356	0.840
rh_G_temp_sup.Plan_tempo	-0.002	-0.024, 0.02	-0.19	0.847	0.931
rh_G_temporal_inf	0.015	-0.019, 0.05	0.88	0.377	0.840
rh_G_temporal_middle	0.003	-0.018, 0.024	0.28	0.782	0.911
rh_Lat_Fis.ant.Horizontal	0.011	-0.015, 0.038	0.85	0.398	0.840
rh_Lat_Fis.ant.Vertical	0.056	0.021, 0.09	3.18	0.002	0.131
rh_Lat_Fis.post	0.003	-0.023, 0.029	0.21	0.832	0.931
rh_Pole_occipital	0.029	-0.01, 0.068	1.46	0.146	0.840
rh_Pole_temporal	0.017	-0.018, 0.051	0.94	0.346	0.840
rh_S_calcarine	-0.013	-0.041, 0.015	-0.89	0.373	0.840
rh_S_central	-0.012	-0.036, 0.013	-0.93	0.355	0.840
rh_S_cingul.Marginalis	-0.007	-0.026, 0.011	-0.79	0.430	0.840
rh_S_circular_insula_ant	0.021	-0.002, 0.045	1.82	0.069	0.840
rh_S_circular_insula_inf	-0.006	-0.025, 0.013	-0.61	0.544	0.840
rh_S_circular_insula_sup	-0.012	-0.032, 0.009	-1.08	0.280	0.840
rh_S_collat_transv_ant	0.005	-0.031, 0.041	0.29	0.775	0.911
rh_S_collat_transv_post	-0.013	-0.04, 0.014	-0.98	0.329	0.840
rh_S_front_inf	0.011	-0.012, 0.033	0.91	0.361	0.840
rh_S_front_middle	-0.011	-0.033, 0.011	-0.98	0.329	0.840
rh_S_front_sup	0.009	-0.007, 0.025	1.09	0.275	0.840
rh_S_interm_prim.Jensen	0.008	-0.015, 0.031	0.66	0.511	0.840
rh_S_intrapariet_and_P_trans	0.002	-0.017, 0.021	0.25	0.803	0.911
rh_S_oc_middle_and_Lunatus	0.013	-0.011, 0.038	1.08	0.283	0.840
rh_S_oc_sup_and_transversal	-0.013	-0.033, 0.008	-1.23	0.218	0.840
rh_S_occipital_ant	0.012	-0.008, 0.032	1.21	0.227	0.840
rh_S_oc.temp_lat	0.015	-0.004, 0.035	1.55	0.122	0.840
rh_S_oc.temp_med_and_Lingual	0.008	-0.011, 0.028	0.81	0.420	0.840
rh_S_orbital_lateral	-0.017	-0.05, 0.015	-1.05	0.294	0.840
rh_S_orbital_med.olfact	0.031	-0.032, 0.094	0.97	0.332	0.840
rh_S_orbital.H_Shaped	0.019	-0.011, 0.048	1.25	0.212	0.840
rh_S_parieto_occipital	-0.004	-0.025, 0.016	-0.42	0.674	0.880
rh_S_pericallosal	0.027	0.004, 0.051	2.27	0.024	0.669
rh_S_postcentral	0.010	-0.008, 0.028	1.12	0.263	0.840

eTable 21. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_precentral.inf.part	0.008	-0.014, 0.031	0.73	0.464	0.840
rh_S_precentral.sup.part	0.009	-0.008, 0.026	1.04	0.298	0.840
rh_S_suborbital	0.030	-0.039, 0.099	0.86	0.390	0.840
rh_S_subparietal	-0.002	-0.024, 0.02	-0.20	0.843	0.931
rh_S_temporal_inf	0.016	-0.019, 0.05	0.89	0.372	0.840
rh_S_temporal_sup	0.006	-0.011, 0.023	0.74	0.462	0.840
rh_S_temporal_transverse	-0.019	-0.05, 0.011	-1.24	0.217	0.840

Bold denotes group differences at p<0.05 uncorrected

eTable 22. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.038	-0.027, 0.102	1.16	0.247	0.989
Left.Thalamus.Proper	0.008	-0.021, 0.037	0.53	0.600	0.989
Left.Caudate	-0.002	-0.04, 0.036	-0.11	0.916	0.989
Left.Putamen	-0.004	-0.037, 0.028	-0.27	0.791	0.989
Left.Pallidum	-0.012	-0.042, 0.018	-0.78	0.435	0.989
Brain.Stem	0.001	-0.034, 0.036	0.07	0.941	0.989
Left.Hippocampus	0.004	-0.026, 0.034	0.27	0.785	0.989
Left.Amygdala	-0.005	-0.044, 0.035	-0.24	0.810	0.989
Left.Accumbens.area	0.009	-0.036, 0.054	0.39	0.695	0.989
Left.VentralDC	0.016	-0.023, 0.055	0.81	0.420	0.989
Right.Cerebellum.Cortex	0.026	-0.031, 0.083	0.91	0.367	0.989
Right.Thalamus.Proper	0.018	-0.013, 0.049	1.17	0.246	0.989
Right.Caudate	-0.014	-0.056, 0.028	-0.64	0.521	0.989
Right.Putamen	0.002	-0.034, 0.038	0.11	0.914	0.989
Right.Pallidum	-0.028	-0.06, 0.005	-1.67	0.098	0.989
Right.Hippocampus	0.013	-0.018, 0.045	0.85	0.397	0.989
Right.Amygdala	0.006	-0.037, 0.05	0.29	0.771	0.989
Right.Accumbens.area	0.022	-0.025, 0.069	0.92	0.362	0.989

eTable 22. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Right.VentralDC	0.012	-0.025, 0.049	0.65	0.516	0.989
Cortical regions					
lh_G_and_S_frontomargin	-0.047	-0.128, 0.033	-1.18	0.243	0.989
lh_G_and_S_occipital_inf	0.027	-0.035, 0.088	0.86	0.392	0.989
lh_G_and_S_paracentral	-0.013	-0.055, 0.029	-0.61	0.541	0.989
lh_G_and_S_subcentral	-0.033	-0.073, 0.007	-1.63	0.107	0.989
lh_G_and_S_transv_frontopol	0.007	-0.072, 0.086	0.17	0.866	0.989
lh_G_and_S_cingul.Ant	0.022	-0.016, 0.059	1.16	0.249	0.989
lh_G_and_S_cingul.Mid.Ant	0.011	-0.024, 0.045	0.61	0.541	0.989
lh_G_and_S_cingul.Mid.Post	0.008	-0.023, 0.038	0.49	0.626	0.989
lh_G_cingul.Post.dorsal	-0.006	-0.046, 0.033	-0.31	0.758	0.989
lh_G_cingul.Post.ventral	-0.023	-0.106, 0.06	-0.56	0.579	0.989
lh_G_cuneus	-0.022	-0.078, 0.034	-0.77	0.443	0.989
lh_G_front_inf.Opercular	-0.015	-0.054, 0.024	-0.78	0.439	0.989
lh_G_front_inf.Orbital	0.011	-0.053, 0.076	0.35	0.729	0.989
lh_G_front_inf.Triangul	0.014	-0.039, 0.067	0.54	0.589	0.989
lh_G_front_middle	-0.034	-0.071, 0.004	-1.79	0.077	0.989
lh_G_front_sup	0.001	-0.027, 0.03	0.10	0.923	0.989
lh_G_Ins_lg_and_S_cent_ins	-0.001	-0.04, 0.038	-0.07	0.946	0.989
lh_G_insular_short	0.020	-0.014, 0.054	1.14	0.255	0.989
lh_G_occipital_middle	-0.012	-0.051, 0.027	-0.61	0.541	0.989
lh_G_occipital_sup	-0.022	-0.063, 0.019	-1.05	0.294	0.989
lh_G_oc.temp_lat.fusifor	0.002	-0.048, 0.051	0.07	0.943	0.989
lh_G_oc.temp_med.Lingual	-0.005	-0.064, 0.054	-0.17	0.863	0.989
lh_G_oc.temp_med.Parahip	0.016	-0.032, 0.064	0.67	0.505	0.989
lh_G_orbital	0.009	-0.066, 0.084	0.24	0.809	0.989
lh_G_pariet_inf.Angular	0.016	-0.025, 0.057	0.75	0.453	0.989
lh_G_pariet_inf.Supramar	-0.005	-0.036, 0.027	-0.29	0.772	0.989
lh_G_parietal_sup	-0.005	-0.039, 0.028	-0.32	0.752	0.989
lh_G_postcentral	-0.010	-0.043, 0.023	-0.59	0.559	0.989
lh_G_precentral	0.001	-0.025, 0.028	0.11	0.914	0.989
lh_G_precuneus	-0.009	-0.045, 0.028	-0.47	0.643	0.989
lh_G_rectus	-0.018	-0.143, 0.107	-0.28	0.776	0.989
lh_G_subcallosal	0.033	-0.036, 0.102	0.94	0.349	0.989
lh_G_temp_sup.G_T_transv	0.003	-0.038, 0.043	0.13	0.893	0.989
lh_G_temp_sup.Lateral	-0.006	-0.036, 0.024	-0.40	0.691	0.989
lh_G_temp_sup.Plan_polar	-0.012	-0.044, 0.02	-0.74	0.461	0.989
lh_G_temp_sup.Plan_tempo	-0.002	-0.043, 0.039	-0.10	0.923	0.989
lh_G_temporal_inf	0.011	-0.051, 0.072	0.34	0.736	0.989
lh_G_temporal_middle	0.046	-0.001, 0.092	1.95	0.054	0.989
lh_Lat_Fis.ant.Horizontal	-0.004	-0.056, 0.048	-0.15	0.883	0.989
lh_Lat_Fis.ant.Vertical	-0.022	-0.086, 0.041	-0.69	0.490	0.989
lh_Lat_Fis.post	-0.001	-0.044, 0.042	-0.05	0.960	0.989

eTable 22. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_Pole_occipital	0.036	-0.047, 0.119	0.86	0.393	0.989
lh_Pole_temporal	-0.013	-0.069, 0.042	-0.47	0.639	0.989
lh_S_calcarine	-0.001	-0.054, 0.051	-0.06	0.956	0.989
lh_S_central	-0.020	-0.06, 0.021	-0.96	0.339	0.989
lh_S_cingul.Marginalis	-0.020	-0.053, 0.014	-1.16	0.249	0.989
lh_S_circular_insula_ant	0.010	-0.034, 0.055	0.47	0.643	0.989
lh_S_circular_insula_inf	0.002	-0.031, 0.035	0.10	0.918	0.989
lh_S_circular_insula_sup	-0.003	-0.04, 0.034	-0.16	0.873	0.989
lh_S_collat_transv_ant	-0.011	-0.076, 0.055	-0.32	0.752	0.989
lh_S_collat_transv_post	-0.043	-0.089, 0.002	-1.88	0.063	0.989
lh_S_front_inf	-0.003	-0.049, 0.043	-0.14	0.893	0.989
lh_S_front_middle	-0.004	-0.05, 0.043	-0.15	0.882	0.989
lh_S_front_sup	-0.001	-0.035, 0.034	-0.04	0.966	0.989
lh_S_interm_prim.Jensen	0.046	0, 0.092	1.98	0.050	0.989
lh_S_intrapariet_and_P_trans	0.000	-0.039, 0.039	0.01	0.994	0.994
lh_S_oc_middle_and_Lunatus	-0.016	-0.06, 0.029	-0.69	0.493	0.989
lh_S_oc_sup_and_transversal	-0.004	-0.044, 0.035	-0.22	0.824	0.989
lh_S_occipital_ant	-0.012	-0.049, 0.025	-0.65	0.519	0.989
lh_S_oc.temp_lat	-0.004	-0.037, 0.028	-0.27	0.790	0.989
lh_S_oc.temp_med_and_Lingual	-0.021	-0.056, 0.013	-1.24	0.217	0.989
lh_S_orbital_lateral	-0.008	-0.087, 0.07	-0.21	0.836	0.989
lh_S_orbital_med.olfact	0.077	-0.044, 0.198	1.27	0.208	0.989
lh_S_orbital.H_Shaped	-0.002	-0.054, 0.05	-0.06	0.949	0.989
lh_S_parieto_occipital	-0.021	-0.059, 0.017	-1.10	0.273	0.989
lh_S_pericallosal	-0.009	-0.052, 0.035	-0.39	0.697	0.989
lh_S_postcentral	-0.009	-0.039, 0.021	-0.59	0.559	0.989
lh_S_precentral.inf.part	0.025	-0.015, 0.065	1.23	0.220	0.989
lh_S_precentral.sup.part	0.002	-0.032, 0.035	0.10	0.921	0.989
lh_S_suborbital	-0.015	-0.108, 0.077	-0.32	0.746	0.989
lh_S_subparietal	0.001	-0.043, 0.044	0.02	0.980	0.992
lh_S_temporal_inf	0.031	-0.029, 0.091	1.02	0.311	0.989
lh_S_temporal_sup	0.028	-0.006, 0.062	1.65	0.103	0.989
lh_S_temporal_transverse	-0.028	-0.086, 0.029	-0.98	0.331	0.989
rh_G_and_S_frontomargin	-0.022	-0.102, 0.059	-0.53	0.594	0.989
rh_G_and_S_occipital_inf	0.039	-0.015, 0.094	1.42	0.159	0.989
rh_G_and_S_paracentral	0.002	-0.044, 0.049	0.10	0.918	0.989
rh_G_and_S_subcentral	-0.016	-0.055, 0.024	-0.79	0.432	0.989
rh_G_and_S_transv_frontopol	-0.023	-0.079, 0.033	-0.83	0.411	0.989
rh_G_and_S_cingul.Ant	0.011	-0.028, 0.049	0.55	0.584	0.989
rh_G_and_S_cingul.Mid.Ant	-0.007	-0.04, 0.025	-0.45	0.654	0.989
rh_G_and_S_cingul.Mid.Post	0.014	-0.015, 0.044	0.98	0.329	0.989
rh_G_cingul.Post.dorsal	0.009	-0.03, 0.048	0.48	0.635	0.989
rh_G_cingul.Post.ventral	-0.027	-0.089, 0.036	-0.85	0.399	0.989


eTable 22. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_cuneus	-0.021	-0.071, 0.028	-0.86	0.391	0.989
rh_G_front_inf.Opercular	-0.009	-0.048, 0.029	-0.49	0.626	0.989
rh_G_front_inf.Orbital	-0.030	-0.085, 0.025	-1.09	0.276	0.989
rh_G_front_inf.Triangul	-0.022	-0.064, 0.02	-1.03	0.304	0.989
rh_G_front_middle	-0.003	-0.039, 0.034	-0.14	0.891	0.989
rh_G_front_sup	-0.009	-0.038, 0.019	-0.67	0.506	0.989
rh_G_Ins_lg_and_S_cent_ins	-0.001	-0.04, 0.038	-0.04	0.972	0.989
rh_G_insular_short	0.010	-0.023, 0.043	0.60	0.551	0.989
rh_G_occipital_middle	0.007	-0.035, 0.05	0.35	0.729	0.989
rh_G_occipital_sup	-0.009	-0.05, 0.032	-0.43	0.672	0.989
rh_G_oc.temp_lat.fusifor	0.006	-0.044, 0.056	0.24	0.810	0.989
rh_G_oc.temp_med.Lingual	-0.008	-0.067, 0.051	-0.27	0.788	0.989
rh_G_oc.temp_med.Parahip	0.006	-0.042, 0.054	0.24	0.808	0.989
rh_G_orbital	0.014	-0.053, 0.082	0.42	0.672	0.989
rh_G_pariet_inf.Angular	0.001	-0.034, 0.036	0.06	0.952	0.989
rh_G_pariet_inf.Supramar	-0.004	-0.035, 0.026	-0.29	0.773	0.989
rh_G_parietal_sup	0.017	-0.018, 0.051	0.97	0.335	0.989
rh_G_postcentral	0.006	-0.031, 0.043	0.30	0.764	0.989
rh_G_precentral	-0.011	-0.038, 0.016	-0.83	0.410	0.989
rh_G_precuneus	-0.009	-0.045, 0.026	-0.51	0.612	0.989
rh_G_rectus	0.092	-0.063, 0.247	1.18	0.241	0.989
rh_G_subcallosal	0.041	-0.024, 0.106	1.26	0.210	0.989
rh_G_temp_sup.G_T_transv	-0.007	-0.049, 0.035	-0.34	0.736	0.989
rh_G_temp_sup.Lateral	-0.002	-0.031, 0.028	-0.10	0.918	0.989
rh_G_temp_sup.Plan_polar	-0.003	-0.035, 0.028	-0.21	0.832	0.989
rh_G_temp_sup.Plan_tempo	-0.008	-0.048, 0.031	-0.42	0.674	0.989
rh_G_temporal_inf	0.001	-0.06, 0.062	0.04	0.965	0.989
rh_G_temporal_middle	0.006	-0.033, 0.046	0.32	0.748	0.989
rh_Lat_Fis.ant.Horizontal	-0.031	-0.082, 0.021	-1.18	0.240	0.989
rh_Lat_Fis.ant.Vertical	0.020	-0.049, 0.088	0.58	0.566	0.989
rh_Lat_Fis.post	0.019	-0.026, 0.065	0.85	0.396	0.989
rh_Pole_occipital	0.059	-0.018, 0.137	1.52	0.132	0.989
rh_Pole_temporal	-0.005	-0.07, 0.06	-0.16	0.877	0.989
rh_S_calcarine	-0.018	-0.071, 0.035	-0.68	0.499	0.989
rh_S_central	-0.002	-0.049, 0.044	-0.10	0.923	0.989
rh_S_cingul.Marginalis	-0.016	-0.047, 0.016	-0.97	0.335	0.989
rh_S_circular_insula_ant	0.011	-0.033, 0.056	0.51	0.611	0.989
rh_S_circular_insula_inf	0.000	-0.032, 0.031	-0.01	0.989	0.994
rh_S_circular_insula_sup	-0.019	-0.059, 0.02	-0.96	0.340	0.989
rh_S_collat_transv_ant	-0.051	-0.118, 0.016	-1.52	0.131	0.989
rh_S_collat_transv_post	-0.021	-0.065, 0.024	-0.92	0.360	0.989
rh_S_front_inf	0.003	-0.038, 0.045	0.15	0.877	0.989
rh_S_front_middle	-0.033	-0.071, 0.005	-1.73	0.086	0.989


eTable 22. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for loss feedback: Welch's t-test

ROIs	Mean	95%CI	t	<i>p</i> (unc)	<i>p</i> FDR
rh_S_front_sup	-0.002	-0.033, 0.029	-0.12	0.905	0.989
rh_S_interm_prim.Jensen	0.027	-0.018, 0.071	1.19	0.238	0.989
rh_S_intrapariet_and_P_trans	-0.004	-0.038, 0.031	-0.21	0.838	0.989
rh_S_oc_middle_and_Lunatus	-0.009	-0.052, 0.033	-0.44	0.663	0.989
rh_S_oc_sup_and_transversal	-0.014	-0.05, 0.023	-0.75	0.452	0.989
rh_S_occipital_ant	-0.011	-0.046, 0.025	-0.59	0.556	0.989
rh_S_oc.temp_lat	0.024	-0.014, 0.062	1.25	0.214	0.989
rh_S_oc.temp_med_and_Lingual	-0.003	-0.043, 0.036	-0.18	0.861	0.989
rh_S_orbital_lateral	-0.039	-0.103, 0.025	-1.22	0.226	0.989
rh_S_orbital_med.olfact	-0.065	-0.177, 0.047	-1.16	0.251	0.989
rh_S_orbital.H_Shaped	-0.011	-0.061, 0.038	-0.45	0.651	0.989
rh_S_parieto_occipital	-0.003	-0.042, 0.036	-0.14	0.892	0.989
rh_S_pericallosal	0.012	-0.031, 0.055	0.55	0.582	0.989
rh_S_postcentral	0.005	-0.028, 0.037	0.29	0.776	0.989
rh_S_precentral.inf.part	0.025	-0.016, 0.065	1.21	0.230	0.989
rh_S_precentral.sup.part	-0.002	-0.034, 0.031	-0.10	0.918	0.989
rh_S_suborbital	0.055	-0.085, 0.195	0.77	0.441	0.989
rh_S_subparietal	-0.025	-0.064, 0.015	-1.25	0.216	0.989
rh_S_temporal_inf	0.017	-0.044, 0.077	0.55	0.583	0.989
rh_S_temporal_sup	0.006	-0.025, 0.037	0.37	0.710	0.989
rh_S_temporal_transverse	0.012	-0.041, 0.066	0.45	0.650	0.989


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 30. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 31 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 31 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 31 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 32. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 32 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 32 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 33 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 34. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 35 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 35 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 35 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for loss feedback. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Stop Signal Task (SST)

Incorrect stop vs correct go

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.007	-0.029, 0.016	-0.59	0.554	0.988
Left.Thalamus.Proper	0.000	-0.011, 0.012	0.07	0.943	0.988
Left.Caudate	0.015	-0.003, 0.032	1.68	0.093	0.988
Left.Putamen	-0.002	-0.016, 0.012	-0.28	0.778	0.988
Left.Pallidum	0.006	-0.005, 0.018	1.11	0.266	0.988
Brain.Stem	-0.003	-0.016, 0.01	-0.39	0.698	0.988
Left.Hippocampus	0.004	-0.009, 0.017	0.60	0.551	0.988
Left.Amygdala	0.012	-0.003, 0.026	1.56	0.120	0.988
Left.Accumbens.area	0.006	-0.012, 0.023	0.66	0.512	0.988
Left.VentralDC	0.006	-0.008, 0.02	0.90	0.369	0.988
Right.Cerebellum.Cortex	0.003	-0.02, 0.025	0.25	0.802	0.988
Right.Thalamus.Proper	0.003	-0.01, 0.015	0.42	0.673	0.988
Right.Caudate	0.008	-0.01, 0.025	0.86	0.391	0.988
Right.Putamen	0.000	-0.015, 0.016	0.04	0.967	0.988
Right.Pallidum	-0.009	-0.021, 0.002	-1.63	0.105	0.988
Right.Hippocampus	-0.001	-0.014, 0.012	-0.15	0.881	0.988
Right.Amygdala	0.009	-0.007, 0.025	1.15	0.251	0.988
Right.Accumbens.area	-0.012	-0.03, 0.006	-1.34	0.182	0.988
Right.VentralDC	0.000	-0.014, 0.014	0.02	0.983	0.989
Cortical regions					
lh_G_and_S_frontomargin	-0.009	-0.037, 0.018	-0.67	0.505	0.988
lh_G_and_S_occipital_inf	-0.011	-0.034, 0.013	-0.89	0.374	0.988
lh_G_and_S_paracentral	-0.013	-0.031, 0.005	-1.43	0.154	0.988
lh_G_and_S_subcentral	0.000	-0.016, 0.016	0.04	0.971	0.988
lh_G_and_S_transv_frontopol	0.014	-0.013, 0.04	1.03	0.304	0.988
lh_G_and_S_cingul.Ant	0.007	-0.009, 0.023	0.82	0.415	0.988
lh_G_and_S_cingul.Mid.Ant	0.003	-0.012, 0.017	0.36	0.721	0.988
lh_G_and_S_cingul.Mid.Post	0.002	-0.012, 0.016	0.27	0.788	0.988
lh_G_cingul.Post.dorsal	0.017	0.001, 0.033	2.11	0.035	0.988
lh_G_cingul.Post.ventral	0.013	-0.018, 0.045	0.82	0.413	0.988
lh_G_cuneus	-0.022	-0.045, 0.002	-1.83	0.067	0.988
lh_G_front_inf.Opercular	0.006	-0.008, 0.02	0.83	0.409	0.988
lh_G_front_inf.Orbital	0.005	-0.019, 0.029	0.42	0.673	0.988

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_front_inf.Triangul	-0.002	-0.022, 0.018	-0.17	0.863	0.988
lh_G_front_middle	0.006	-0.009, 0.021	0.85	0.398	0.988
lh_G_front_sup	0.007	-0.004, 0.018	1.25	0.213	0.988
lh_G_Ins_lg_and_S_cent_ins	0.003	-0.012, 0.019	0.44	0.660	0.988
lh_G_insular_short	-0.002	-0.015, 0.01	-0.35	0.728	0.988
lh_G_occipital_middle	-0.003	-0.02, 0.013	-0.39	0.696	0.988
lh_G_occipital_sup	-0.023	-0.042, -0.003	-2.32	0.021	0.988
lh_G_oc.temp_lat.fusifor	-0.006	-0.025, 0.013	-0.61	0.542	0.988
lh_G_oc.temp_med.Lingual	-0.003	-0.029, 0.022	-0.26	0.798	0.988
lh_G_oc.temp_med.Parahip	0.004	-0.013, 0.021	0.46	0.646	0.988
lh_G_orbital	-0.010	-0.037, 0.016	-0.75	0.451	0.988
lh_G_pariet_inf.Angular	0.001	-0.015, 0.017	0.13	0.895	0.988
lh_G_pariet_inf.Supramar	-0.002	-0.016, 0.012	-0.23	0.821	0.988
lh_G_parietal_sup	-0.001	-0.017, 0.015	-0.09	0.924	0.988
lh_G_postcentral	-0.004	-0.021, 0.013	-0.49	0.626	0.988
lh_G_precentral	0.004	-0.008, 0.016	0.60	0.550	0.988
lh_G_precuneus	-0.003	-0.018, 0.012	-0.41	0.684	0.988
lh_G_rectus	-0.019	-0.067, 0.029	-0.78	0.434	0.988
lh_G_subcallosal	0.009	-0.017, 0.035	0.65	0.515	0.988
lh_G_temp_sup.G_T_transv	0.003	-0.013, 0.019	0.38	0.704	0.988
lh_G_temp_sup.Lateral	-0.004	-0.017, 0.008	-0.65	0.517	0.988
lh_G_temp_sup.Plan_polar	-0.002	-0.013, 0.009	-0.36	0.716	0.988
lh_G_temp_sup.Plan_tempo	0.010	-0.004, 0.024	1.38	0.170	0.988
lh_G_temporal_inf	-0.003	-0.027, 0.02	-0.29	0.775	0.988
lh_G_temporal_middle	-0.006	-0.022, 0.01	-0.72	0.473	0.988
lh_Lat_Fis.ant.Horizontal	0.005	-0.014, 0.024	0.50	0.619	0.988
lh_Lat_Fis.ant.Vertical	0.003	-0.021, 0.027	0.23	0.822	0.988
lh_Lat_Fis.post	-0.012	-0.028, 0.004	-1.45	0.149	0.988
lh_Pole_occipital	-0.006	-0.036, 0.024	-0.41	0.680	0.988
lh_Pole_temporal	0.001	-0.019, 0.02	0.06	0.956	0.988
lh_S_calcarine	-0.008	-0.032, 0.016	-0.66	0.507	0.988
lh_S_central	-0.001	-0.018, 0.017	-0.10	0.924	0.988
lh_S_cingul.Marginalis	0.010	-0.004, 0.025	1.38	0.170	0.988
lh_S_circular_insula_ant	0.004	-0.013, 0.022	0.48	0.631	0.988
lh_S_circular_insula_inf	0.002	-0.011, 0.015	0.29	0.770	0.988
lh_S_circular_insula_sup	0.007	-0.009, 0.022	0.85	0.396	0.988
lh_S_collat_transv_ant	-0.003	-0.025, 0.019	-0.27	0.786	0.988
lh_S_collat_transv_post	0.009	-0.011, 0.029	0.87	0.386	0.988
lh_S_front_inf	0.007	-0.011, 0.026	0.76	0.445	0.988
lh_S_front_middle	0.010	-0.008, 0.028	1.10	0.274	0.988
lh_S_front_sup	0.007	-0.005, 0.02	1.16	0.245	0.988
lh_S_interm_prim.Jensen	0.007	-0.013, 0.026	0.66	0.508	0.988
lh_S_intrapariet_and_P_trans	-0.008	-0.026, 0.009	-0.97	0.333	0.988

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_oc_middle_and_Lunatus	-0.017	-0.04, 0.005	-1.53	0.126	0.988
lh_S_oc_sup_and_transversal	-0.023	-0.042, -0.005	-2.54	0.011	0.988
lh_S_occipital_ant	-0.009	-0.027, 0.009	-0.98	0.328	0.988
lh_S_oc.temp_lat	-0.001	-0.017, 0.015	-0.13	0.895	0.988
lh_S_oc.temp_med_and_Lingual	0.007	-0.008, 0.022	0.95	0.345	0.988
lh_S_orbital_lateral	0.004	-0.021, 0.029	0.32	0.753	0.988
lh_S_orbital_med.olfact	-0.030	-0.067, 0.008	-1.55	0.123	0.988
lh_S_orbital.H_Shaped	0.003	-0.017, 0.024	0.31	0.760	0.988
lh_S_parieto_occipital	0.002	-0.014, 0.018	0.23	0.820	0.988
lh_S_pericallosal	0.004	-0.015, 0.023	0.42	0.676	0.988
lh_S_postcentral	0.000	-0.015, 0.014	-0.06	0.953	0.988
lh_S_precentral.inf.part	0.004	-0.013, 0.021	0.47	0.635	0.988
lh_S_precentral.sup.part	-0.004	-0.018, 0.01	-0.57	0.571	0.988
lh_S_suborbital	-0.007	-0.042, 0.028	-0.40	0.686	0.988
lh_S_subparietal	0.002	-0.017, 0.022	0.25	0.804	0.988
lh_S_temporal_inf	-0.010	-0.032, 0.012	-0.87	0.383	0.988
lh_S_temporal_sup	-0.003	-0.016, 0.011	-0.39	0.698	0.988
lh_S_temporal_transverse	0.005	-0.015, 0.025	0.46	0.646	0.988
rh_G_and_S_frontomargin	-0.003	-0.032, 0.026	-0.23	0.819	0.988
rh_G_and_S_occipital_inf	-0.011	-0.033, 0.011	-0.95	0.342	0.988
rh_G_and_S_paracentral	-0.012	-0.031, 0.007	-1.24	0.215	0.988
rh_G_and_S_subcentral	0.005	-0.012, 0.022	0.56	0.576	0.988
rh_G_and_S_transv_frontopol	0.003	-0.023, 0.028	0.21	0.835	0.988
rh_G_and_S_cingul.Ant	0.009	-0.008, 0.026	1.07	0.285	0.988
rh_G_and_S_cingul.Mid.Ant	0.002	-0.012, 0.017	0.32	0.753	0.988
rh_G_and_S_cingul.Mid.Post	0.000	-0.013, 0.012	-0.03	0.976	0.988
rh_G_cingul.Post.dorsal	0.006	-0.01, 0.022	0.70	0.482	0.988
rh_G_cingul.Post.ventral	-0.001	-0.026, 0.024	-0.08	0.938	0.988
rh_G_cuneus	-0.002	-0.025, 0.021	-0.19	0.849	0.988
rh_G_front_inf.Opercular	0.005	-0.011, 0.02	0.57	0.571	0.988
rh_G_front_inf.Orbital	-0.010	-0.032, 0.012	-0.88	0.382	0.988
rh_G_front_inf.Triangul	0.000	-0.019, 0.018	-0.03	0.977	0.988
rh_G_front_middle	-0.001	-0.018, 0.015	-0.16	0.872	0.988
rh_G_front_sup	0.000	-0.011, 0.011	-0.04	0.969	0.988
rh_G_Ins_lg_and_S_cent_ins	0.003	-0.012, 0.019	0.44	0.663	0.988
rh_G_insular_short	-0.001	-0.016, 0.014	-0.15	0.879	0.988
rh_G_occipital_middle	-0.002	-0.019, 0.014	-0.28	0.780	0.988
rh_G_occipital_sup	-0.008	-0.026, 0.011	-0.80	0.426	0.988
rh_G_oc.temp_lat.fusifor	0.009	-0.01, 0.028	0.92	0.356	0.988
rh_G_oc.temp_med.Lingual	-0.005	-0.031, 0.021	-0.39	0.695	0.988
rh_G_oc.temp_med.Parahip	-0.005	-0.024, 0.014	-0.49	0.622	0.988
rh_G_orbital	0.002	-0.023, 0.028	0.18	0.856	0.988
rh_G_pariet_inf.Angular	0.002	-0.014, 0.017	0.21	0.836	0.988

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_pariet_inf.Supramar	-0.001	-0.015, 0.013	-0.14	0.889	0.988
rh_G_parietal_sup	-0.003	-0.018, 0.013	-0.35	0.723	0.988
rh_G_postcentral	-0.004	-0.02, 0.013	-0.42	0.672	0.988
rh_G_precentral	0.005	-0.007, 0.017	0.87	0.384	0.988
rh_G_precuneus	0.006	-0.011, 0.022	0.68	0.496	0.988
rh_G_rectus	-0.010	-0.067, 0.047	-0.34	0.737	0.988
rh_G_subcallosal	0.008	-0.019, 0.034	0.56	0.575	0.988
rh_G_temp_sup.G_T_transv	0.010	-0.007, 0.026	1.16	0.248	0.988
rh_G_temp_sup.Lateral	0.004	-0.009, 0.016	0.55	0.580	0.988
rh_G_temp_sup.Plan_polar	-0.007	-0.019, 0.005	-1.10	0.270	0.988
rh_G_temp_sup.Plan_tempo	0.004	-0.011, 0.02	0.55	0.584	0.988
rh_G_temporal_inf	0.001	-0.022, 0.024	0.11	0.914	0.988
rh_G_temporal_middle	-0.004	-0.021, 0.012	-0.51	0.612	0.988
rh_Lat_Fis.ant.Horizontal	0.007	-0.013, 0.027	0.67	0.506	0.988
rh_Lat_Fis.ant.Vertical	-0.011	-0.035, 0.013	-0.90	0.368	0.988
rh_Lat_Fis.post	0.001	-0.018, 0.02	0.09	0.931	0.988
rh_Pole_occipital	0.004	-0.025, 0.032	0.24	0.811	0.988
rh_Pole_temporal	0.012	-0.01, 0.034	1.08	0.279	0.988
rh_S_calcarine	-0.006	-0.029, 0.017	-0.49	0.623	0.988
rh_S_central	0.001	-0.016, 0.018	0.09	0.928	0.988
rh_S_cingul.Marginalis	0.004	-0.01, 0.018	0.60	0.548	0.988
rh_S_circular_insula_ant	-0.002	-0.019, 0.016	-0.21	0.837	0.988
rh_S_circular_insula_inf	0.002	-0.012, 0.017	0.28	0.781	0.988
rh_S_circular_insula_sup	-0.002	-0.018, 0.014	-0.27	0.787	0.988
rh_S_collat_transv_ant	-0.016	-0.04, 0.008	-1.29	0.200	0.988
rh_S_collat_transv_post	-0.011	-0.031, 0.009	-1.09	0.274	0.988
rh_S_front_inf	-0.004	-0.022, 0.014	-0.48	0.632	0.988
rh_S_front_middle	0.008	-0.011, 0.026	0.83	0.407	0.988
rh_S_front_sup	0.000	-0.013, 0.013	0.03	0.975	0.988
rh_S_interm_prim.Jensen	-0.002	-0.022, 0.017	-0.25	0.806	0.988
rh_S_intrapariet_and_P_trans	-0.004	-0.021, 0.013	-0.43	0.666	0.988
rh_S_oc_middle_and_Lunatus	-0.019	-0.04, 0.002	-1.76	0.079	0.988
rh_S_oc_sup_and_transversal	-0.015	-0.033, 0.003	-1.65	0.099	0.988
rh_S_occipital_ant	-0.005	-0.022, 0.012	-0.59	0.558	0.988
rh_S_oc.temp_lat	-0.015	-0.031, 0	-1.94	0.053	0.988
rh_S_oc.temp_med_and_Lingual	-0.006	-0.02, 0.007	-0.92	0.357	0.988
rh_S_orbital_lateral	-0.015	-0.04, 0.009	-1.22	0.223	0.988
rh_S_orbital_med.olfact	0.002	-0.039, 0.044	0.12	0.905	0.988
rh_S_orbital.H_Shaped	-0.009	-0.03, 0.012	-0.81	0.417	0.988
rh_S_parieto_occipital	-0.007	-0.024, 0.01	-0.84	0.401	0.988
rh_S_pericallosal	0.006	-0.015, 0.028	0.56	0.579	0.988
rh_S_postcentral	0.002	-0.011, 0.016	0.33	0.739	0.988
rh_S_precentral.inf.part	-0.006	-0.024, 0.012	-0.62	0.533	0.988

eTable 23. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_precentral.sup.part	0.003	-0.011, 0.016	0.36	0.721	0.988
rh_S_suborbital	-0.018	-0.066, 0.03	-0.73	0.464	0.988
rh_S_subparietal	0.006	-0.011, 0.023	0.66	0.511	0.988
rh_S_temporal_inf	-0.029	-0.053, -0.005	-2.38	0.018	0.988
rh_S_temporal_sup	0.000	-0.013, 0.013	-0.01	0.992	0.992
rh_S_temporal_transverse	-0.007	-0.029, 0.014	-0.68	0.498	0.988

Bold denotes group differences at p<0.05 uncorrected

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.011	-0.011, 0.032	0.96	0.337	0.910
Left.Thalamus.Proper	-0.004	-0.015, 0.008	-0.62	0.532	0.910
Left.Caudate	-0.010	-0.027, 0.007	-1.18	0.238	0.910
Left.Putamen	-0.004	-0.017, 0.009	-0.57	0.566	0.910
Left.Pallidum	-0.003	-0.015, 0.009	-0.53	0.597	0.910
Brain.Stem	0.004	-0.009, 0.017	0.55	0.584	0.910
Left.Hippocampus	0.005	-0.009, 0.018	0.72	0.474	0.910
Left.Amygdala	0.004	-0.011, 0.018	0.49	0.624	0.921
Left.Accumbens.area	-0.005	-0.023, 0.014	-0.51	0.609	0.910
Left.VentralDC	-0.005	-0.017, 0.008	-0.70	0.483	0.910
Right.Cerebellum.Cortex	0.014	-0.009, 0.037	1.17	0.245	0.910
Right.Thalamus.Proper	-0.003	-0.015, 0.009	-0.53	0.597	0.910
Right.Caudate	-0.013	-0.031, 0.005	-1.40	0.161	0.910
Right.Putamen	-0.003	-0.017, 0.011	-0.37	0.711	0.935
Right.Pallidum	-0.010	-0.022, 0.001	-1.81	0.072	0.910
Right.Hippocampus	-0.002	-0.015, 0.011	-0.25	0.803	0.974
Right.Amygdala	0.013	-0.003, 0.03	1.57	0.116	0.910
Right.Accumbens.area	-0.008	-0.025, 0.009	-0.90	0.370	0.910
Right.VentralDC	0.004	-0.009, 0.017	0.60	0.551	0.910
Cortical regions					
lh_G_and_S_frontomargin	0.013	-0.016, 0.043	0.90	0.371	0.910
lh_G_and_S_occipital_inf	-0.009	-0.032, 0.014	-0.80	0.423	0.910
lh_G_and_S_paracentral	-0.011	-0.028, 0.006	-1.23	0.220	0.910
lh_G_and_S_subcentral	-0.004	-0.019, 0.012	-0.45	0.656	0.924
lh_G_and_S_transv_frontopol	0.013	-0.016, 0.042	0.87	0.386	0.910
lh_G_and_S_cingul.Ant	0.000	-0.017, 0.016	-0.05	0.958	0.993
lh_G_and_S_cingul.Mid.Ant	-0.011	-0.025, 0.004	-1.43	0.153	0.910
lh_G_and_S_cingul.Mid.Post	-0.001	-0.014, 0.012	-0.15	0.883	0.974

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_cingul.Post.dorsal	0.010	-0.007, 0.027	1.13	0.258	0.910
lh_G_cingul.Post.ventral	-0.010	-0.044, 0.024	-0.60	0.550	0.910
lh_G_cuneus	-0.013	-0.036, 0.009	-1.15	0.249	0.910
lh_G_front_inf.Opercular	-0.008	-0.022, 0.006	-1.14	0.257	0.910
lh_G_front_inf.Orbital	-0.009	-0.031, 0.014	-0.77	0.445	0.910
lh_G_front_inf.Triangul	-0.004	-0.023, 0.015	-0.39	0.700	0.935
lh_G_front_middle	0.000	-0.015, 0.016	0.05	0.964	0.993
lh_G_front_sup	-0.005	-0.016, 0.007	-0.81	0.421	0.910
lh_G_Ins_lg_and_S_cent_ins	0.007	-0.007, 0.022	0.99	0.323	0.910
lh_G_insular_short	-0.006	-0.019, 0.007	-0.94	0.347	0.910
lh_G_occipital_middle	-0.005	-0.021, 0.012	-0.53	0.594	0.910
lh_G_occipital_sup	-0.012	-0.031, 0.007	-1.24	0.216	0.910
lh_G_oc.temp_lat.fusifor	-0.005	-0.024, 0.014	-0.53	0.596	0.910
lh_G_oc.temp_med.Lingual	0.013	-0.012, 0.038	0.99	0.320	0.910
lh_G_oc.temp_med.Parahip	0.001	-0.016, 0.018	0.12	0.903	0.974
lh_G_orbital	-0.006	-0.035, 0.023	-0.41	0.680	0.931
lh_G_pariet_inf.Angular	0.006	-0.01, 0.023	0.73	0.468	0.910
lh_G_pariet_inf.Supramar	0.004	-0.01, 0.017	0.55	0.585	0.910
lh_G_parietal_sup	-0.002	-0.017, 0.013	-0.29	0.773	0.972
lh_G_postcentral	-0.005	-0.021, 0.011	-0.58	0.561	0.910
lh_G_precentral	-0.005	-0.017, 0.006	-0.91	0.364	0.910
lh_G_precuneus	0.010	-0.005, 0.025	1.30	0.194	0.910
lh_G_rectus	0.018	-0.032, 0.067	0.70	0.485	0.910
lh_G_subcallosal	0.020	-0.006, 0.045	1.53	0.128	0.910
lh_G_temp_sup.G_T_transv	0.004	-0.01, 0.019	0.59	0.552	0.910
lh_G_temp_sup.Lateral	-0.005	-0.017, 0.008	-0.73	0.467	0.910
lh_G_temp_sup.Plan_polar	0.002	-0.01, 0.013	0.28	0.777	0.972
lh_G_temp_sup.Plan_tempo	0.010	-0.005, 0.025	1.32	0.188	0.910
lh_G_temporal_inf	0.006	-0.018, 0.029	0.48	0.629	0.921
lh_G_temporal_middle	-0.004	-0.02, 0.013	-0.44	0.662	0.924
lh_Lat_Fis.ant.Horizontal	-0.009	-0.03, 0.011	-0.91	0.364	0.910
lh_Lat_Fis.ant.Vertical	-0.001	-0.026, 0.023	-0.12	0.905	0.974
lh_Lat_Fis.post	-0.004	-0.02, 0.012	-0.51	0.610	0.910
lh_Pole_occipital	0.012	-0.016, 0.039	0.85	0.398	0.910
lh_Pole_temporal	0.017	-0.005, 0.038	1.53	0.128	0.910
lh_S_calcarine	-0.011	-0.034, 0.012	-0.96	0.335	0.910
lh_S_central	0.004	-0.014, 0.022	0.44	0.663	0.924
lh_S_cingul.Marginalis	0.009	-0.005, 0.024	1.24	0.214	0.910
lh_S_circular_insula_ant	-0.002	-0.019, 0.016	-0.18	0.859	0.974
lh_S_circular_insula_inf	-0.006	-0.018, 0.007	-0.90	0.370	0.910
lh_S_circular_insula_sup	-0.001	-0.016, 0.014	-0.12	0.901	0.974
lh_S_collat_transv_ant	-0.013	-0.036, 0.011	-1.06	0.290	0.910
lh_S_collat_transv_post	-0.002	-0.022, 0.017	-0.26	0.798	0.974

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_front_inf	0.002	-0.015, 0.02	0.25	0.806	0.974
lh_S_front_middle	0.006	-0.012, 0.024	0.63	0.531	0.910
lh_S_front_sup	0.001	-0.012, 0.015	0.23	0.821	0.974
lh_S_interm_prim.Jensen	0.006	-0.014, 0.027	0.60	0.547	0.910
lh_S_intrapariet_and_P_trans	0.008	-0.008, 0.025	0.99	0.321	0.910
lh_S_oc_middle_and_Lunatus	-0.025	-0.046, -0.005	-2.46	0.014	0.910
lh_S_oc_sup_and_transversal	-0.006	-0.024, 0.012	-0.64	0.525	0.910
lh_S_occipital_ant	-0.008	-0.026, 0.01	-0.87	0.383	0.910
lh_S_oc.temp_lat	0.002	-0.013, 0.017	0.28	0.780	0.972
lh_S_oc.temp_med_and_Lingual	0.004	-0.01, 0.018	0.62	0.533	0.910
lh_S_orbital_lateral	-0.011	-0.038, 0.015	-0.84	0.403	0.910
lh_S_orbital_med.olfact	-0.032	-0.074, 0.01	-1.52	0.131	0.910
lh_S_orbital.H_Shaped	0.005	-0.016, 0.026	0.47	0.637	0.924
lh_S_parieto_occipital	0.014	-0.002, 0.03	1.70	0.090	0.910
lh_S_pericallosal	0.000	-0.019, 0.018	-0.04	0.969	0.993
lh_S_postcentral	0.001	-0.012, 0.014	0.16	0.871	0.974
lh_S_precentral.inf.part	0.009	-0.008, 0.026	1.08	0.283	0.910
lh_S_precentral.sup.part	-0.004	-0.018, 0.009	-0.60	0.550	0.910
lh_S_suborbital	0.013	-0.023, 0.05	0.72	0.472	0.910
lh_S_subparietal	0.007	-0.013, 0.027	0.71	0.478	0.910
lh_S_temporal_inf	-0.014	-0.037, 0.009	-1.17	0.242	0.910
lh_S_temporal_sup	-0.003	-0.016, 0.011	-0.38	0.706	0.935
lh_S_temporal_transverse	0.008	-0.013, 0.029	0.75	0.452	0.910
rh_G_and_S_frontomargin	-0.002	-0.035, 0.03	-0.14	0.886	0.974
rh_G_and_S_occipital_inf	-0.002	-0.025, 0.02	-0.21	0.831	0.974
rh_G_and_S_paracentral	0.000	-0.018, 0.018	-0.01	0.993	0.993
rh_G_and_S_subcentral	0.006	-0.011, 0.022	0.67	0.502	0.910
rh_G_and_S_transv_frontopol	-0.016	-0.04, 0.008	-1.29	0.199	0.910
rh_G_and_S_cingul.Ant	-0.001	-0.018, 0.016	-0.15	0.882	0.974
rh_G_and_S_cingul.Mid.Ant	0.004	-0.01, 0.018	0.53	0.596	0.910
rh_G_and_S_cingul.Mid.Post	0.007	-0.005, 0.019	1.21	0.227	0.910
rh_G_cingul.Post.dorsal	0.001	-0.015, 0.017	0.11	0.915	0.974
rh_G_cingul.Post.ventral	-0.002	-0.028, 0.025	-0.11	0.910	0.974
rh_G_cuneus	-0.002	-0.025, 0.021	-0.19	0.849	0.974
rh_G_front_inf.Opercular	-0.010	-0.025, 0.005	-1.32	0.187	0.910
rh_G_front_inf.Orbital	-0.014	-0.037, 0.009	-1.23	0.220	0.910
rh_G_front_inf.Triangul	-0.006	-0.024, 0.012	-0.70	0.487	0.910
rh_G_front_middle	-0.013	-0.029, 0.003	-1.59	0.114	0.910
rh_G_front_sup	-0.004	-0.016, 0.008	-0.66	0.509	0.910
rh_G_Ins_lg_and_S_cent_ins	0.003	-0.012, 0.018	0.36	0.723	0.940
rh_G_insular_short	0.000	-0.014, 0.013	-0.06	0.954	0.993
rh_G_occipital_middle	0.000	-0.016, 0.017	0.01	0.991	0.993
rh_G_occipital_sup	0.005	-0.013, 0.023	0.55	0.582	0.910

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_oc.temp_lat.fusifor	0.006	-0.013, 0.026	0.64	0.521	0.910
rh_G_oc.temp_med.Lingual	0.000	-0.026, 0.025	-0.02	0.982	0.993
rh_G_oc.temp_med.Parahip	-0.005	-0.025, 0.014	-0.55	0.580	0.910
rh_G_orbital	0.009	-0.018, 0.036	0.67	0.506	0.910
rh_G_pariet_inf.Angular	-0.003	-0.019, 0.014	-0.34	0.736	0.940
rh_G_pariet_inf.Supramar	0.007	-0.006, 0.02	1.02	0.307	0.910
rh_G_parietal_sup	-0.003	-0.018, 0.011	-0.44	0.664	0.924
rh_G_postcentral	0.009	-0.007, 0.024	1.05	0.293	0.910
rh_G_precentral	0.003	-0.008, 0.014	0.52	0.602	0.910
rh_G_precuneus	0.003	-0.013, 0.019	0.34	0.737	0.940
rh_G_rectus	0.000	-0.061, 0.062	0.01	0.993	0.993
rh_G_subcallosal	-0.006	-0.033, 0.021	-0.45	0.652	0.924
rh_G_temp_sup.G_T_transv	0.006	-0.01, 0.021	0.73	0.465	0.910
rh_G_temp_sup.Lateral	-0.001	-0.014, 0.012	-0.12	0.904	0.974
rh_G_temp_sup.Plan_polar	0.001	-0.011, 0.013	0.16	0.870	0.974
rh_G_temp_sup.Plan_tempo	0.013	-0.002, 0.029	1.70	0.090	0.910
rh_G_temporal_inf	0.001	-0.022, 0.025	0.11	0.909	0.974
rh_G_temporal_middle	-0.009	-0.025, 0.008	-1.00	0.316	0.910
rh_Lat_Fis.ant.Horizontal	0.003	-0.016, 0.023	0.34	0.734	0.940
rh_Lat_Fis.ant.Vertical	-0.027	-0.052, -0.002	-2.10	0.037	0.910
rh_Lat_Fis.post	0.005	-0.013, 0.023	0.53	0.598	0.910
rh_Pole_occipital	0.003	-0.024, 0.03	0.22	0.829	0.974
rh_Pole_temporal	-0.003	-0.025, 0.02	-0.22	0.825	0.974
rh_S_calcarine	-0.003	-0.026, 0.02	-0.25	0.803	0.974
rh_S_central	0.007	-0.01, 0.024	0.77	0.442	0.910
rh_S_cingul.Marginalis	0.004	-0.01, 0.019	0.57	0.572	0.910
rh_S_circular_insula_ant	-0.012	-0.029, 0.006	-1.29	0.197	0.910
rh_S_circular_insula_inf	0.008	-0.006, 0.023	1.18	0.239	0.910
rh_S_circular_insula_sup	-0.006	-0.021, 0.01	-0.73	0.468	0.910
rh_S_collat_transv_ant	-0.027	-0.052, -0.003	-2.23	0.027	0.910
rh_S_collat_transv_post	0.001	-0.019, 0.02	0.06	0.955	0.993
rh_S_front_inf	-0.008	-0.027, 0.01	-0.86	0.392	0.910
rh_S_front_middle	-0.008	-0.028, 0.012	-0.79	0.429	0.910
rh_S_front_sup	-0.003	-0.017, 0.011	-0.38	0.705	0.935
rh_S_interm_prim.Jensen	0.001	-0.019, 0.021	0.09	0.927	0.980
rh_S_intrapariet_and_P_trans	0.008	-0.008, 0.024	1.02	0.310	0.910
rh_S_oc_middle_and_Lunatus	-0.014	-0.034, 0.007	-1.32	0.187	0.910
rh_S_oc_sup_and_transversal	0.008	-0.01, 0.025	0.89	0.377	0.910
rh_S_occipital_ant	0.014	-0.004, 0.032	1.49	0.136	0.910
rh_S_oc.temp_lat	-0.005	-0.021, 0.011	-0.60	0.549	0.910
rh_S_oc.temp_med_and_Lingual	0.006	-0.008, 0.02	0.85	0.398	0.910
rh_S_orbital_lateral	-0.018	-0.042, 0.007	-1.40	0.163	0.910
rh_S_orbital_med.olfact	-0.011	-0.054, 0.032	-0.52	0.606	0.910

eTable 24. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_orbital.H_Shaped	-0.017	-0.039, 0.005	-1.50	0.134	0.910
rh_S_parieto_occipital	0.008	-0.009, 0.024	0.94	0.350	0.910
rh_S_pericallosal	0.004	-0.016, 0.024	0.40	0.686	0.931
rh_S_postcentral	0.006	-0.008, 0.02	0.82	0.413	0.910
rh_S_precentral.inf.part	-0.011	-0.029, 0.007	-1.24	0.215	0.910
rh_S_precentral.sup.part	0.001	-0.013, 0.014	0.12	0.903	0.974
rh_S_suborbital	-0.017	-0.071, 0.037	-0.63	0.529	0.910
rh_S_subparietal	0.015	-0.002, 0.033	1.75	0.081	0.910
rh_S_temporal_inf	-0.030	-0.053, -0.007	-2.51	0.012	0.910
rh_S_temporal_sup	0.006	-0.007, 0.02	0.91	0.363	0.910
rh_S_temporal_transverse	0.004	-0.017, 0.026	0.41	0.685	0.931

Bold denotes group differences at p<0.05 uncorrected

eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.011	-0.028, 0.049	0.56	0.579	0.973
Left.Thalamus.Proper	-0.004	-0.022, 0.014	-0.44	0.660	0.973
Left.Caudate	0.002	-0.028, 0.032	0.12	0.904	0.973
Left.Putamen	-0.010	-0.035, 0.014	-0.83	0.408	0.973
Left.Pallidum	0.014	-0.005, 0.034	1.44	0.153	0.973
Brain.Stem	-0.003	-0.025, 0.019	-0.24	0.811	0.973
Left.Hippocampus	0.013	-0.009, 0.035	1.20	0.233	0.973
Left.Amygdala	0.006	-0.02, 0.031	0.43	0.670	0.973
Left.Accumbens.area	0.005	-0.026, 0.036	0.31	0.757	0.973
Left.VentralDC	0.004	-0.019, 0.027	0.35	0.724	0.973
Right.Cerebellum.Cortex	0.029	-0.013, 0.07	1.37	0.175	0.973
Right.Thalamus.Proper	-0.004	-0.026, 0.017	-0.37	0.711	0.973
Right.Caudate	-0.011	-0.041, 0.019	-0.71	0.480	0.973
Right.Putamen	-0.008	-0.034, 0.018	-0.60	0.552	0.973
Right.Pallidum	-0.018	-0.038, 0.003	-1.71	0.091	0.973
Right.Hippocampus	0.008	-0.014, 0.031	0.72	0.475	0.973
Right.Amygdala	0.019	-0.01, 0.048	1.32	0.190	0.973
Right.Accumbens.area	-0.012	-0.045, 0.021	-0.70	0.487	0.973
Right.VentralDC	-0.003	-0.028, 0.022	-0.21	0.835	0.973
Cortical regions					
lh_G_and_S_frontomargin	0.010	-0.042, 0.062	0.40	0.692	0.973
lh_G_and_S_occipital_inf	0.004	-0.036, 0.045	0.22	0.827	0.973
lh_G_and_S_paracentral	-0.011	-0.041, 0.02	-0.69	0.490	0.973

eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_and_S_subcentral	-0.017	-0.047, 0.013	-1.13	0.262	0.973
lh_G_and_S_transv_frontopol	0.013	-0.04, 0.066	0.48	0.631	0.973
lh_G_and_S_cingul.Ant	-0.001	-0.029, 0.026	-0.09	0.927	0.973
lh_G_and_S_cingul.Mid.Ant	-0.020	-0.047, 0.006	-1.52	0.132	0.973
lh_G_and_S_cingul.Mid.Post	-0.005	-0.031, 0.021	-0.39	0.695	0.973
lh_G_cingul.Post.dorsal	0.022	-0.004, 0.048	1.69	0.094	0.973
lh_G_cingul.Post.ventral	0.022	-0.035, 0.079	0.77	0.442	0.973
lh_G_cuneus	-0.029	-0.07, 0.012	-1.41	0.163	0.973
lh_G_front_inf.Opercular	-0.003	-0.027, 0.022	-0.21	0.833	0.973
lh_G_front_inf.Orbital	0.002	-0.042, 0.046	0.09	0.931	0.973
lh_G_front_inf.Triangul	0.012	-0.024, 0.049	0.66	0.509	0.973
lh_G_front_middle	0.005	-0.02, 0.03	0.43	0.671	0.973
lh_G_front_sup	0.001	-0.02, 0.021	0.07	0.944	0.973
lh_G_Ins_lg_and_S_cent_ins	0.006	-0.021, 0.034	0.45	0.656	0.973
lh_G_insular_short	-0.004	-0.027, 0.019	-0.32	0.746	0.973
lh_G_occipital_middle	0.000	-0.026, 0.026	-0.01	0.993	0.993
lh_G_occipital_sup	-0.014	-0.049, 0.021	-0.80	0.426	0.973
lh_G_oc.temp_lat.fusifor	0.002	-0.033, 0.037	0.13	0.899	0.973
lh_G_oc.temp_med.Lingual	0.015	-0.029, 0.058	0.67	0.502	0.973
lh_G_oc.temp_med.Parahip	-0.007	-0.035, 0.02	-0.53	0.599	0.973
lh_G_orbital	-0.009	-0.063, 0.046	-0.31	0.754	0.973
lh_G_pariet_inf.Angular	0.005	-0.023, 0.033	0.34	0.734	0.973
lh_G_pariet_inf.Supramar	0.009	-0.014, 0.032	0.80	0.423	0.973
lh_G_parietal_sup	0.004	-0.022, 0.029	0.28	0.778	0.973
lh_G_postcentral	0.003	-0.025, 0.032	0.24	0.812	0.973
lh_G_precentral	-0.005	-0.026, 0.015	-0.53	0.595	0.973
lh_G_precuneus	0.016	-0.009, 0.042	1.28	0.204	0.973
lh_G_rectus	0.005	-0.087, 0.096	0.10	0.922	0.973
lh_G_subcallosal	0.011	-0.037, 0.06	0.47	0.638	0.973
lh_G_temp_sup.G_T_transv	-0.002	-0.028, 0.024	-0.14	0.890	0.973
lh_G_temp_sup.Lateral	0.002	-0.018, 0.023	0.23	0.815	0.973
lh_G_temp_sup.Plan_polar	-0.001	-0.022, 0.02	-0.11	0.912	0.973
lh_G_temp_sup.Plan_tempo	0.023	-0.001, 0.047	1.90	0.061	0.973
lh_G_temporal_inf	0.008	-0.036, 0.051	0.35	0.730	0.973
lh_G_temporal_middle	0.002	-0.026, 0.03	0.16	0.872	0.973
lh_Lat_Fis.ant.Horizontal	0.000	-0.034, 0.033	-0.02	0.983	0.989
lh_Lat_Fis.ant.Vertical	0.024	-0.022, 0.069	1.04	0.303	0.973
lh_Lat_Fis.post	-0.019	-0.047, 0.008	-1.39	0.166	0.973
lh_Pole_occipital	0.014	-0.036, 0.063	0.55	0.586	0.973
lh_Pole_temporal	0.009	-0.026, 0.044	0.50	0.618	0.973
lh_S_calcarine	-0.029	-0.07, 0.012	-1.39	0.169	0.973
lh_S_central	-0.011	-0.042, 0.02	-0.70	0.484	0.973
lh_S_cingul.Marginalis	0.012	-0.012, 0.037	1.02	0.309	0.973

eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_circular_insula_ant	-0.009	-0.039, 0.021	-0.60	0.552	0.973
lh_S_circular_insula_inf	-0.010	-0.032, 0.012	-0.91	0.367	0.973
lh_S_circular_insula_sup	0.001	-0.029, 0.031	0.08	0.936	0.973
lh_S_collat_transv_ant	0.021	-0.021, 0.062	1.00	0.321	0.973
lh_S_collat_transv_post	0.005	-0.029, 0.039	0.30	0.762	0.973
lh_S_front_inf	0.018	-0.015, 0.05	1.08	0.283	0.973
lh_S_front_middle	0.017	-0.014, 0.049	1.09	0.279	0.973
lh_S_front_sup	-0.002	-0.025, 0.021	-0.17	0.865	0.973
lh_S_interm_prim.Jensen	0.016	-0.019, 0.05	0.90	0.370	0.973
lh_S_intrapariet_and_P_trans	0.009	-0.02, 0.039	0.63	0.530	0.973
lh_S_oc_middle_and_Lunatus	-0.031	-0.063, 0	-1.96	0.053	0.973
lh_S_oc_sup_and_transversal	-0.016	-0.045, 0.014	-1.03	0.305	0.973
lh_S_occipital_ant	0.001	-0.027, 0.029	0.08	0.933	0.973
lh_S_oc.temp_lat	0.005	-0.023, 0.033	0.38	0.707	0.973
lh_S_oc.temp_med_and_Lingual	-0.003	-0.027, 0.021	-0.21	0.830	0.973
lh_S_orbital_lateral	0.010	-0.032, 0.051	0.45	0.652	0.973
lh_S_orbital_med.olfact	-0.003	-0.074, 0.067	-0.10	0.923	0.973
lh_S_orbital.H_Shaped	0.034	-0.002, 0.069	1.90	0.061	0.973
lh_S_parieto_occipital	0.011	-0.018, 0.04	0.77	0.441	0.973
lh_S_pericallosal	-0.007	-0.04, 0.026	-0.43	0.671	0.973
lh_S_postcentral	-0.003	-0.027, 0.021	-0.23	0.822	0.973
lh_S_precentral.inf.part	0.005	-0.025, 0.036	0.35	0.727	0.973
lh_S_precentral.sup.part	0.005	-0.018, 0.028	0.40	0.687	0.973
lh_S_suborbital	-0.001	-0.07, 0.068	-0.03	0.974	0.985
lh_S_subparietal	-0.005	-0.039, 0.028	-0.31	0.758	0.973
lh_S_temporal_inf	-0.012	-0.051, 0.026	-0.62	0.534	0.973
lh_S_temporal_sup	-0.003	-0.026, 0.021	-0.24	0.808	0.973
lh_S_temporal_transverse	0.006	-0.029, 0.042	0.36	0.721	0.973
rh_G_and_S_frontomargin	0.003	-0.046, 0.052	0.12	0.907	0.973
rh_G_and_S_occipital_inf	-0.003	-0.043, 0.038	-0.14	0.892	0.973
rh_G_and_S_paracentral	-0.006	-0.038, 0.026	-0.35	0.727	0.973
rh_G_and_S_subcentral	0.007	-0.026, 0.039	0.41	0.681	0.973
rh_G_and_S_transv_frontopol	-0.036	-0.082, 0.01	-1.56	0.123	0.973
rh_G_and_S_cingul.Ant	0.001	-0.027, 0.03	0.09	0.929	0.973
rh_G_and_S_cingul.Mid.Ant	-0.013	-0.038, 0.012	-1.03	0.304	0.973
rh_G_and_S_cingul.Mid.Post	0.002	-0.016, 0.02	0.23	0.816	0.973
rh_G_cingul.Post.dorsal	-0.003	-0.029, 0.023	-0.24	0.812	0.973
rh_G_cingul.Post.ventral	-0.003	-0.048, 0.042	-0.14	0.887	0.973
rh_G_cuneus	-0.018	-0.057, 0.021	-0.89	0.373	0.973
rh_G_front_inf.Opercular	-0.015	-0.041, 0.012	-1.10	0.275	0.973
rh_G_front_inf.Orbital	-0.035	-0.073, 0.004	-1.79	0.077	0.973
rh_G_front_inf.Triangul	0.010	-0.025, 0.045	0.57	0.572	0.973
rh_G_front_middle	-0.011	-0.041, 0.018	-0.76	0.450	0.973


eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_front_sup	-0.006	-0.025, 0.014	-0.57	0.570	0.973
rh_G_Ins_lg_and_S_cent_ins	-0.003	-0.031, 0.025	-0.23	0.817	0.973
rh_G_insular_short	-0.003	-0.027, 0.021	-0.25	0.806	0.973
rh_G_occipital_middle	0.014	-0.015, 0.043	0.99	0.325	0.973
rh_G_occipital_sup	-0.005	-0.035, 0.025	-0.31	0.754	0.973
rh_G_oc.temp_lat.fusifor	0.011	-0.023, 0.044	0.63	0.532	0.973
rh_G_oc.temp_med.Lingual	-0.017	-0.061, 0.027	-0.78	0.440	0.973
rh_G_oc.temp_med.Parahip	-0.018	-0.05, 0.014	-1.14	0.257	0.973
rh_G_orbital	0.003	-0.046, 0.052	0.12	0.905	0.973
rh_G_pariet_inf.Angular	-0.003	-0.031, 0.026	-0.21	0.838	0.973
rh_G_pariet_inf.Supramar	0.007	-0.019, 0.032	0.51	0.615	0.973
rh_G_parietal_sup	-0.002	-0.029, 0.026	-0.13	0.898	0.973
rh_G_postcentral	0.013	-0.017, 0.043	0.87	0.386	0.973
rh_G_precentral	0.006	-0.015, 0.028	0.59	0.560	0.973
rh_G_precuneus	0.013	-0.015, 0.042	0.94	0.349	0.973
rh_G_rectus	-0.050	-0.164, 0.064	-0.87	0.384	0.973
rh_G_subcallosal	-0.001	-0.053, 0.052	-0.03	0.973	0.985
rh_G_temp_sup.G_T_transv	-0.002	-0.03, 0.025	-0.17	0.863	0.973
rh_G_temp_sup.Lateral	-0.002	-0.024, 0.021	-0.15	0.880	0.973
rh_G_temp_sup.Plan_polar	0.014	-0.007, 0.035	1.31	0.192	0.973
rh_G_temp_sup.Plan_tempo	0.003	-0.024, 0.03	0.22	0.824	0.973
rh_G_temporal_inf	-0.002	-0.041, 0.037	-0.10	0.917	0.973
rh_G_temporal_middle	-0.013	-0.04, 0.014	-0.94	0.352	0.973
rh_Lat_Fis.ant.Horizontal	0.013	-0.024, 0.049	0.68	0.498	0.973
rh_Lat_Fis.ant.Vertical	-0.029	-0.069, 0.012	-1.40	0.165	0.973
rh_Lat_Fis.post	-0.003	-0.037, 0.031	-0.18	0.856	0.973
rh_Pole_occipital	0.023	-0.028, 0.074	0.91	0.367	0.973
rh_Pole_temporal	0.025	-0.014, 0.064	1.29	0.201	0.973
rh_S_calcarine	-0.011	-0.052, 0.03	-0.53	0.596	0.973
rh_S_central	-0.005	-0.036, 0.026	-0.33	0.739	0.973
rh_S_cingul.Marginalis	0.005	-0.019, 0.03	0.44	0.663	0.973
rh_S_circular_insula_ant	-0.021	-0.052, 0.009	-1.38	0.172	0.973
rh_S_circular_insula_inf	0.004	-0.02, 0.027	0.31	0.756	0.973
rh_S_circular_insula_sup	-0.011	-0.04, 0.019	-0.73	0.464	0.973
rh_S_collat_transv_ant	-0.019	-0.058, 0.02	-0.96	0.338	0.973
rh_S_collat_transv_post	-0.017	-0.046, 0.013	-1.13	0.261	0.973
rh_S_front_inf	-0.005	-0.038, 0.028	-0.29	0.771	0.973
rh_S_front_middle	-0.001	-0.037, 0.034	-0.07	0.941	0.973
rh_S_front_sup	0.005	-0.019, 0.028	0.39	0.698	0.973
rh_S_interprim.Jensen	-0.008	-0.042, 0.026	-0.49	0.628	0.973
rh_S_intrapariet_and_P_trans	0.012	-0.018, 0.041	0.79	0.433	0.973
rh_S_oc_middle_and_Lunatus	-0.021	-0.058, 0.017	-1.09	0.276	0.973
rh_S_oc_sup_and_transversal	0.001	-0.031, 0.032	0.05	0.961	0.984


eTable 25. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for failed stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_occipital_ant	0.010	-0.021, 0.04	0.62	0.540	0.973
rh_S_oc.temp_lat	-0.006	-0.033, 0.02	-0.46	0.645	0.973
rh_S_oc.temp_med_and_Lingual	-0.004	-0.026, 0.018	-0.35	0.729	0.973
rh_S_orbital_lateral	-0.030	-0.076, 0.016	-1.30	0.198	0.973
rh_S_orbital_med.olfact	-0.003	-0.076, 0.07	-0.08	0.938	0.973
rh_S_orbital.H_Shaped	-0.010	-0.046, 0.027	-0.52	0.607	0.973
rh_S_parieto_occipital	-0.005	-0.032, 0.021	-0.40	0.687	0.973
rh_S_pericallosal	0.004	-0.032, 0.041	0.24	0.813	0.973
rh_S_postcentral	0.008	-0.016, 0.033	0.69	0.492	0.973
rh_S_precentral.inf.part	-0.014	-0.045, 0.018	-0.87	0.385	0.973
rh_S_precentral.sup.part	0.008	-0.017, 0.033	0.63	0.532	0.973
rh_S_suborbital	-0.032	-0.134, 0.07	-0.62	0.534	0.973
rh_S_subparietal	0.025	-0.007, 0.058	1.56	0.123	0.973
rh_S_temporal_inf	-0.041	-0.086, 0.004	-1.82	0.072	0.973
rh_S_temporal_sup	0.006	-0.016, 0.028	0.53	0.598	0.973
rh_S_temporal_transverse	-0.013	-0.048, 0.022	-0.72	0.471	0.973


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 36. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 37 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 37 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 37 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 38. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 39 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 39 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 39 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 40. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 41 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 41 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 41 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for failed stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Correct stop vs correct go

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.007	-0.026, 0.013	-0.68	0.499	0.985
Left.Thalamus.Proper	0.005	-0.006, 0.016	0.86	0.390	0.985
Left.Caudate	0.009	-0.007, 0.025	1.11	0.270	0.925
Left.Putamen	0.008	-0.006, 0.022	1.18	0.239	0.925
Left.Pallidum	0.003	-0.008, 0.015	0.57	0.569	0.985
Brain.Stem	-0.004	-0.016, 0.009	-0.62	0.537	0.985
Left.Hippocampus	0.005	-0.007, 0.017	0.81	0.418	0.985
Left.Amygdala	0.004	-0.011, 0.019	0.50	0.616	0.985
Left.Accumbens.area	0.011	-0.007, 0.028	1.22	0.224	0.925
Left.VentralDC	0.009	-0.004, 0.022	1.32	0.189	0.925
Right.Cerebellum.Cortex	0.005	-0.015, 0.025	0.52	0.604	0.985
Right.Thalamus.Proper	0.002	-0.01, 0.014	0.35	0.728	0.985
Right.Caudate	-0.002	-0.019, 0.015	-0.19	0.853	0.985
Right.Putamen	-0.001	-0.015, 0.013	-0.09	0.927	0.985
Right.Pallidum	0.005	-0.006, 0.016	0.90	0.366	0.985
Right.Hippocampus	0.002	-0.01, 0.014	0.31	0.758	0.985
Right.Amygdala	0.009	-0.007, 0.024	1.06	0.291	0.925
Right.Accumbens.area	0.004	-0.013, 0.021	0.44	0.662	0.985
Right.VentralDC	0.003	-0.01, 0.016	0.41	0.684	0.985
Cortical regions					
lh_G_and_S_frontomargin	-0.003	-0.03, 0.024	-0.21	0.836	0.985
lh_G_and_S_occipital_inf	0.001	-0.021, 0.023	0.10	0.921	0.985
lh_G_and_S_paracentral	-0.002	-0.016, 0.013	-0.26	0.799	0.985
lh_G_and_S_subcentral	0.015	0, 0.03	2.01	0.045	0.925
lh_G_and_S_transv_frontopol	-0.001	-0.026, 0.024	-0.07	0.944	0.985
lh_G_and_S_cingul.Ant	0.012	-0.003, 0.027	1.57	0.116	0.925
lh_G_and_S_cingul.Mid.Ant	0.010	-0.004, 0.023	1.40	0.164	0.925
lh_G_and_S_cingul.Mid.Post	0.008	-0.004, 0.019	1.30	0.194	0.925
lh_G_cingul.Post.dorsal	0.003	-0.012, 0.019	0.42	0.673	0.985
lh_G_cingul.Post.ventral	-0.003	-0.035, 0.029	-0.19	0.852	0.985
lh_G_cuneus	0.002	-0.019, 0.022	0.15	0.879	0.985
lh_G_front_inf.Opercular	0.004	-0.01, 0.017	0.54	0.591	0.985
lh_G_front_inf.Orbital	0.013	-0.009, 0.035	1.19	0.234	0.925
lh_G_front_inf.Triangul	0.007	-0.011, 0.025	0.74	0.459	0.985
lh_G_front_middle	0.003	-0.012, 0.017	0.34	0.731	0.985
lh_G_front_sup	0.009	-0.001, 0.02	1.70	0.090	0.925
lh_G_Ins_lg_and_S_cent_ins	0.007	-0.007, 0.021	1.02	0.310	0.925
lh_G_insular_short	0.007	-0.006, 0.019	1.05	0.292	0.925
lh_G_occipital_middle	0.012	-0.005, 0.03	1.37	0.170	0.925

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_occipital_sup	0.000	-0.018, 0.018	-0.01	0.993	0.993
lh_G_oc.temp_lat.fusifor	0.004	-0.014, 0.023	0.46	0.644	0.985
lh_G_oc.temp_med.Lingual	-0.001	-0.024, 0.022	-0.06	0.952	0.985
lh_G_oc.temp_med.Parahip	-0.003	-0.02, 0.014	-0.32	0.750	0.985
lh_G_orbital	0.011	-0.013, 0.035	0.90	0.369	0.985
lh_G_pariet_inf.Angular	0.009	-0.006, 0.025	1.19	0.233	0.925
lh_G_pariet_inf.Supramar	0.011	-0.002, 0.024	1.69	0.092	0.925
lh_G_parietal_sup	0.000	-0.014, 0.014	-0.04	0.970	0.988
lh_G_postcentral	0.012	-0.003, 0.027	1.63	0.105	0.925
lh_G_precentral	0.004	-0.007, 0.014	0.64	0.524	0.985
lh_G_precuneus	0.001	-0.014, 0.016	0.14	0.890	0.985
lh_G_rectus	-0.013	-0.055, 0.03	-0.57	0.566	0.985
lh_G_subcallosal	0.020	-0.004, 0.045	1.62	0.105	0.925
lh_G_temp_sup.G_T_transv	0.004	-0.011, 0.018	0.51	0.609	0.985
lh_G_temp_sup.Lateral	-0.002	-0.015, 0.011	-0.28	0.779	0.985
lh_G_temp_sup.Plan_polar	-0.002	-0.014, 0.01	-0.37	0.710	0.985
lh_G_temp_sup.Plan_tempo	0.010	-0.005, 0.024	1.33	0.185	0.925
lh_G_temporal_inf	0.022	0.001, 0.043	2.08	0.038	0.925
lh_G_temporal_middle	0.005	-0.01, 0.02	0.68	0.497	0.985
lh_Lat_Fis.ant.Horizontal	0.005	-0.014, 0.024	0.52	0.602	0.985
lh_Lat_Fis.ant.Vertical	-0.002	-0.022, 0.019	-0.18	0.861	0.985
lh_Lat_Fis.post	0.009	-0.005, 0.024	1.26	0.210	0.925
lh_Pole_occipital	0.004	-0.025, 0.033	0.27	0.785	0.985
lh_Pole_temporal	0.013	-0.007, 0.032	1.29	0.197	0.925
lh_S_calcarine	0.001	-0.019, 0.022	0.14	0.886	0.985
lh_S_central	0.014	-0.001, 0.028	1.90	0.059	0.925
lh_S_cingul.Marginalis	0.011	-0.003, 0.024	1.57	0.118	0.925
lh_S_circular_insula_ant	0.005	-0.011, 0.022	0.62	0.536	0.985
lh_S_circular_insula_inf	0.001	-0.012, 0.013	0.08	0.937	0.985
lh_S_circular_insula_sup	0.019	0.006, 0.033	2.76	0.006	0.925
lh_S_collat_transv_ant	-0.004	-0.026, 0.018	-0.33	0.739	0.985
lh_S_collat_transv_post	-0.003	-0.022, 0.017	-0.25	0.801	0.985
lh_S_front_inf	0.008	-0.007, 0.024	1.05	0.294	0.925
lh_S_front_middle	0.008	-0.009, 0.025	0.96	0.339	0.976
lh_S_front_sup	0.001	-0.011, 0.014	0.21	0.835	0.985
lh_S_interm_prim.Jensen	0.013	-0.006, 0.032	1.35	0.179	0.925
lh_S_intrapariet_and_P_trans	0.004	-0.012, 0.021	0.51	0.612	0.985
lh_S_oc_middle_and_Lunatus	0.002	-0.018, 0.023	0.22	0.824	0.985
lh_S_oc_sup_and_transversal	-0.012	-0.03, 0.006	-1.32	0.186	0.925
lh_S_occipital_ant	0.002	-0.014, 0.019	0.28	0.778	0.985
lh_S_oc.temp_lat	0.016	0.001, 0.031	2.14	0.033	0.925
lh_S_oc.temp_med_and_Lingual	0.007	-0.007, 0.021	0.93	0.354	0.985
lh_S_orbital_lateral	0.018	-0.004, 0.04	1.58	0.115	0.925

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_orbital_med.olfact	0.002	-0.036, 0.039	0.10	0.920	0.985
lh_S_orbital.H_Shaped	0.009	-0.01, 0.027	0.93	0.351	0.985
lh_S_parieto_occipital	0.004	-0.012, 0.021	0.55	0.583	0.985
lh_S_pericallosal	0.009	-0.009, 0.026	0.99	0.324	0.950
lh_S_postcentral	0.005	-0.007, 0.018	0.86	0.388	0.985
lh_S_precentral.inf.part	0.011	-0.005, 0.026	1.39	0.167	0.925
lh_S_precentral.sup.part	0.006	-0.007, 0.018	0.84	0.400	0.985
lh_S_suborbital	0.013	-0.021, 0.047	0.75	0.453	0.985
lh_S_subparietal	0.002	-0.017, 0.02	0.17	0.865	0.985
lh_S_temporal_inf	0.013	-0.006, 0.033	1.38	0.169	0.925
lh_S_temporal_sup	0.011	-0.003, 0.024	1.49	0.136	0.925
lh_S_temporal_transverse	0.001	-0.017, 0.02	0.15	0.878	0.985
rh_G_and_S_frontomargin	-0.004	-0.031, 0.023	-0.32	0.751	0.985
rh_G_and_S_occipital_inf	-0.004	-0.026, 0.018	-0.34	0.734	0.985
rh_G_and_S_paracentral	-0.004	-0.019, 0.011	-0.52	0.606	0.985
rh_G_and_S_subcentral	0.009	-0.006, 0.024	1.17	0.241	0.925
rh_G_and_S_transv_frontopol	0.008	-0.013, 0.029	0.76	0.451	0.985
rh_G_and_S_cingul.Ant	0.018	0.001, 0.034	2.13	0.033	0.925
rh_G_and_S_cingul.Mid.Ant	0.008	-0.006, 0.021	1.12	0.265	0.925
rh_G_and_S_cingul.Mid.Post	0.009	-0.002, 0.019	1.59	0.113	0.925
rh_G_cingul.Post.dorsal	0.001	-0.015, 0.016	0.12	0.908	0.985
rh_G_cingul.Post.ventral	0.001	-0.027, 0.029	0.05	0.961	0.985
rh_G_cuneus	0.008	-0.013, 0.028	0.71	0.476	0.985
rh_G_front_inf.Opercular	0.013	-0.001, 0.028	1.77	0.078	0.925
rh_G_front_inf.Orbital	-0.002	-0.025, 0.02	-0.20	0.840	0.985
rh_G_front_inf.Triangul	0.016	-0.001, 0.033	1.81	0.071	0.925
rh_G_front_middle	-0.001	-0.017, 0.015	-0.13	0.894	0.985
rh_G_front_sup	0.007	-0.004, 0.017	1.22	0.222	0.925
rh_G_Ins_lg_and_S_cent_ins	0.003	-0.012, 0.017	0.35	0.726	0.985
rh_G_insular_short	0.012	-0.001, 0.025	1.86	0.063	0.925
rh_G_occipital_middle	0.004	-0.012, 0.02	0.49	0.625	0.985
rh_G_occipital_sup	-0.005	-0.022, 0.013	-0.51	0.611	0.985
rh_G_oc.temp_lat.fusifor	-0.003	-0.021, 0.014	-0.36	0.716	0.985
rh_G_oc.temp_med.Lingual	0.001	-0.022, 0.023	0.07	0.947	0.985
rh_G_oc.temp_med.Parahip	0.001	-0.017, 0.02	0.13	0.896	0.985
rh_G_orbital	-0.002	-0.026, 0.021	-0.20	0.840	0.985
rh_G_pariet_inf.Angular	0.008	-0.008, 0.024	1.02	0.310	0.925
rh_G_pariet_inf.Supramar	0.004	-0.009, 0.018	0.61	0.545	0.985
rh_G_parietal_sup	0.003	-0.011, 0.018	0.48	0.633	0.985
rh_G_postcentral	0.006	-0.008, 0.019	0.83	0.410	0.985
rh_G_precentral	0.004	-0.006, 0.014	0.77	0.444	0.985
rh_G_precuneus	0.012	-0.003, 0.027	1.56	0.120	0.925
rh_G_rectus	0.007	-0.047, 0.061	0.25	0.803	0.985

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_subcallosal	0.022	-0.002, 0.046	1.78	0.076	0.925
rh_G_temp_sup.G_T_transv	0.007	-0.009, 0.022	0.86	0.393	0.985
rh_G_temp_sup.Lateral	0.004	-0.008, 0.017	0.67	0.506	0.985
rh_G_temp_sup.Plan_polar	0.001	-0.01, 0.012	0.14	0.887	0.985
rh_G_temp_sup.Plan_tempo	0.011	-0.004, 0.026	1.49	0.137	0.925
rh_G_temporal_inf	0.002	-0.02, 0.024	0.18	0.860	0.985
rh_G_temporal_middle	0.007	-0.009, 0.024	0.89	0.372	0.985
rh_Lat_Fis.ant.Horizontal	0.013	-0.006, 0.031	1.34	0.181	0.925
rh_Lat_Fis.ant.Vertical	0.008	-0.015, 0.031	0.67	0.501	0.985
rh_Lat_Fis.post	0.009	-0.006, 0.025	1.15	0.250	0.925
rh_Pole_occipital	-0.001	-0.028, 0.026	-0.09	0.927	0.985
rh_Pole_temporal	0.023	0.001, 0.044	2.07	0.039	0.925
rh_S_calcarine	-0.001	-0.021, 0.019	-0.06	0.950	0.985
rh_S_central	0.008	-0.007, 0.022	1.06	0.289	0.925
rh_S_cingul.Marginalis	0.004	-0.009, 0.017	0.64	0.525	0.985
rh_S_circular_insula_ant	0.009	-0.008, 0.025	1.03	0.305	0.925
rh_S_circular_insula_inf	0.001	-0.013, 0.014	0.09	0.929	0.985
rh_S_circular_insula_sup	0.005	-0.01, 0.019	0.63	0.529	0.985
rh_S_collat_transv_ant	-0.002	-0.023, 0.02	-0.14	0.892	0.985
rh_S_collat_transv_post	-0.012	-0.03, 0.006	-1.28	0.201	0.925
rh_S_front_inf	0.004	-0.014, 0.022	0.44	0.658	0.985
rh_S_front_middle	0.009	-0.007, 0.026	1.11	0.268	0.925
rh_S_front_sup	0.007	-0.006, 0.02	1.08	0.280	0.925
rh_S_interm_prim.Jensen	0.005	-0.014, 0.023	0.48	0.631	0.985
rh_S_intrapariet_and_P_trans	0.005	-0.011, 0.02	0.57	0.569	0.985
rh_S_oc_middle_and_Lunatus	-0.001	-0.021, 0.02	-0.05	0.956	0.985
rh_S_oc_sup_and_transversal	0.001	-0.016, 0.018	0.14	0.889	0.985
rh_S_occipital_ant	-0.005	-0.022, 0.013	-0.52	0.604	0.985
rh_S_oc.temp_lat	0.008	-0.007, 0.024	1.04	0.299	0.925
rh_S_oc.temp_med_and_Lingual	-0.001	-0.014, 0.013	-0.12	0.907	0.985
rh_S_orbital_lateral	0.016	-0.007, 0.039	1.38	0.169	0.925
rh_S_orbital_med.olfact	0.009	-0.031, 0.049	0.46	0.644	0.985
rh_S_orbital.H_Shaped	0.004	-0.016, 0.023	0.37	0.712	0.985
rh_S_parieto_occipital	0.004	-0.012, 0.02	0.45	0.650	0.985
rh_S_pericallosal	0.016	-0.004, 0.035	1.57	0.117	0.925
rh_S_postcentral	0.000	-0.012, 0.012	0.01	0.992	0.993
rh_S_precentral.inf.part	0.000	-0.017, 0.017	-0.02	0.984	0.993
rh_S_precentral.sup.part	0.004	-0.008, 0.017	0.64	0.524	0.985
rh_S_suborbital	0.006	-0.039, 0.052	0.27	0.784	0.985
rh_S_subparietal	0.005	-0.013, 0.023	0.51	0.612	0.985
rh_S_temporal_inf	0.005	-0.018, 0.028	0.41	0.685	0.985
rh_S_temporal_sup	0.006	-0.008, 0.019	0.79	0.432	0.985
rh_S_temporal_transverse	0.005	-0.015, 0.025	0.51	0.608	0.985

eTable 26. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
------	------	-------	---	---------	-------

Bold denotes group differences at p<0.05 uncorrected

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.003	-0.022, 0.016	-0.29	0.770	0.940
Left.Thalamus.Proper	-0.003	-0.014, 0.009	-0.44	0.658	0.940
Left.Caudate	-0.002	-0.019, 0.015	-0.23	0.818	0.940
Left.Putamen	0.006	-0.009, 0.02	0.77	0.445	0.917
Left.Pallidum	0.008	-0.003, 0.019	1.37	0.170	0.801
Brain.Stem	0.006	-0.006, 0.019	0.97	0.334	0.916
Left.Hippocampus	0.011	-0.001, 0.023	1.87	0.062	0.801
Left.Amygdala	0.008	-0.006, 0.022	1.17	0.242	0.877
Left.Accumbens.area	0.000	-0.017, 0.018	0.05	0.962	0.989
Left.VentralDC	0.000	-0.013, 0.014	0.07	0.948	0.989
Right.Cerebellum.Cortex	0.017	-0.003, 0.038	1.68	0.095	0.801
Right.Thalamus.Proper	0.001	-0.011, 0.013	0.23	0.819	0.940
Right.Caudate	-0.004	-0.023, 0.014	-0.47	0.638	0.940
Right.Putamen	-0.002	-0.017, 0.012	-0.30	0.761	0.940
Right.Pallidum	0.004	-0.007, 0.016	0.77	0.443	0.917
Right.Hippocampus	0.006	-0.006, 0.019	1.05	0.293	0.903
Right.Amygdala	0.019	0.004, 0.033	2.49	0.013	0.801
Right.Accumbens.area	0.003	-0.014, 0.021	0.37	0.712	0.940
Right.VentralDC	0.003	-0.011, 0.017	0.39	0.700	0.940
Cortical regions					
lh_G_and_S_frontomargin	0.019	-0.007, 0.046	1.44	0.150	0.801
lh_G_and_S_occipital_inf	-0.004	-0.027, 0.02	-0.32	0.746	0.940
lh_G_and_S_paracentral	0.004	-0.011, 0.019	0.58	0.564	0.937
lh_G_and_S_subcentral	0.003	-0.012, 0.019	0.39	0.696	0.940
lh_G_and_S_transv_frontopol	0.013	-0.012, 0.039	1.01	0.314	0.903
lh_G_and_S_cingul.Ant	0.009	-0.007, 0.025	1.08	0.281	0.903
lh_G_and_S_cingul.Mid.Ant	-0.001	-0.014, 0.012	-0.15	0.884	0.958
lh_G_and_S_cingul.Mid.Post	0.005	-0.007, 0.017	0.80	0.421	0.916
lh_G_cingul.Post.dorsal	0.012	-0.003, 0.028	1.57	0.116	0.801
lh_G_cingul.Post.ventral	0.012	-0.02, 0.043	0.72	0.472	0.932
lh_G_cuneus	-0.014	-0.034, 0.006	-1.34	0.180	0.801
lh_G_front_inf.Opercular	0.002	-0.012, 0.016	0.24	0.813	0.940
lh_G_front_inf.Orbital	0.010	-0.012, 0.032	0.86	0.390	0.916

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_front_inf.Triangul	0.006	-0.013, 0.026	0.64	0.521	0.937
lh_G_front_middle	0.001	-0.014, 0.016	0.12	0.907	0.972
lh_G_front_sup	0.002	-0.008, 0.013	0.46	0.642	0.940
lh_G_Ins_lg_and_S_cent_ins	0.015	0, 0.03	2.03	0.043	0.801
lh_G_insular_short	0.003	-0.01, 0.016	0.42	0.672	0.940
lh_G_occipital_middle	0.002	-0.016, 0.019	0.21	0.836	0.940
lh_G_occipital_sup	-0.003	-0.02, 0.014	-0.34	0.737	0.940
lh_G_oc.temp_lat.fusifor	0.009	-0.01, 0.027	0.92	0.356	0.916
lh_G_oc.temp_med.Lingual	0.008	-0.016, 0.032	0.63	0.529	0.937
lh_G_oc.temp_med.Parahip	0.007	-0.009, 0.023	0.84	0.399	0.916
lh_G_orbital	0.021	-0.003, 0.046	1.71	0.089	0.801
lh_G_pariet_inf.Angular	0.013	-0.003, 0.029	1.63	0.103	0.801
lh_G_pariet_inf.Supramar	0.007	-0.006, 0.021	1.04	0.297	0.903
lh_G_parietal_sup	0.009	-0.006, 0.023	1.21	0.225	0.855
lh_G_postcentral	-0.006	-0.021, 0.01	-0.71	0.481	0.932
lh_G_precentral	0.001	-0.011, 0.012	0.11	0.913	0.972
lh_G_precuneus	0.016	0.001, 0.031	2.04	0.042	0.801
lh_G_rectus	0.024	-0.023, 0.071	1.02	0.310	0.903
lh_G_subcallosal	0.004	-0.022, 0.029	0.28	0.778	0.940
lh_G_temp_sup.G_T_transv	0.014	0, 0.029	1.90	0.058	0.801
lh_G_temp_sup.Lateral	0.005	-0.008, 0.018	0.72	0.470	0.932
lh_G_temp_sup.Plan_polar	0.007	-0.004, 0.019	1.24	0.215	0.850
lh_G_temp_sup.Plan_tempo	0.006	-0.009, 0.021	0.80	0.423	0.916
lh_G_temporal_inf	0.007	-0.014, 0.028	0.63	0.532	0.937
lh_G_temporal_middle	0.012	-0.004, 0.027	1.45	0.147	0.801
lh_Lat_Fis.ant.Horizontal	-0.005	-0.024, 0.014	-0.55	0.586	0.937
lh_Lat_Fis.ant.Vertical	-0.005	-0.028, 0.018	-0.44	0.662	0.940
lh_Lat_Fis.post	0.012	-0.004, 0.028	1.43	0.154	0.801
lh_Pole_occipital	0.003	-0.023, 0.03	0.26	0.794	0.940
lh_Pole_temporal	0.017	-0.002, 0.036	1.78	0.076	0.801
lh_S_calcarine	-0.007	-0.028, 0.014	-0.67	0.502	0.932
lh_S_central	0.008	-0.008, 0.023	0.96	0.338	0.916
lh_S_cingul.Marginalis	0.011	-0.003, 0.026	1.56	0.119	0.801
lh_S_circular_insula_ant	-0.005	-0.023, 0.012	-0.59	0.555	0.937
lh_S_circular_insula_inf	0.004	-0.009, 0.017	0.61	0.539	0.937
lh_S_circular_insula_sup	0.006	-0.009, 0.022	0.84	0.401	0.916
lh_S_collat_transv_ant	0.003	-0.019, 0.025	0.30	0.766	0.940
lh_S_collat_transv_post	0.000	-0.019, 0.02	0.02	0.980	0.989
lh_S_front_inf	0.003	-0.014, 0.02	0.35	0.729	0.940
lh_S_front_middle	0.007	-0.01, 0.024	0.78	0.433	0.916
lh_S_front_sup	0.002	-0.011, 0.015	0.28	0.783	0.940
lh_S_interm_prim.Jensen	-0.002	-0.021, 0.018	-0.16	0.870	0.956
lh_S_intrapariet_and_P_trans	0.003	-0.014, 0.02	0.34	0.731	0.940

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_oc_middle_and_Lunatus	0.000	-0.021, 0.021	-0.02	0.983	0.989
lh_S_oc_sup_and_transversal	0.005	-0.012, 0.023	0.58	0.563	0.937
lh_S_occipital_ant	-0.001	-0.019, 0.016	-0.16	0.876	0.956
lh_S_oc.temp_lat	0.008	-0.007, 0.024	1.03	0.306	0.903
lh_S_oc.temp_med_and_Lingual	0.010	-0.005, 0.025	1.31	0.192	0.801
lh_S_orbital_lateral	0.006	-0.017, 0.03	0.53	0.595	0.937
lh_S_orbital_med.olfact	0.011	-0.028, 0.05	0.55	0.583	0.937
lh_S_orbital.H_Shaped	0.007	-0.012, 0.027	0.74	0.458	0.932
lh_S_parieto_occipital	0.012	-0.005, 0.028	1.41	0.159	0.801
lh_S_pericallosal	0.007	-0.011, 0.025	0.79	0.430	0.916
lh_S_postcentral	0.000	-0.013, 0.014	0.02	0.982	0.989
lh_S_precentral.inf.part	0.014	-0.003, 0.031	1.63	0.103	0.801
lh_S_precentral.sup.part	0.011	-0.003, 0.025	1.53	0.127	0.801
lh_S_suborbital	0.033	0, 0.067	1.94	0.053	0.801
lh_S_subparietal	0.014	-0.005, 0.033	1.42	0.158	0.801
lh_S_temporal_inf	0.010	-0.011, 0.031	0.95	0.343	0.916
lh_S_temporal_sup	0.008	-0.007, 0.022	1.05	0.295	0.903
lh_S_temporal_transverse	-0.002	-0.021, 0.017	-0.20	0.839	0.940
rh_G_and_S_frontomargin	-0.010	-0.038, 0.018	-0.69	0.491	0.932
rh_G_and_S_occipital_inf	-0.010	-0.033, 0.013	-0.85	0.398	0.916
rh_G_and_S_paracentral	0.013	-0.004, 0.029	1.53	0.127	0.801
rh_G_and_S_subcentral	0.011	-0.005, 0.027	1.34	0.182	0.801
rh_G_and_S_transv_frontopol	-0.001	-0.024, 0.023	-0.05	0.957	0.989
rh_G_and_S_cingul.Ant	0.011	-0.005, 0.027	1.33	0.184	0.801
rh_G_and_S_cingul.Mid.Ant	0.002	-0.012, 0.016	0.28	0.779	0.940
rh_G_and_S_cingul.Mid.Post	0.013	0.001, 0.025	2.15	0.032	0.801
rh_G_cingul.Post.dorsal	0.010	-0.005, 0.025	1.31	0.191	0.801
rh_G_cingul.Post.ventral	0.015	-0.012, 0.042	1.11	0.267	0.903
rh_G_cuneus	0.002	-0.019, 0.023	0.21	0.835	0.940
rh_G_front_inf.Opercular	0.006	-0.009, 0.022	0.79	0.429	0.916
rh_G_front_inf.Orbital	-0.024	-0.046, -0.003	-2.20	0.028	0.801
rh_G_front_inf.Triangul	0.007	-0.013, 0.026	0.68	0.497	0.932
rh_G_front_middle	-0.012	-0.028, 0.005	-1.42	0.157	0.801
rh_G_front_sup	0.003	-0.008, 0.014	0.51	0.613	0.940
rh_G_Ins_lg_and_S_cent_ins	0.007	-0.009, 0.022	0.88	0.381	0.916
rh_G_insular_short	-0.002	-0.016, 0.012	-0.25	0.800	0.940
rh_G_occipital_middle	-0.002	-0.018, 0.015	-0.20	0.839	0.940
rh_G_occipital_sup	0.012	-0.006, 0.029	1.27	0.205	0.835
rh_G_oc.temp_lat.fusifor	0.010	-0.008, 0.029	1.14	0.254	0.884
rh_G_oc.temp_med.Lingual	0.003	-0.021, 0.026	0.23	0.815	0.940
rh_G_oc.temp_med.Parahip	-0.005	-0.024, 0.013	-0.54	0.591	0.937
rh_G_orbital	-0.001	-0.028, 0.025	-0.10	0.919	0.972
rh_G_pariet_inf.Angular	0.003	-0.014, 0.02	0.35	0.727	0.940

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_pariet_inf.Supramar	0.003	-0.011, 0.017	0.39	0.695	0.940
rh_G_parietal_sup	0.001	-0.013, 0.015	0.16	0.870	0.956
rh_G_postcentral	0.005	-0.01, 0.021	0.69	0.493	0.932
rh_G_precentral	0.006	-0.006, 0.018	1.04	0.299	0.903
rh_G_precuneus	0.015	0.001, 0.03	2.06	0.040	0.801
rh_G_rectus	0.015	-0.045, 0.075	0.48	0.630	0.940
rh_G_subcallosal	0.011	-0.013, 0.036	0.91	0.361	0.916
rh_G_temp_sup.G_T_transv	0.005	-0.012, 0.021	0.56	0.577	0.937
rh_G_temp_sup.Lateral	0.005	-0.007, 0.018	0.83	0.405	0.916
rh_G_temp_sup.Plan_polar	0.001	-0.01, 0.012	0.10	0.919	0.972
rh_G_temp_sup.Plan_tempo	0.018	0.003, 0.033	2.36	0.019	0.801
rh_G_temporal_inf	-0.003	-0.026, 0.021	-0.22	0.830	0.940
rh_G_temporal_middle	0.011	-0.005, 0.027	1.39	0.165	0.801
rh_Lat_Fis.ant.Horizontal	-0.017	-0.036, 0.002	-1.71	0.088	0.801
rh_Lat_Fis.ant.Vertical	-0.001	-0.026, 0.024	-0.09	0.926	0.973
rh_Lat_Fis.post	0.011	-0.007, 0.029	1.23	0.219	0.850
rh_Pole_occipital	-0.011	-0.037, 0.015	-0.82	0.415	0.916
rh_Pole_temporal	0.004	-0.019, 0.027	0.31	0.754	0.940
rh_S_calcarine	0.002	-0.019, 0.024	0.22	0.826	0.940
rh_S_central	0.012	-0.004, 0.027	1.49	0.136	0.801
rh_S_cingul.Marginalis	0.010	-0.003, 0.024	1.49	0.138	0.801
rh_S_circular_insula_ant	-0.012	-0.029, 0.006	-1.34	0.182	0.801
rh_S_circular_insula_inf	-0.003	-0.016, 0.01	-0.44	0.663	0.940
rh_S_circular_insula_sup	0.000	-0.016, 0.015	-0.02	0.983	0.989
rh_S_collat_transv_ant	0.003	-0.021, 0.026	0.21	0.830	0.940
rh_S_collat_transv_post	0.003	-0.016, 0.022	0.30	0.762	0.940
rh_S_front_inf	-0.006	-0.025, 0.014	-0.58	0.563	0.937
rh_S_front_middle	0.002	-0.016, 0.02	0.20	0.845	0.941
rh_S_front_sup	0.000	-0.013, 0.013	-0.01	0.992	0.992
rh_S_interm_prim.Jensen	0.004	-0.016, 0.023	0.36	0.716	0.940
rh_S_intrapariet_and_P_trans	0.003	-0.013, 0.019	0.35	0.725	0.940
rh_S_oc_middle_and_Lunatus	-0.006	-0.025, 0.014	-0.56	0.573	0.937
rh_S_oc_sup_and_transversal	0.013	-0.005, 0.03	1.44	0.150	0.801
rh_S_occipital_ant	0.005	-0.013, 0.023	0.55	0.580	0.937
rh_S_oc.temp_lat	0.007	-0.008, 0.023	0.93	0.353	0.916
rh_S_oc.temp_med_and_Lingual	0.009	-0.006, 0.023	1.16	0.248	0.880
rh_S_orbital_lateral	-0.011	-0.036, 0.013	-0.94	0.350	0.916
rh_S_orbital_med.olfact	-0.010	-0.051, 0.031	-0.47	0.642	0.940
rh_S_orbital.H_Shaped	-0.011	-0.031, 0.009	-1.10	0.270	0.903
rh_S_parieto_occipital	0.016	-0.001, 0.032	1.90	0.058	0.801
rh_S_pericallosal	0.005	-0.014, 0.025	0.55	0.584	0.937
rh_S_postcentral	-0.002	-0.015, 0.011	-0.36	0.718	0.940
rh_S_precentral.inf.part	0.007	-0.012, 0.025	0.71	0.481	0.932

eTable 27. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_precentral.sup.part	0.006	-0.008, 0.019	0.81	0.419	0.916
rh_S_suborbital	0.021	-0.026, 0.068	0.90	0.370	0.916
rh_S_subparietal	0.018	0, 0.037	1.99	0.048	0.801
rh_S_temporal_inf	-0.018	-0.041, 0.006	-1.47	0.143	0.801
rh_S_temporal_sup	0.009	-0.006, 0.023	1.19	0.235	0.871
rh_S_temporal_transverse	0.003	-0.016, 0.023	0.31	0.756	0.940

Bold denotes group differences at p<0.05 uncorrected

eTable 28. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.023	-0.083, 0.037	-0.76	0.450	0.994
Left.Thalamus.Proper	0.007	-0.024, 0.037	0.42	0.672	0.994
Left.Caudate	-0.003	-0.043, 0.036	-0.17	0.869	0.994
Left.Putamen	-0.001	-0.037, 0.034	-0.08	0.933	0.999
Left.Pallidum	0.004	-0.027, 0.034	0.25	0.801	0.994
Brain.Stem	-0.007	-0.043, 0.029	-0.37	0.709	0.994
Left.Hippocampus	-0.025	-0.059, 0.009	-1.44	0.152	0.912
Left.Amygdala	-0.012	-0.048, 0.024	-0.65	0.516	0.994
Left.Accumbens.area	-0.008	-0.052, 0.036	-0.37	0.711	0.994
Left.VentralDC	-0.016	-0.053, 0.021	-0.87	0.385	0.994
Right.Cerebellum.Cortex	0.015	-0.041, 0.07	0.52	0.603	0.994
Right.Thalamus.Proper	-0.012	-0.045, 0.021	-0.72	0.472	0.994
Right.Caudate	-0.017	-0.062, 0.027	-0.76	0.446	0.994
Right.Putamen	-0.013	-0.048, 0.021	-0.77	0.446	0.994
Right.Pallidum	-0.018	-0.049, 0.012	-1.20	0.233	0.994
Right.Hippocampus	-0.009	-0.042, 0.024	-0.54	0.590	0.994
Right.Amygdala	-0.023	-0.059, 0.013	-1.25	0.214	0.994
Right.Accumbens.area	0.003	-0.046, 0.052	0.12	0.904	0.999
Right.VentralDC	-0.028	-0.066, 0.01	-1.45	0.151	0.912
Cortical regions					
lh_G_and_S_frontomargin	-0.019	-0.099, 0.061	-0.47	0.642	0.994
lh_G_and_S_occipital_inf	-0.020	-0.088, 0.049	-0.57	0.571	0.994
lh_G_and_S_paracentral	-0.014	-0.053, 0.025	-0.72	0.474	0.994
lh_G_and_S_subcentral	-0.034	-0.076, 0.008	-1.63	0.107	0.912
lh_G_and_S_transv_frontopol	-0.004	-0.094, 0.086	-0.09	0.925	0.999
lh_G_and_S_cingul.Ant	-0.022	-0.065, 0.022	-0.99	0.323	0.994
lh_G_and_S_cingul.Mid.Ant	-0.011	-0.047, 0.024	-0.63	0.532	0.994
lh_G_and_S_cingul.Mid.Post	-0.007	-0.039, 0.026	-0.40	0.689	0.994
lh_G_cingul.Post.dorsal	0.001	-0.036, 0.039	0.06	0.954	0.999
lh_G_cingul.Post.ventral	0.008	-0.076, 0.093	0.19	0.847	0.994

eTable 28. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_cuneus	-0.029	-0.084, 0.026	-1.06	0.291	0.994
lh_G_front_inf.Opercular	0.029	-0.01, 0.067	1.49	0.140	0.912
lh_G_front_inf.Orbital	-0.026	-0.086, 0.034	-0.85	0.396	0.994
lh_G_front_inf.Triangul	-0.005	-0.056, 0.045	-0.21	0.832	0.994
lh_G_front_middle	0.008	-0.036, 0.052	0.36	0.718	0.994
lh_G_front_sup	-0.021	-0.047, 0.005	-1.59	0.116	0.912
lh_G_Ins_lg_and_S_cent_ins	-0.017	-0.055, 0.021	-0.88	0.382	0.994
lh_G_insular_short	0.013	-0.019, 0.046	0.80	0.424	0.994
lh_G_occipital_middle	-0.015	-0.058, 0.028	-0.70	0.484	0.994
lh_G_occipital_sup	-0.005	-0.052, 0.043	-0.19	0.849	0.994
lh_G_oc.temp_lat.fusifor	-0.021	-0.07, 0.027	-0.87	0.386	0.994
lh_G_oc.temp_med.Lingual	-0.060	-0.131, 0.011	-1.69	0.095	0.912
lh_G_oc.temp_med.Parahip	0.006	-0.048, 0.061	0.23	0.821	0.994
lh_G_orbital	-0.013	-0.088, 0.061	-0.36	0.720	0.994
lh_G_pariet_inf.Angular	0.014	-0.032, 0.06	0.61	0.546	0.994
lh_G_pariet_inf.Supramar	0.002	-0.034, 0.039	0.13	0.899	0.999
lh_G_parietal_sup	0.001	-0.038, 0.041	0.05	0.957	0.999
lh_G_postcentral	-0.008	-0.047, 0.032	-0.40	0.691	0.994
lh_G_precentral	-0.010	-0.039, 0.018	-0.73	0.468	0.994
lh_G_precuneus	0.001	-0.037, 0.038	0.03	0.975	0.999
lh_G_rectus	-0.036	-0.196, 0.125	-0.44	0.659	0.994
lh_G_subcallosal	-0.039	-0.118, 0.039	-0.99	0.323	0.994
lh_G_temp_sup.G_T_transv	-0.016	-0.055, 0.023	-0.82	0.415	0.994
lh_G_temp_sup.Lateral	-0.019	-0.053, 0.014	-1.14	0.258	0.994
lh_G_temp_sup.Plan_polar	-0.027	-0.058, 0.004	-1.74	0.085	0.912
lh_G_temp_sup.Plan_tempo	0.000	-0.042, 0.042	0.00	0.999	0.999
lh_G_temporal_inf	0.048	-0.018, 0.114	1.44	0.153	0.912
lh_G_temporal_middle	0.003	-0.039, 0.046	0.15	0.883	0.997
lh_Lat_Fis.ant.Horizontal	-0.064	-0.118, -0.011	-2.39	0.019	0.641
lh_Lat_Fis.ant.Vertical	0.046	-0.02, 0.112	1.38	0.170	0.936
lh_Lat_Fis.post	-0.008	-0.05, 0.034	-0.37	0.714	0.994
lh_Pole_occipital	-0.036	-0.106, 0.035	-1.01	0.316	0.994
lh_Pole_temporal	-0.046	-0.102, 0.009	-1.66	0.100	0.912
lh_S_calcarine	-0.016	-0.077, 0.045	-0.52	0.606	0.994
lh_S_central	-0.020	-0.062, 0.022	-0.93	0.357	0.994
lh_S_cingul.Marginalis	0.002	-0.031, 0.034	0.10	0.919	0.999
lh_S_circular_insula_ant	-0.009	-0.061, 0.043	-0.34	0.731	0.994
lh_S_circular_insula_inf	-0.019	-0.053, 0.014	-1.14	0.257	0.994
lh_S_circular_insula_sup	0.000	-0.039, 0.038	-0.01	0.991	0.999
lh_S_collat_transv_ant	0.020	-0.055, 0.095	0.53	0.597	0.994
lh_S_collat_transv_post	-0.046	-0.109, 0.018	-1.42	0.158	0.912
lh_S_front_inf	0.000	-0.055, 0.056	0.01	0.994	0.999
lh_S_front_middle	-0.039	-0.092, 0.014	-1.46	0.147	0.912

eTable 28. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_front_sup	0.008	-0.03, 0.045	0.41	0.683	0.994
lh_S_interm_prim.Jensen	0.014	-0.04, 0.069	0.52	0.601	0.994
lh_S_intrapariet_and_P_trans	0.008	-0.041, 0.058	0.33	0.741	0.994
lh_S_oc_middle_and_Lunatus	-0.061	-0.116, -0.006	-2.19	0.031	0.865
lh_S_oc_sup_and_transversal	-0.009	-0.059, 0.041	-0.37	0.709	0.994
lh_S_occipital_ant	-0.008	-0.048, 0.032	-0.39	0.699	0.994
lh_S_oc.temp_lat	-0.024	-0.069, 0.021	-1.05	0.298	0.994
lh_S_oc.temp_med_and_Lingual	-0.006	-0.045, 0.034	-0.29	0.769	0.994
lh_S_orbital_lateral	0.000	-0.072, 0.071	-0.01	0.991	0.999
lh_S_orbital_med.olfact	0.085	-0.039, 0.209	1.36	0.179	0.936
lh_S_orbital.H_Shaped	-0.028	-0.088, 0.031	-0.95	0.345	0.994
lh_S_parieto_occipital	-0.004	-0.046, 0.038	-0.20	0.839	0.994
lh_S_pericallosal	-0.006	-0.056, 0.044	-0.23	0.819	0.994
lh_S_postcentral	-0.001	-0.04, 0.038	-0.05	0.957	0.999
lh_S_precentral.inf.part	0.018	-0.033, 0.069	0.70	0.483	0.994
lh_S_precentral.sup.part	0.004	-0.038, 0.045	0.18	0.859	0.994
lh_S_suborbital	-0.026	-0.133, 0.081	-0.49	0.628	0.994
lh_S_subparietal	-0.013	-0.063, 0.036	-0.53	0.595	0.994
lh_S_temporal_inf	0.025	-0.037, 0.087	0.80	0.426	0.994
lh_S_temporal_sup	-0.008	-0.043, 0.027	-0.46	0.649	0.994
lh_S_temporal_transverse	-0.074	-0.129, -0.02	-2.70	0.008	0.464
rh_G_and_S_frontomargin	-0.008	-0.103, 0.087	-0.17	0.867	0.994
rh_G_and_S_occipital_inf	-0.014	-0.086, 0.058	-0.40	0.693	0.994
rh_G_and_S_paracentral	-0.024	-0.065, 0.017	-1.16	0.250	0.994
rh_G_and_S_subcentral	-0.064	-0.105, -0.023	-3.09	0.003	0.439
rh_G_and_S_transv_frontopol	-0.033	-0.108, 0.042	-0.87	0.387	0.994
rh_G_and_S_cingul.Ant	-0.035	-0.084, 0.013	-1.44	0.154	0.912
rh_G_and_S_cingul.Mid.Ant	0.001	-0.038, 0.04	0.04	0.972	0.999
rh_G_and_S_cingul.Mid.Post	-0.004	-0.036, 0.029	-0.23	0.817	0.994
rh_G_cingul.Post.dorsal	0.003	-0.034, 0.04	0.17	0.869	0.994
rh_G_cingul.Post.ventral	-0.009	-0.072, 0.054	-0.28	0.780	0.994
rh_G_cuneus	-0.048	-0.104, 0.008	-1.69	0.095	0.912
rh_G_front_inf.Opercular	0.004	-0.032, 0.04	0.21	0.835	0.994
rh_G_front_inf.Orbital	-0.029	-0.094, 0.036	-0.90	0.372	0.994
rh_G_front_inf.Triangul	-0.013	-0.06, 0.034	-0.55	0.585	0.994
rh_G_front_middle	-0.022	-0.07, 0.027	-0.89	0.377	0.994
rh_G_front_sup	-0.019	-0.05, 0.012	-1.24	0.218	0.994
rh_G_Ins_lg_and_S_cent_ins	-0.028	-0.066, 0.01	-1.49	0.140	0.912
rh_G_insular_short	-0.001	-0.035, 0.034	-0.03	0.974	0.999
rh_G_occipital_middle	-0.009	-0.051, 0.034	-0.41	0.683	0.994
rh_G_occipital_sup	-0.014	-0.06, 0.032	-0.59	0.557	0.994
rh_G_oc.temp_lat.fusifor	-0.013	-0.073, 0.047	-0.42	0.673	0.994
rh_G_oc.temp_med.Lingual	-0.019	-0.092, 0.055	-0.50	0.617	0.994


eTable 28. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_oc.temp_med.Parahip	-0.021	-0.071, 0.029	-0.84	0.403	0.994
rh_G_orbital	-0.008	-0.084, 0.068	-0.21	0.837	0.994
rh_G_pariet_inf.Angular	-0.002	-0.047, 0.043	-0.09	0.930	0.999
rh_G_pariet_inf.Supramar	-0.013	-0.048, 0.021	-0.77	0.445	0.994
rh_G_parietal_sup	-0.007	-0.046, 0.032	-0.37	0.714	0.994
rh_G_postcentral	-0.038	-0.078, 0.001	-1.91	0.059	0.912
rh_G_precentral	-0.015	-0.046, 0.015	-0.99	0.324	0.994
rh_G_precuneus	0.018	-0.025, 0.061	0.82	0.414	0.994
rh_G_rectus	0.089	-0.086, 0.263	1.01	0.315	0.994
rh_G_subcallosal	-0.008	-0.086, 0.07	-0.20	0.838	0.994
rh_G_temp_sup.G_T_transv	-0.022	-0.069, 0.026	-0.91	0.367	0.994
rh_G_temp_sup.Lateral	-0.039	-0.071, -0.007	-2.43	0.017	0.641
rh_G_temp_sup.Plan_polar	0.007	-0.029, 0.042	0.38	0.707	0.994
rh_G_temp_sup.Plan_tempo	-0.023	-0.064, 0.017	-1.16	0.248	0.994
rh_G_temporal_inf	0.055	-0.011, 0.121	1.66	0.101	0.912
rh_G_temporal_middle	-0.005	-0.046, 0.036	-0.24	0.810	0.994
rh_Lat_Fis.ant.Horizontal	-0.035	-0.087, 0.016	-1.35	0.179	0.936
rh_Lat_Fis.ant.Vertical	0.003	-0.07, 0.076	0.08	0.937	0.999
rh_Lat_Fis.post	-0.037	-0.084, 0.01	-1.56	0.122	0.912
rh_Pole_occipital	-0.026	-0.105, 0.053	-0.65	0.517	0.994
rh_Pole_temporal	-0.016	-0.08, 0.048	-0.50	0.618	0.994
rh_S_calcarine	-0.025	-0.088, 0.039	-0.78	0.440	0.994
rh_S_central	-0.026	-0.072, 0.02	-1.13	0.262	0.994
rh_S_cingul.Marginalis	0.003	-0.032, 0.037	0.15	0.879	0.997
rh_S_circular_insula_ant	-0.019	-0.067, 0.03	-0.77	0.443	0.994
rh_S_circular_insula_inf	-0.028	-0.065, 0.008	-1.56	0.123	0.912
rh_S_circular_insula_sup	0.009	-0.028, 0.046	0.49	0.626	0.994
rh_S_collat_transv_ant	-0.010	-0.078, 0.057	-0.31	0.761	0.994
rh_S_collat_transv_post	-0.005	-0.06, 0.051	-0.17	0.869	0.994
rh_S_front_inf	-0.011	-0.06, 0.037	-0.45	0.652	0.994
rh_S_front_middle	-0.030	-0.085, 0.024	-1.11	0.272	0.994
rh_S_front_sup	0.006	-0.035, 0.046	0.28	0.782	0.994
rh_S_interm_prim.Jensen	0.000	-0.056, 0.057	0.02	0.987	0.999
rh_S_intrapariet_and_P_trans	0.031	-0.018, 0.08	1.27	0.206	0.994
rh_S_oc_middle_and_Lunatus	-0.050	-0.104, 0.003	-1.87	0.064	0.912
rh_S_oc_sup_and_transversal	-0.037	-0.079, 0.004	-1.78	0.079	0.912
rh_S_occipital_ant	-0.044	-0.089, 0.002	-1.89	0.062	0.912
rh_S_oc.temp_lat	-0.066	-0.112, -0.02	-2.86	0.005	0.439
rh_S_oc.temp_med_and_Lingual	-0.018	-0.061, 0.025	-0.82	0.416	0.994
rh_S_orbital_lateral	-0.049	-0.112, 0.014	-1.55	0.124	0.912
rh_S_orbital_med.olfact	-0.037	-0.17, 0.096	-0.55	0.583	0.994
rh_S_orbital.H_Shaped	0.035	-0.027, 0.098	1.12	0.266	0.994
rh_S_parieto_occipital	-0.006	-0.047, 0.036	-0.26	0.793	0.994


eTable 28. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for correct stops: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_pericallosal	-0.017	-0.068, 0.033	-0.69	0.493	0.994
rh_S_postcentral	-0.004	-0.04, 0.032	-0.23	0.819	0.994
rh_S_precentral.inf.part	0.011	-0.038, 0.06	0.45	0.655	0.994
rh_S_precentral.sup.part	0.014	-0.024, 0.053	0.74	0.459	0.994
rh_S_suborbital	-0.033	-0.185, 0.119	-0.43	0.668	0.994
rh_S_subparietal	-0.024	-0.076, 0.028	-0.91	0.366	0.994
rh_S_temporal_inf	-0.002	-0.073, 0.069	-0.06	0.955	0.999
rh_S_temporal_sup	-0.031	-0.064, 0.001	-1.90	0.060	0.912
rh_S_temporal_transverse	-0.028	-0.085, 0.029	-0.97	0.335	0.994


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 42. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 43 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 43 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 43 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 44. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 45 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 45 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 45 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 46. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 47 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 47 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 47 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for correct stops. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Emotional n-back

2-back vs 0-back

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	0.000	-0.031, 0.032	0.01	0.996	0.996
Left.Thalamus.Proper	0.005	-0.012, 0.021	0.55	0.581	0.928
Left.Caudate	-0.005	-0.028, 0.017	-0.48	0.633	0.928
Left.Putamen	-0.010	-0.03, 0.009	-1.05	0.296	0.925
Left.Pallidum	-0.003	-0.021, 0.014	-0.38	0.703	0.944
Brain.Stem	0.002	-0.018, 0.022	0.21	0.830	0.968
Left.Hippocampus	-0.006	-0.026, 0.013	-0.65	0.517	0.928
Left.Amygdala	0.009	-0.014, 0.032	0.78	0.437	0.925
Left.Accumbens.area	-0.002	-0.028, 0.024	-0.14	0.885	0.969
Left.VentralDC	-0.007	-0.027, 0.012	-0.77	0.444	0.925
Right.Cerebellum.Cortex	-0.002	-0.031, 0.026	-0.16	0.870	0.969
Right.Thalamus.Proper	-0.005	-0.023, 0.013	-0.57	0.570	0.928
Right.Caudate	-0.017	-0.042, 0.007	-1.38	0.169	0.925
Right.Putamen	-0.012	-0.033, 0.008	-1.20	0.231	0.925
Right.Pallidum	-0.009	-0.026, 0.007	-1.12	0.264	0.925
Right.Hippocampus	-0.005	-0.024, 0.014	-0.51	0.607	0.928
Right.Amygdala	0.001	-0.022, 0.025	0.12	0.907	0.969
Right.Accumbens.area	0.007	-0.019, 0.033	0.52	0.604	0.928
Right.VentralDC	-0.004	-0.024, 0.017	-0.37	0.711	0.944
Cortical regions					
lh_G_and_S_frontomargin	-0.016	-0.061, 0.03	-0.68	0.495	0.928
lh_G_and_S_occipital_inf	-0.020	-0.058, 0.017	-1.08	0.281	0.925
lh_G_and_S_paracentral	-0.006	-0.027, 0.016	-0.49	0.622	0.928
lh_G_and_S_subcentral	-0.012	-0.037, 0.012	-0.98	0.328	0.925
lh_G_and_S_transv_frontopol	-0.011	-0.056, 0.035	-0.46	0.647	0.937
lh_G_and_S_cingul.Ant	-0.010	-0.032, 0.013	-0.85	0.393	0.925
lh_G_and_S_cingul.Mid.Ant	-0.008	-0.029, 0.013	-0.76	0.446	0.925
lh_G_and_S_cingul.Mid.Post	-0.003	-0.021, 0.016	-0.27	0.784	0.968
lh_G_cingul.Post.dorsal	-0.001	-0.023, 0.021	-0.09	0.926	0.969
lh_G_cingul.Post.ventral	-0.024	-0.068, 0.021	-1.06	0.292	0.925
lh_G_cuneus	-0.021	-0.054, 0.012	-1.24	0.215	0.925
lh_G_front_inf.Opercular	0.003	-0.019, 0.025	0.28	0.779	0.968
lh_G_front_inf.Orbital	0.016	-0.018, 0.05	0.94	0.348	0.925
lh_G_front_inf.Triangul	0.006	-0.022, 0.035	0.43	0.671	0.942
lh_G_front_middle	-0.013	-0.038, 0.012	-1.03	0.302	0.925
lh_G_front_sup	-0.009	-0.025, 0.007	-1.12	0.263	0.925
lh_G_Ins_lg_and_S_cent_ins	-0.004	-0.026, 0.018	-0.33	0.743	0.948

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_insular_short	-0.002	-0.02, 0.016	-0.18	0.855	0.968
lh_G_occipital_middle	-0.024	-0.048, 0	-1.98	0.048	0.925
lh_G_occipital_sup	-0.008	-0.035, 0.019	-0.58	0.562	0.928
lh_G_oc.temp_lat.fusifor	0.000	-0.029, 0.028	-0.02	0.981	0.996
lh_G_oc.temp_med.Lingual	-0.039	-0.077, 0	-1.98	0.048	0.925
lh_G_oc.temp_med.Parahip	-0.006	-0.034, 0.021	-0.45	0.651	0.937
lh_G_orbital	-0.029	-0.07, 0.011	-1.44	0.152	0.925
lh_G_pariet_inf.Angular	-0.004	-0.029, 0.021	-0.34	0.735	0.948
lh_G_pariet_inf.Supramar	0.004	-0.015, 0.024	0.43	0.670	0.942
lh_G_parietal_sup	-0.023	-0.044, -0.001	-2.09	0.037	0.925
lh_G_postcentral	0.003	-0.021, 0.026	0.22	0.827	0.968
lh_G_precentral	0.001	-0.014, 0.017	0.18	0.858	0.968
lh_G_precuneus	-0.010	-0.032, 0.012	-0.92	0.356	0.925
lh_G_rectus	-0.033	-0.114, 0.049	-0.79	0.433	0.925
lh_G_subcallosal	-0.015	-0.056, 0.025	-0.74	0.458	0.925
lh_G_temp_sup.G_T_transv	-0.013	-0.036, 0.009	-1.15	0.251	0.925
lh_G_temp_sup.Lateral	-0.007	-0.026, 0.013	-0.68	0.498	0.928
lh_G_temp_sup.Plan_polar	-0.010	-0.028, 0.007	-1.13	0.259	0.925
lh_G_temp_sup.Plan_tempo	0.000	-0.022, 0.022	0.01	0.996	0.996
lh_G_temporal_inf	0.009	-0.027, 0.045	0.49	0.623	0.928
lh_G_temporal_middle	0.012	-0.012, 0.036	0.98	0.327	0.925
lh_Lat_Fis.ant.Horizontal	-0.011	-0.042, 0.02	-0.72	0.473	0.925
lh_Lat_Fis.ant.Vertical	0.006	-0.028, 0.041	0.36	0.718	0.944
lh_Lat_Fis.post	0.007	-0.017, 0.031	0.58	0.563	0.928
lh_Pole_occipital	-0.033	-0.076, 0.009	-1.54	0.125	0.925
lh_Pole_temporal	-0.002	-0.034, 0.029	-0.13	0.897	0.969
lh_S_calcarine	-0.030	-0.063, 0.002	-1.85	0.065	0.925
lh_S_central	0.000	-0.023, 0.024	0.02	0.980	0.996
lh_S_cingul.Marginalis	0.005	-0.015, 0.024	0.49	0.623	0.928
lh_S_circular_insula_ant	-0.010	-0.037, 0.017	-0.73	0.466	0.925
lh_S_circular_insula_inf	0.003	-0.016, 0.022	0.28	0.780	0.968
lh_S_circular_insula_sup	-0.001	-0.021, 0.019	-0.10	0.918	0.969
lh_S_collat_transv_ant	0.013	-0.025, 0.05	0.66	0.509	0.928
lh_S_collat_transv_post	0.004	-0.03, 0.037	0.21	0.834	0.968
lh_S_front_inf	-0.013	-0.041, 0.016	-0.88	0.381	0.925
lh_S_front_middle	-0.023	-0.051, 0.006	-1.57	0.117	0.925
lh_S_front_sup	-0.005	-0.026, 0.015	-0.52	0.607	0.928
lh_S_interm_prim.Jensen	-0.001	-0.03, 0.029	-0.06	0.953	0.984
lh_S_intrapariet_and_P_trans	-0.017	-0.042, 0.009	-1.28	0.203	0.925
lh_S_oc_middle_and_Lunatus	-0.021	-0.053, 0.011	-1.30	0.195	0.925
lh_S_oc_sup_and_transversal	-0.012	-0.038, 0.014	-0.89	0.377	0.925
lh_S_occipital_ant	-0.017	-0.04, 0.006	-1.42	0.155	0.925
lh_S_oc.temp_lat	-0.014	-0.038, 0.01	-1.16	0.248	0.925

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_oc.temp_med_and_Lingual	-0.003	-0.025, 0.019	-0.26	0.796	0.968
lh_S_orbital_lateral	-0.024	-0.065, 0.018	-1.13	0.259	0.925
lh_S_orbital_med.olfact	-0.009	-0.074, 0.057	-0.26	0.797	0.968
lh_S_orbital.H_Shaped	-0.009	-0.041, 0.023	-0.56	0.576	0.928
lh_S_parieto_occipital	-0.007	-0.03, 0.017	-0.57	0.567	0.928
lh_S_pericallosal	-0.008	-0.034, 0.018	-0.62	0.535	0.928
lh_S_postcentral	-0.002	-0.022, 0.017	-0.25	0.805	0.968
lh_S_precentral.inf.part	-0.008	-0.034, 0.019	-0.57	0.568	0.928
lh_S_precentral.sup.part	-0.012	-0.033, 0.009	-1.10	0.271	0.925
lh_S_suborbital	-0.027	-0.081, 0.028	-0.96	0.340	0.925
lh_S_subparietal	-0.011	-0.038, 0.016	-0.81	0.418	0.925
lh_S_temporal_inf	0.016	-0.019, 0.051	0.92	0.360	0.925
lh_S_temporal_sup	-0.003	-0.022, 0.015	-0.34	0.737	0.948
lh_S_temporal_transverse	-0.036	-0.065, -0.007	-2.43	0.015	0.925
rh_G_and_S_frontomargin	-0.012	-0.062, 0.037	-0.49	0.627	0.928
rh_G_and_S_occipital_inf	0.002	-0.035, 0.039	0.09	0.928	0.969
rh_G_and_S_paracentral	0.005	-0.019, 0.028	0.38	0.704	0.944
rh_G_and_S_subcentral	-0.018	-0.043, 0.007	-1.38	0.168	0.925
rh_G_and_S_transv_frontopol	-0.016	-0.055, 0.024	-0.78	0.438	0.925
rh_G_and_S_cingul.Ant	-0.010	-0.036, 0.015	-0.82	0.414	0.925
rh_G_and_S_cingul.Mid.Ant	0.004	-0.017, 0.024	0.36	0.716	0.944
rh_G_and_S_cingul.Mid.Post	0.007	-0.011, 0.025	0.73	0.467	0.925
rh_G_cingul.Post.dorsal	-0.003	-0.023, 0.018	-0.26	0.796	0.968
rh_G_cingul.Post.ventral	-0.020	-0.056, 0.017	-1.05	0.296	0.925
rh_G_cuneus	-0.029	-0.061, 0.003	-1.81	0.071	0.925
rh_G_front_inf.Opercular	-0.002	-0.024, 0.02	-0.18	0.854	0.968
rh_G_front_inf.Orbital	-0.010	-0.045, 0.025	-0.56	0.573	0.928
rh_G_front_inf.Triangul	0.003	-0.024, 0.03	0.22	0.822	0.968
rh_G_front_middle	-0.007	-0.033, 0.02	-0.50	0.620	0.928
rh_G_front_sup	-0.002	-0.018, 0.015	-0.19	0.847	0.968
rh_G_Ins_lg_and_S_cent_ins	-0.010	-0.033, 0.012	-0.91	0.362	0.925
rh_G_insular_short	-0.001	-0.021, 0.018	-0.10	0.918	0.969
rh_G_occipital_middle	-0.013	-0.037, 0.01	-1.10	0.270	0.925
rh_G_occipital_sup	-0.013	-0.039, 0.014	-0.92	0.360	0.925
rh_G_oc.temp_lat.fusifor	-0.003	-0.036, 0.029	-0.19	0.847	0.968
rh_G_oc.temp_med.Lingual	-0.021	-0.059, 0.016	-1.11	0.267	0.925
rh_G_oc.temp_med.Parahip	-0.015	-0.044, 0.014	-1.04	0.297	0.925
rh_G_orbital	-0.042	-0.082, -0.001	-2.00	0.047	0.925
rh_G_pariet_inf.Angular	-0.013	-0.038, 0.013	-0.96	0.336	0.925
rh_G_pariet_inf.Supramar	-0.007	-0.025, 0.012	-0.72	0.473	0.925
rh_G_parietal_sup	-0.016	-0.037, 0.005	-1.52	0.130	0.925
rh_G_postcentral	0.005	-0.018, 0.028	0.44	0.662	0.942
rh_G_precentral	0.008	-0.009, 0.025	0.94	0.350	0.925

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_precuneus	-0.008	-0.033, 0.016	-0.69	0.492	0.928
rh_G_rectus	0.016	-0.079, 0.111	0.33	0.743	0.948
rh_G_subcallosal	-0.019	-0.057, 0.019	-0.97	0.334	0.925
rh_G_temp_sup.G_T_transv	-0.019	-0.045, 0.006	-1.49	0.136	0.925
rh_G_temp_sup.Lateral	-0.012	-0.031, 0.007	-1.22	0.222	0.925
rh_G_temp_sup.Plan_polar	-0.011	-0.031, 0.008	-1.14	0.255	0.925
rh_G_temp_sup.Plan_tempo	-0.015	-0.038, 0.007	-1.32	0.189	0.925
rh_G_temporal_inf	0.002	-0.035, 0.04	0.11	0.911	0.969
rh_G_temporal_middle	0.000	-0.023, 0.022	-0.01	0.990	0.996
rh_Lat_Fis.ant.Horizontal	0.006	-0.022, 0.034	0.40	0.687	0.944
rh_Lat_Fis.ant.Vertical	0.001	-0.036, 0.039	0.06	0.954	0.984
rh_Lat_Fis.post	-0.010	-0.035, 0.016	-0.74	0.462	0.925
rh_Pole_occipital	-0.026	-0.068, 0.016	-1.21	0.229	0.925
rh_Pole_temporal	-0.002	-0.038, 0.035	-0.09	0.925	0.969
rh_S_calcarine	-0.031	-0.066, 0.003	-1.80	0.072	0.925
rh_S_central	0.005	-0.02, 0.029	0.38	0.704	0.944
rh_S_cingul.Marginalis	0.008	-0.012, 0.027	0.76	0.450	0.925
rh_S_circular_insula_ant	-0.012	-0.038, 0.015	-0.86	0.392	0.925
rh_S_circular_insula_inf	-0.007	-0.029, 0.014	-0.65	0.518	0.928
rh_S_circular_insula_sup	0.010	-0.011, 0.03	0.93	0.353	0.925
rh_S_collat_transv_ant	-0.031	-0.07, 0.008	-1.58	0.115	0.925
rh_S_collat_transv_post	-0.002	-0.034, 0.029	-0.14	0.886	0.969
rh_S_front_inf	-0.016	-0.042, 0.01	-1.22	0.224	0.925
rh_S_front_middle	-0.012	-0.041, 0.018	-0.79	0.432	0.925
rh_S_front_sup	-0.010	-0.032, 0.013	-0.85	0.397	0.925
rh_S_interm_prim.Jensen	-0.014	-0.046, 0.018	-0.85	0.397	0.925
rh_S_intrapariet_and_P_trans	-0.001	-0.026, 0.023	-0.11	0.911	0.969
rh_S_oc_middle_and_Lunatus	-0.023	-0.055, 0.008	-1.44	0.150	0.925
rh_S_oc_sup_and_transversal	-0.019	-0.044, 0.007	-1.46	0.146	0.925
rh_S_occipital_ant	-0.018	-0.043, 0.007	-1.41	0.159	0.925
rh_S_oc.temp_lat	-0.012	-0.037, 0.014	-0.90	0.367	0.925
rh_S_oc.temp_med_and_Lingual	-0.011	-0.035, 0.013	-0.90	0.366	0.925
rh_S_orbital_lateral	-0.017	-0.055, 0.02	-0.90	0.367	0.925
rh_S_orbital_med.olfact	-0.007	-0.075, 0.061	-0.21	0.833	0.968
rh_S_orbital.H_Shaped	0.008	-0.023, 0.039	0.53	0.597	0.928
rh_S_parieto_occipital	-0.007	-0.031, 0.016	-0.61	0.542	0.928
rh_S_pericallosal	-0.010	-0.038, 0.018	-0.71	0.476	0.925
rh_S_postcentral	0.005	-0.015, 0.025	0.51	0.607	0.928
rh_S_precentral.inf.part	-0.006	-0.033, 0.021	-0.41	0.680	0.944
rh_S_precentral.sup.part	0.005	-0.016, 0.027	0.48	0.634	0.928
rh_S_suborbital	-0.048	-0.122, 0.027	-1.26	0.210	0.925
rh_S_subparietal	-0.011	-0.036, 0.015	-0.82	0.412	0.925
rh_S_temporal_inf	-0.030	-0.066, 0.006	-1.65	0.101	0.925

eTable 29. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_sup	-0.016	-0.036, 0.003	-1.65	0.100	0.925
rh_S_temporal_transverse	-0.026	-0.057, 0.005	-1.65	0.100	0.925

Bold denotes group differences at p<0.05 uncorrected

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.011	-0.044, 0.022	-0.66	0.509	0.890
Left.Thalamus.Proper	-0.002	-0.018, 0.015	-0.18	0.855	0.992
Left.Caudate	0.000	-0.023, 0.023	0.02	0.982	0.992
Left.Putamen	0.000	-0.02, 0.02	-0.03	0.978	0.992
Left.Pallidum	0.009	-0.008, 0.027	1.03	0.306	0.890
Brain.Stem	0.000	-0.021, 0.021	0.01	0.990	0.992
Left.Hippocampus	-0.008	-0.028, 0.012	-0.78	0.438	0.890
Left.Amygdala	0.005	-0.02, 0.029	0.37	0.714	0.992
Left.Accumbens.area	0.000	-0.028, 0.028	0.01	0.989	0.992
Left.VentralDC	-0.015	-0.037, 0.007	-1.34	0.181	0.890
Right.Cerebellum.Cortex	-0.011	-0.044, 0.022	-0.67	0.501	0.890
Right.Thalamus.Proper	-0.007	-0.025, 0.011	-0.76	0.447	0.890
Right.Caudate	-0.016	-0.04, 0.008	-1.30	0.194	0.890
Right.Putamen	-0.008	-0.028, 0.012	-0.78	0.433	0.890
Right.Pallidum	-0.013	-0.03, 0.005	-1.44	0.151	0.890
Right.Hippocampus	-0.007	-0.027, 0.012	-0.74	0.459	0.890
Right.Amygdala	0.009	-0.015, 0.034	0.75	0.455	0.890
Right.Accumbens.area	-0.012	-0.039, 0.015	-0.88	0.379	0.890
Right.VentralDC	-0.023	-0.044, -0.001	-2.07	0.039	0.890
Cortical regions					
lh_G_and_S_frontomargin	-0.002	-0.048, 0.045	-0.08	0.939	0.992
lh_G_and_S_occipital_inf	-0.007	-0.044, 0.03	-0.38	0.704	0.992
lh_G_and_S_paracentral	0.000	-0.022, 0.023	0.04	0.970	0.992
lh_G_and_S_subcentral	-0.015	-0.039, 0.009	-1.23	0.218	0.890
lh_G_and_S_transv_frontopol	0.030	-0.018, 0.077	1.23	0.221	0.890
lh_G_and_S_cingul.Ant	-0.003	-0.029, 0.022	-0.25	0.805	0.992
lh_G_and_S_cingul.Mid.Ant	-0.012	-0.032, 0.008	-1.16	0.246	0.890
lh_G_and_S_cingul.Mid.Post	-0.011	-0.028, 0.007	-1.18	0.238	0.890
lh_G_cingul.Post.dorsal	-0.005	-0.027, 0.018	-0.41	0.681	0.992

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_cingul.Post.ventral	-0.002	-0.05, 0.045	-0.10	0.919	0.992
lh_G_cuneus	-0.030	-0.062, 0.003	-1.79	0.074	0.890
lh_G_front_inf.Opercular	0.013	-0.008, 0.034	1.18	0.237	0.890
lh_G_front_inf.Orbital	0.002	-0.031, 0.036	0.12	0.901	0.992
lh_G_front_inf.Triangul	0.001	-0.027, 0.03	0.10	0.919	0.992
lh_G_front_middle	-0.015	-0.04, 0.01	-1.16	0.248	0.890
lh_G_front_sup	-0.007	-0.024, 0.009	-0.89	0.375	0.890
lh_G_Ins_lg_and_S_cent_ins	-0.001	-0.024, 0.022	-0.07	0.948	0.992
lh_G_insular_short	0.004	-0.015, 0.023	0.41	0.685	0.992
lh_G_occipital_middle	-0.009	-0.033, 0.016	-0.70	0.485	0.890
lh_G_occipital_sup	-0.015	-0.042, 0.011	-1.12	0.262	0.890
lh_G_oc.temp_lat.fusifor	-0.002	-0.032, 0.029	-0.12	0.903	0.992
lh_G_oc.temp_med.Lingual	-0.031	-0.071, 0.008	-1.57	0.118	0.890
lh_G_oc.temp_med.Parahip	-0.003	-0.031, 0.025	-0.23	0.820	0.992
lh_G_orbital	-0.023	-0.067, 0.021	-1.02	0.307	0.890
lh_G_pariet_inf.Angular	0.006	-0.02, 0.032	0.47	0.640	0.967
lh_G_pariet_inf.Supramar	0.001	-0.018, 0.021	0.12	0.903	0.992
lh_G_parietal_sup	-0.008	-0.03, 0.015	-0.66	0.511	0.890
lh_G_postcentral	-0.017	-0.039, 0.006	-1.43	0.155	0.890
lh_G_precentral	-0.007	-0.024, 0.01	-0.78	0.435	0.890
lh_G_precuneus	-0.002	-0.025, 0.02	-0.22	0.828	0.992
lh_G_rectus	-0.032	-0.117, 0.053	-0.73	0.463	0.890
lh_G_subcallosal	-0.010	-0.052, 0.032	-0.47	0.637	0.967
lh_G_temp_sup.G_T_transv	-0.002	-0.025, 0.022	-0.14	0.890	0.992
lh_G_temp_sup.Lateral	-0.011	-0.03, 0.009	-1.08	0.282	0.890
lh_G_temp_sup.Plan_polar	-0.023	-0.042, -0.004	-2.40	0.017	0.890
lh_G_temp_sup.Plan_tempo	0.008	-0.015, 0.03	0.68	0.497	0.890
lh_G_temporal_inf	0.017	-0.021, 0.054	0.89	0.377	0.890
lh_G_temporal_middle	-0.003	-0.029, 0.023	-0.21	0.830	0.992
lh_Lat_Fis.ant.Horizontal	-0.013	-0.045, 0.018	-0.85	0.398	0.890
lh_Lat_Fis.ant.Vertical	0.026	-0.011, 0.062	1.38	0.170	0.890
lh_Lat_Fis.post	0.004	-0.02, 0.027	0.31	0.754	0.992
lh_Pole_occipital	-0.028	-0.075, 0.02	-1.15	0.250	0.890
lh_Pole_temporal	-0.008	-0.041, 0.025	-0.48	0.634	0.967
lh_S_calcarine	-0.009	-0.042, 0.025	-0.50	0.617	0.967
lh_S_central	0.001	-0.023, 0.025	0.08	0.938	0.992
lh_S_cingul.Marginalis	0.007	-0.014, 0.027	0.64	0.520	0.890
lh_S_circular_insula_ant	0.000	-0.027, 0.028	0.03	0.976	0.992
lh_S_circular_insula_inf	0.002	-0.019, 0.022	0.15	0.880	0.992
lh_S_circular_insula_sup	0.004	-0.017, 0.025	0.36	0.719	0.992
lh_S_collat_transv_ant	0.019	-0.021, 0.058	0.94	0.350	0.890
lh_S_collat_transv_post	-0.031	-0.065, 0.003	-1.80	0.072	0.890
lh_S_front_inf	0.010	-0.019, 0.039	0.65	0.516	0.890

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_front_middle	-0.027	-0.057, 0.003	-1.75	0.082	0.890
lh_S_front_sup	-0.001	-0.022, 0.02	-0.11	0.915	0.992
lh_S_interm_prim.Jensen	-0.003	-0.035, 0.03	-0.16	0.873	0.992
lh_S_intrapariet_and_P_trans	-0.001	-0.027, 0.025	-0.06	0.954	0.992
lh_S_oc_middle_and_Lunatus	-0.025	-0.058, 0.008	-1.51	0.133	0.890
lh_S_oc_sup_and_transversal	-0.003	-0.03, 0.024	-0.22	0.827	0.992
lh_S_occipital_ant	-0.008	-0.033, 0.017	-0.61	0.544	0.895
lh_S_oc.temp_lat	0.006	-0.018, 0.03	0.50	0.619	0.967
lh_S_oc.temp_med_and_Lingual	0.001	-0.023, 0.025	0.08	0.932	0.992
lh_S_orbital_lateral	-0.006	-0.047, 0.034	-0.31	0.760	0.992
lh_S_orbital_med.olfact	0.066	-0.004, 0.135	1.87	0.063	0.890
lh_S_orbital.H_Shaped	-0.019	-0.052, 0.014	-1.14	0.256	0.890
lh_S_parieto_occipital	0.000	-0.024, 0.024	-0.01	0.992	0.992
lh_S_pericallosal	-0.015	-0.042, 0.012	-1.08	0.279	0.890
lh_S_postcentral	-0.014	-0.035, 0.007	-1.31	0.191	0.890
lh_S_precentral.inf.part	0.011	-0.017, 0.04	0.78	0.437	0.890
lh_S_precentral.sup.part	-0.010	-0.031, 0.011	-0.92	0.356	0.890
lh_S_suborbital	0.020	-0.037, 0.077	0.69	0.491	0.890
lh_S_subparietal	-0.011	-0.038, 0.016	-0.80	0.426	0.890
lh_S_temporal_inf	0.004	-0.032, 0.04	0.24	0.812	0.992
lh_S_temporal_sup	-0.002	-0.023, 0.019	-0.19	0.846	0.992
lh_S_temporal_transverse	-0.019	-0.05, 0.013	-1.17	0.242	0.890
rh_G_and_S_frontomargin	-0.040	-0.092, 0.013	-1.49	0.138	0.890
rh_G_and_S_occipital_inf	-0.015	-0.053, 0.022	-0.81	0.417	0.890
rh_G_and_S_paracentral	-0.011	-0.035, 0.013	-0.91	0.363	0.890
rh_G_and_S_subcentral	-0.020	-0.045, 0.004	-1.63	0.105	0.890
rh_G_and_S_transv_frontopol	-0.031	-0.073, 0.01	-1.48	0.139	0.890
rh_G_and_S_cingul.Ant	-0.015	-0.043, 0.013	-1.05	0.295	0.890
rh_G_and_S_cingul.Mid.Ant	-0.011	-0.032, 0.01	-1.04	0.300	0.890
rh_G_and_S_cingul.Mid.Post	-0.015	-0.032, 0.003	-1.68	0.095	0.890
rh_G_cingul.Post.dorsal	-0.001	-0.022, 0.02	-0.11	0.913	0.992
rh_G_cingul.Post.ventral	-0.012	-0.05, 0.025	-0.64	0.522	0.890
rh_G_cuneus	-0.005	-0.036, 0.027	-0.29	0.773	0.992
rh_G_front_inf.Opercular	0.003	-0.018, 0.024	0.28	0.781	0.992
rh_G_front_inf.Orbital	0.005	-0.029, 0.039	0.27	0.789	0.992
rh_G_front_inf.Triangul	-0.010	-0.038, 0.017	-0.74	0.459	0.890
rh_G_front_middle	-0.010	-0.037, 0.017	-0.71	0.481	0.890
rh_G_front_sup	-0.013	-0.029, 0.004	-1.47	0.141	0.890
rh_G_Ins_lg_and_S_cent_ins	0.002	-0.02, 0.025	0.19	0.846	0.992
rh_G_insular_short	-0.008	-0.028, 0.012	-0.75	0.452	0.890
rh_G_occipital_middle	-0.004	-0.028, 0.021	-0.30	0.766	0.992
rh_G_occipital_sup	-0.006	-0.032, 0.021	-0.41	0.678	0.992
rh_G_oc.temp_lat.fusifor	-0.022	-0.054, 0.01	-1.35	0.179	0.890

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_oc.temp_med.Lingual	-0.019	-0.057, 0.019	-0.98	0.330	0.890
rh_G_oc.temp_med.Parahip	-0.017	-0.045, 0.012	-1.16	0.245	0.890
rh_G_orbital	-0.021	-0.064, 0.022	-0.98	0.329	0.890
rh_G_pariet_inf.Angular	-0.013	-0.038, 0.012	-1.00	0.316	0.890
rh_G_pariet_inf.Supramar	-0.008	-0.027, 0.012	-0.77	0.444	0.890
rh_G_parietal_sup	-0.014	-0.036, 0.008	-1.26	0.210	0.890
rh_G_postcentral	-0.024	-0.047, -0.002	-2.17	0.031	0.890
rh_G_precentral	-0.013	-0.03, 0.004	-1.52	0.130	0.890
rh_G_precuneus	0.005	-0.02, 0.029	0.37	0.710	0.992
rh_G_rectus	0.032	-0.073, 0.137	0.60	0.548	0.895
rh_G_subcallosal	0.007	-0.035, 0.048	0.32	0.753	0.992
rh_G_temp_sup.G_T_transv	0.008	-0.017, 0.033	0.64	0.523	0.890
rh_G_temp_sup.Lateral	-0.017	-0.035, 0.002	-1.80	0.073	0.890
rh_G_temp_sup.Plan_polar	-0.014	-0.033, 0.006	-1.40	0.163	0.890
rh_G_temp_sup.Plan_tempo	0.002	-0.02, 0.024	0.19	0.850	0.992
rh_G_temporal_inf	0.028	-0.014, 0.07	1.29	0.196	0.890
rh_G_temporal_middle	-0.022	-0.047, 0.003	-1.71	0.088	0.890
rh_Lat_Fis.ant.Horizontal	0.008	-0.023, 0.038	0.49	0.621	0.967
rh_Lat_Fis.ant.Vertical	0.012	-0.027, 0.051	0.60	0.552	0.895
rh_Lat_Fis.post	-0.009	-0.035, 0.017	-0.67	0.505	0.890
rh_Pole_occipital	-0.010	-0.054, 0.034	-0.46	0.642	0.967
rh_Pole_temporal	-0.012	-0.048, 0.025	-0.62	0.533	0.892
rh_S_calcarine	-0.009	-0.042, 0.024	-0.55	0.584	0.937
rh_S_central	-0.009	-0.033, 0.015	-0.75	0.456	0.890
rh_S_cingul.Marginalis	-0.002	-0.021, 0.018	-0.16	0.874	0.992
rh_S_circular_insula_ant	-0.010	-0.038, 0.018	-0.68	0.495	0.890
rh_S_circular_insula_inf	-0.009	-0.029, 0.012	-0.85	0.396	0.890
rh_S_circular_insula_sup	0.009	-0.011, 0.029	0.89	0.373	0.890
rh_S_collat_transv_ant	-0.013	-0.052, 0.026	-0.68	0.500	0.890
rh_S_collat_transv_post	-0.007	-0.039, 0.025	-0.43	0.670	0.992
rh_S_front_inf	0.001	-0.027, 0.029	0.06	0.949	0.992
rh_S_front_middle	-0.027	-0.056, 0.003	-1.77	0.077	0.890
rh_S_front_sup	-0.020	-0.043, 0.003	-1.70	0.091	0.890
rh_S_interm_prim.Jensen	0.004	-0.028, 0.035	0.22	0.824	0.992
rh_S_intrapariet_and_P_trans	0.009	-0.016, 0.035	0.72	0.471	0.890
rh_S_oc_middle_and_Lunatus	-0.039	-0.07, -0.007	-2.41	0.016	0.890
rh_S_oc_sup_and_transversal	-0.019	-0.044, 0.006	-1.51	0.132	0.890
rh_S_occipital_ant	-0.009	-0.032, 0.015	-0.73	0.466	0.890
rh_S_oc.temp_lat	-0.029	-0.054, -0.004	-2.27	0.024	0.890
rh_S_oc.temp_med_and_Lingual	-0.004	-0.029, 0.02	-0.35	0.729	0.992
rh_S_orbital_lateral	-0.015	-0.051, 0.021	-0.80	0.423	0.890
rh_S_orbital_med.olfact	-0.006	-0.074, 0.062	-0.18	0.857	0.992
rh_S_orbital.H_Shaped	0.002	-0.032, 0.035	0.10	0.922	0.992

eTable 30. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_parieto_occipital	-0.008	-0.031, 0.015	-0.68	0.499	0.890
rh_S_pericallosal	-0.010	-0.04, 0.02	-0.67	0.502	0.890
rh_S_postcentral	-0.014	-0.034, 0.006	-1.34	0.180	0.890
rh_S_precentral.inf.part	0.010	-0.017, 0.037	0.74	0.460	0.890
rh_S_precentral.sup.part	-0.004	-0.024, 0.017	-0.34	0.737	0.992
rh_S_suborbital	0.007	-0.075, 0.088	0.16	0.873	0.992
rh_S_subparietal	-0.010	-0.038, 0.017	-0.73	0.465	0.890
rh_S_temporal_inf	-0.004	-0.046, 0.039	-0.18	0.853	0.992
rh_S_temporal_sup	-0.015	-0.034, 0.004	-1.58	0.116	0.890
rh_S_temporal_transverse	-0.010	-0.042, 0.022	-0.62	0.534	0.892

Bold denotes group differences at $p < 0.05$ uncorrected

eTable 31. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.011	-0.067, 0.046	-0.38	0.706	0.984
Left.Thalamus.Proper	0.016	-0.013, 0.045	1.09	0.280	0.984
Left.Caudate	0.002	-0.034, 0.037	0.09	0.929	0.984
Left.Putamen	0.007	-0.025, 0.04	0.46	0.650	0.984
Left.Pallidum	0.015	-0.014, 0.043	1.00	0.320	0.984
Brain.Stem	0.004	-0.031, 0.04	0.24	0.812	0.984
Left.Hippocampus	-0.008	-0.041, 0.025	-0.47	0.641	0.984
Left.Amygdala	-0.007	-0.042, 0.029	-0.36	0.717	0.984
Left.Accumbens.area	-0.014	-0.055, 0.027	-0.70	0.488	0.984
Left.VentralDC	-0.002	-0.038, 0.033	-0.13	0.895	0.984
Right.Cerebellum.Cortex	0.018	-0.035, 0.071	0.69	0.491	0.984
Right.Thalamus.Proper	-0.006	-0.036, 0.025	-0.36	0.719	0.984
Right.Caudate	-0.009	-0.05, 0.032	-0.44	0.661	0.984
Right.Putamen	-0.004	-0.037, 0.029	-0.23	0.822	0.984
Right.Pallidum	-0.016	-0.046, 0.013	-1.09	0.280	0.984
Right.Hippocampus	0.008	-0.024, 0.04	0.50	0.617	0.984
Right.Amygdala	-0.011	-0.048, 0.026	-0.59	0.558	0.984
Right.Accumbens.area	-0.004	-0.05, 0.043	-0.15	0.880	0.984
Right.VentralDC	-0.014	-0.049, 0.022	-0.78	0.440	0.984

eTable 31. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Cortical regions					
lh_G_and_S_frontomargin	-0.021	-0.107, 0.065	-0.48	0.629	0.984
lh_G_and_S_occipital_inf	-0.010	-0.073, 0.053	-0.31	0.757	0.984
lh_G_and_S_paracentral	-0.015	-0.052, 0.021	-0.83	0.409	0.984
lh_G_and_S_subcentral	-0.019	-0.06, 0.021	-0.94	0.349	0.984
lh_G_and_S_transv_frontopol	0.016	-0.074, 0.105	0.35	0.728	0.984
lh_G_and_S_cingul.Ant	-0.011	-0.051, 0.028	-0.57	0.568	0.984
lh_G_and_S_cingul.Mid.Ant	-0.003	-0.036, 0.03	-0.18	0.859	0.984
lh_G_and_S_cingul.Mid.Post	-0.001	-0.031, 0.029	-0.06	0.949	0.984
lh_G_cingul.Post.dorsal	0.006	-0.027, 0.04	0.38	0.708	0.984
lh_G_cingul.Post.ventral	0.009	-0.07, 0.087	0.22	0.826	0.984
lh_G_cuneus	-0.029	-0.081, 0.023	-1.10	0.275	0.984
lh_G_front_inf.Opercular	0.032	-0.005, 0.069	1.71	0.091	0.984
lh_G_front_inf.Orbital	-0.028	-0.089, 0.033	-0.91	0.364	0.984
lh_G_front_inf.Triangul	-0.012	-0.059, 0.036	-0.48	0.631	0.984
lh_G_front_middle	0.019	-0.021, 0.06	0.95	0.346	0.984
lh_G_front_sup	-0.018	-0.045, 0.009	-1.33	0.188	0.984
lh_G_Ins_lg_and_S_cent_ins	-0.010	-0.048, 0.027	-0.54	0.589	0.984
lh_G_insular_short	0.007	-0.024, 0.039	0.46	0.647	0.984
lh_G_occipital_middle	-0.023	-0.064, 0.019	-1.08	0.283	0.984
lh_G_occipital_sup	-0.007	-0.051, 0.038	-0.30	0.762	0.984
lh_G_oc.temp_lat.fusifor	-0.016	-0.061, 0.028	-0.73	0.465	0.984
lh_G_oc.temp_med.Lingual	-0.037	-0.102, 0.029	-1.12	0.267	0.984
lh_G_oc.temp_med.Parahip	0.003	-0.05, 0.056	0.12	0.908	0.984
lh_G_orbital	-0.012	-0.08, 0.057	-0.33	0.740	0.984
lh_G_pariet_inf.Angular	0.025	-0.018, 0.068	1.17	0.246	0.984
lh_G_pariet_inf.Supramar	0.009	-0.026, 0.043	0.49	0.626	0.984
lh_G_parietal_sup	0.001	-0.036, 0.037	0.04	0.967	0.989
lh_G_postcentral	-0.009	-0.047, 0.029	-0.46	0.650	0.984
lh_G_precentral	-0.008	-0.037, 0.02	-0.58	0.562	0.984
lh_G_precuneus	0.011	-0.025, 0.047	0.61	0.545	0.984
lh_G_rectus	-0.006	-0.15, 0.138	-0.09	0.930	0.984
lh_G_subcallosal	-0.044	-0.113, 0.025	-1.27	0.208	0.984
lh_G_temp_sup.G_T_transv	-0.012	-0.051, 0.027	-0.62	0.533	0.984
lh_G_temp_sup.Lateral	-0.025	-0.059, 0.008	-1.50	0.137	0.984
lh_G_temp_sup.Plan_polar	-0.023	-0.053, 0.007	-1.52	0.131	0.984
lh_G_temp_sup.Plan_tempo	0.005	-0.034, 0.043	0.24	0.812	0.984
lh_G_temporal_inf	0.042	-0.019, 0.104	1.36	0.176	0.984
lh_G_temporal_middle	0.011	-0.03, 0.052	0.53	0.599	0.984
lh_Lat_Fis.ant.Horizontal	-0.050	-0.101, 0.001	-1.96	0.053	0.984
lh_Lat_Fis.ant.Vertical	0.052	-0.006, 0.11	1.76	0.081	0.984
lh_Lat_Fis.post	-0.001	-0.04, 0.038	-0.04	0.970	0.989
lh_Pole_occipital	-0.034	-0.105, 0.036	-0.97	0.335	0.984

eTable 31. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_Pole_temporal	-0.038	-0.098, 0.023	-1.24	0.216	0.984
lh_S_calcarine	-0.012	-0.067, 0.043	-0.44	0.660	0.984
lh_S_central	-0.012	-0.051, 0.028	-0.58	0.560	0.984
lh_S_cingul.Marginalis	0.003	-0.027, 0.033	0.20	0.843	0.984
lh_S_circular_insula_ant	-0.008	-0.057, 0.042	-0.30	0.763	0.984
lh_S_circular_insula_inf	-0.016	-0.049, 0.017	-0.98	0.328	0.984
lh_S_circular_insula_sup	0.002	-0.033, 0.038	0.14	0.893	0.984
lh_S_collat_transv_ant	0.016	-0.05, 0.081	0.48	0.634	0.984
lh_S_collat_transv_post	-0.028	-0.087, 0.031	-0.93	0.355	0.984
lh_S_front_inf	0.004	-0.046, 0.054	0.16	0.871	0.984
lh_S_front_middle	-0.027	-0.077, 0.023	-1.07	0.287	0.984
lh_S_front_sup	0.018	-0.016, 0.052	1.04	0.302	0.984
lh_S_interm_prim.Jensen	0.027	-0.024, 0.077	1.05	0.297	0.984
lh_S_intrapariet_and_P_trans	0.012	-0.033, 0.058	0.53	0.596	0.984
lh_S_oc_middle_and_Lunatus	-0.048	-0.099, 0.003	-1.87	0.064	0.984
lh_S_oc_sup_and_transversal	0.003	-0.043, 0.05	0.14	0.888	0.984
lh_S_occipital_ant	0.001	-0.036, 0.039	0.08	0.940	0.984
lh_S_oc.temp_lat	-0.019	-0.06, 0.022	-0.91	0.363	0.984
lh_S_oc.temp_med_and_Lingual	-0.002	-0.039, 0.035	-0.11	0.911	0.984
lh_S_orbital_lateral	0.029	-0.041, 0.099	0.81	0.419	0.984
lh_S_orbital_med.olfact	0.080	-0.028, 0.187	1.47	0.144	0.984
lh_S_orbital.H_Shaped	-0.034	-0.089, 0.021	-1.22	0.226	0.984
lh_S_parieto_occipital	0.003	-0.035, 0.042	0.17	0.869	0.984
lh_S_pericallosal	-0.003	-0.048, 0.042	-0.13	0.894	0.984
lh_S_postcentral	0.005	-0.03, 0.041	0.30	0.762	0.984
lh_S_precentral.inf.part	0.028	-0.018, 0.074	1.20	0.233	0.984
lh_S_precentral.sup.part	0.006	-0.03, 0.042	0.34	0.737	0.984
lh_S_suborbital	-0.014	-0.112, 0.084	-0.29	0.773	0.984
lh_S_subparietal	-0.018	-0.06, 0.023	-0.88	0.380	0.984
lh_S_temporal_inf	0.027	-0.027, 0.082	1.00	0.320	0.984
lh_S_temporal_sup	0.007	-0.025, 0.04	0.44	0.660	0.984
lh_S_temporal_transverse	-0.063	-0.116, -0.011	-2.39	0.019	0.984
rh_G_and_S_frontomargin	0.031	-0.065, 0.127	0.64	0.524	0.984
rh_G_and_S_occipital_inf	-0.025	-0.093, 0.043	-0.73	0.464	0.984
rh_G_and_S_paracentral	-0.023	-0.066, 0.02	-1.07	0.285	0.984
rh_G_and_S_subcentral	-0.041	-0.083, 0.001	-1.95	0.054	0.984
rh_G_and_S_transv_frontopol	-0.025	-0.098, 0.048	-0.68	0.500	0.984
rh_G_and_S_cingul.Ant	-0.029	-0.072, 0.014	-1.33	0.188	0.984
rh_G_and_S_cingul.Mid.Ant	0.005	-0.03, 0.041	0.30	0.763	0.984
rh_G_and_S_cingul.Mid.Post	-0.002	-0.033, 0.028	-0.16	0.875	0.984
rh_G_cingul.Post.dorsal	0.014	-0.02, 0.049	0.83	0.409	0.984
rh_G_cingul.Post.ventral	0.007	-0.052, 0.065	0.23	0.819	0.984
rh_G_cuneus	-0.023	-0.075, 0.029	-0.88	0.384	0.984


eTable 31. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_front_inf.Opercular	0.013	-0.021, 0.047	0.77	0.444	0.984
rh_G_front_inf.Orbital	0.000	-0.065, 0.066	0.01	0.990	0.990
rh_G_front_inf.Triangul	0.002	-0.044, 0.049	0.10	0.919	0.984
rh_G_front_middle	-0.019	-0.065, 0.027	-0.84	0.405	0.984
rh_G_front_sup	-0.016	-0.048, 0.015	-1.02	0.312	0.984
rh_G_Ins_lg_and_S_cent_ins	-0.022	-0.06, 0.015	-1.19	0.236	0.984
rh_G_insular_short	0.000	-0.033, 0.032	-0.02	0.983	0.989
rh_G_occipital_middle	-0.002	-0.043, 0.039	-0.09	0.929	0.984
rh_G_occipital_sup	-0.002	-0.046, 0.043	-0.07	0.944	0.984
rh_G_oc.temp_lat.fusifor	-0.009	-0.063, 0.045	-0.33	0.745	0.984
rh_G_oc.temp_med.Lingual	-0.001	-0.068, 0.066	-0.03	0.978	0.989
rh_G_oc.temp_med.Parahip	-0.015	-0.063, 0.034	-0.61	0.544	0.984
rh_G_orbital	0.017	-0.057, 0.091	0.46	0.646	0.984
rh_G_pariet_inf.Angular	0.006	-0.037, 0.049	0.29	0.773	0.984
rh_G_pariet_inf.Supramar	0.002	-0.031, 0.035	0.11	0.909	0.984
rh_G_parietal_sup	0.004	-0.032, 0.041	0.24	0.814	0.984
rh_G_postcentral	-0.022	-0.061, 0.017	-1.13	0.260	0.984
rh_G_precentral	-0.010	-0.04, 0.021	-0.63	0.529	0.984
rh_G_precuneus	0.022	-0.017, 0.061	1.14	0.258	0.984
rh_G_rectus	0.116	-0.034, 0.265	1.54	0.127	0.984
rh_G_subcallosal	-0.014	-0.082, 0.054	-0.40	0.690	0.984
rh_G_temp_sup.G_T_transv	-0.010	-0.055, 0.036	-0.42	0.674	0.984
rh_G_temp_sup.Lateral	-0.034	-0.066, -0.001	-2.07	0.041	0.984
rh_G_temp_sup.Plan_polar	0.006	-0.027, 0.04	0.38	0.705	0.984
rh_G_temp_sup.Plan_tempo	-0.014	-0.052, 0.025	-0.71	0.478	0.984
rh_G_temporal_inf	0.066	0.004, 0.127	2.13	0.036	0.984
rh_G_temporal_middle	0.001	-0.039, 0.041	0.03	0.973	0.989
rh_Lat_Fis.ant.Horizontal	-0.014	-0.063, 0.035	-0.57	0.571	0.984
rh_Lat_Fis.ant.Vertical	0.006	-0.06, 0.072	0.19	0.849	0.984
rh_Lat_Fis.post	-0.032	-0.074, 0.01	-1.49	0.138	0.984
rh_Pole_occipital	-0.006	-0.083, 0.07	-0.16	0.872	0.984
rh_Pole_temporal	-0.032	-0.102, 0.037	-0.92	0.360	0.984
rh_S_calcarine	-0.012	-0.07, 0.045	-0.42	0.673	0.984
rh_S_central	-0.020	-0.061, 0.022	-0.94	0.350	0.984
rh_S_cingul.Marginalis	0.007	-0.025, 0.039	0.43	0.670	0.984
rh_S_circular_insula_ant	-0.009	-0.053, 0.035	-0.40	0.688	0.984
rh_S_circular_insula_inf	-0.021	-0.059, 0.018	-1.05	0.297	0.984
rh_S_circular_insula_sup	0.013	-0.019, 0.046	0.83	0.411	0.984
rh_S_collat_transv_ant	0.004	-0.057, 0.065	0.14	0.890	0.984
rh_S_collat_transv_post	0.011	-0.04, 0.062	0.43	0.665	0.984
rh_S_front_inf	-0.011	-0.056, 0.035	-0.47	0.639	0.984
rh_S_front_middle	-0.029	-0.081, 0.023	-1.12	0.265	0.984
rh_S_front_sup	0.011	-0.026, 0.048	0.57	0.572	0.984


eTable 31. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_interm_prim.Jensen	0.007	-0.045, 0.058	0.26	0.797	0.984
rh_S_intrapariet_and_P_trans	0.036	-0.009, 0.08	1.60	0.114	0.984
rh_S_oc_middle_and_Lunatus	-0.031	-0.082, 0.02	-1.21	0.230	0.984
rh_S_oc_sup_and_transversal	-0.024	-0.063, 0.014	-1.25	0.213	0.984
rh_S_occipital_ant	-0.023	-0.067, 0.021	-1.05	0.298	0.984
rh_S_oc.temp_lat	-0.056	-0.099, -0.013	-2.60	0.011	0.984
rh_S_oc.temp_med_and_Lingual	-0.002	-0.042, 0.039	-0.08	0.934	0.984
rh_S_orbital_lateral	-0.031	-0.093, 0.031	-1.00	0.321	0.984
rh_S_orbital_med.olfact	-0.026	-0.144, 0.092	-0.44	0.663	0.984
rh_S_orbital.H_Shaped	0.037	-0.021, 0.095	1.26	0.211	0.984
rh_S_parieto_occipital	0.004	-0.034, 0.042	0.20	0.843	0.984
rh_S_pericallosal	-0.002	-0.046, 0.042	-0.10	0.924	0.984
rh_S_postcentral	0.005	-0.029, 0.038	0.28	0.782	0.984
rh_S_precentral.inf.part	0.013	-0.031, 0.058	0.61	0.545	0.984
rh_S_precentral.sup.part	0.019	-0.016, 0.055	1.08	0.283	0.984
rh_S_suborbital	0.012	-0.12, 0.145	0.18	0.856	0.984
rh_S_subparietal	-0.005	-0.051, 0.04	-0.23	0.821	0.984
rh_S_temporal_inf	0.005	-0.059, 0.069	0.16	0.875	0.984
rh_S_temporal_sup	-0.015	-0.047, 0.016	-0.97	0.336	0.984
rh_S_temporal_transverse	-0.030	-0.084, 0.024	-1.11	0.271	0.984


Bold denotes group differences at p<0.05 uncorrected




eFigure 48. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 49 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 49 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 49 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 50. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 51 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 51 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 51 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 52. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 53 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 53 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 53 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for 2-back vs 0-back. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).

Emotional vs neutral faces

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test					
ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.032	-0.073, 0.009	-1.53	0.127	0.922
Left.Thalamus.Proper	0.001	-0.019, 0.02	0.05	0.957	0.994
Left.Caudate	-0.014	-0.042, 0.013	-1.02	0.310	0.993
Left.Putamen	-0.001	-0.026, 0.024	-0.10	0.923	0.993
Left.Pallidum	0.011	-0.01, 0.033	1.02	0.310	0.993
Brain.Stem	-0.025	-0.051, 0.001	-1.90	0.059	0.875
Left.Hippocampus	-0.020	-0.041, 0.001	-1.90	0.058	0.875
Left.Amygdala	0.005	-0.022, 0.031	0.34	0.737	0.993
Left.Accumbens.area	-0.002	-0.035, 0.031	-0.10	0.918	0.993
Left.VentralDC	-0.005	-0.028, 0.019	-0.39	0.697	0.993
Right.Cerebellum.Cortex	-0.006	-0.045, 0.034	-0.29	0.772	0.993
Right.Thalamus.Proper	-0.011	-0.033, 0.012	-0.94	0.347	0.993
Right.Caudate	-0.013	-0.041, 0.015	-0.90	0.371	0.993
Right.Putamen	-0.021	-0.047, 0.005	-1.62	0.107	0.875
Right.Pallidum	-0.001	-0.022, 0.021	-0.08	0.937	0.994
Right.Hippocampus	-0.005	-0.027, 0.018	-0.42	0.678	0.993
Right.Amygdala	-0.009	-0.037, 0.018	-0.66	0.511	0.993
Right.Accumbens.area	-0.018	-0.046, 0.01	-1.28	0.201	0.993
Right.VentralDC	-0.016	-0.039, 0.008	-1.29	0.197	0.993
Cortical regions					
lh_G_and_S_frontomargin	0.003	-0.05, 0.056	0.10	0.918	0.993
lh_G_and_S_occipital_inf	-0.024	-0.068, 0.02	-1.09	0.279	0.993
lh_G_and_S_paracentral	0.001	-0.024, 0.026	0.06	0.950	0.994
lh_G_and_S_subcentral	-0.025	-0.053, 0.002	-1.81	0.071	0.875
lh_G_and_S_transv_frontopol	-0.007	-0.055, 0.041	-0.29	0.772	0.993
lh_G_and_S_cingul.Ant	-0.002	-0.029, 0.025	-0.14	0.892	0.993
lh_G_and_S_cingul.Mid.Ant	-0.019	-0.041, 0.004	-1.64	0.102	0.875
lh_G_and_S_cingul.Mid.Post	0.001	-0.021, 0.022	0.06	0.951	0.994
lh_G_cingul.Post.dorsal	-0.009	-0.035, 0.018	-0.63	0.527	0.993
lh_G_cingul.Post.ventral	0.026	-0.026, 0.077	0.99	0.322	0.993
lh_G_cuneus	-0.017	-0.053, 0.02	-0.90	0.370	0.993
lh_G_front_inf.Opercular	-0.014	-0.038, 0.011	-1.12	0.264	0.993
lh_G_front_inf.Orbital	-0.035	-0.073, 0.004	-1.77	0.078	0.875
lh_G_front_inf.Triangul	-0.001	-0.034, 0.032	-0.05	0.964	0.994
lh_G_front_middle	-0.009	-0.036, 0.018	-0.64	0.524	0.993
lh_G_front_sup	0.006	-0.013, 0.026	0.62	0.534	0.993
lh_G_Ins_lg_and_S_cent_ins	0.003	-0.022, 0.028	0.23	0.819	0.993

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_insular_short	-0.005	-0.027, 0.017	-0.44	0.660	0.993
lh_G_occipital_middle	0.005	-0.024, 0.034	0.33	0.743	0.993
lh_G_occipital_sup	-0.003	-0.034, 0.029	-0.17	0.865	0.993
lh_G_oc.temp_lat.fusifor	-0.015	-0.05, 0.02	-0.86	0.393	0.993
lh_G_oc.temp_med.Lingual	-0.019	-0.061, 0.024	-0.86	0.390	0.993
lh_G_oc.temp_med.Parahip	0.013	-0.021, 0.046	0.75	0.455	0.993
lh_G_orbital	-0.008	-0.055, 0.04	-0.31	0.755	0.993
lh_G_pariet_inf.Angular	0.000	-0.026, 0.027	0.03	0.973	0.994
lh_G_pariet_inf.Supramar	-0.013	-0.036, 0.009	-1.19	0.236	0.993
lh_G_parietal_sup	-0.002	-0.027, 0.022	-0.19	0.852	0.993
lh_G_postcentral	-0.010	-0.036, 0.016	-0.75	0.453	0.993
lh_G_precentral	-0.011	-0.032, 0.009	-1.11	0.266	0.993
lh_G_precuneus	0.006	-0.018, 0.031	0.50	0.615	0.993
lh_G_rectus	-0.090	-0.182, 0.003	-1.91	0.057	0.875
lh_G_subcallosal	0.017	-0.026, 0.061	0.78	0.437	0.993
lh_G_temp_sup.G_T_transv	-0.007	-0.036, 0.022	-0.46	0.649	0.993
lh_G_temp_sup.Lateral	0.001	-0.021, 0.024	0.12	0.908	0.993
lh_G_temp_sup.Plan_polar	0.017	-0.004, 0.037	1.60	0.110	0.875
lh_G_temp_sup.Plan_tempo	-0.010	-0.036, 0.015	-0.80	0.425	0.993
lh_G_temporal_inf	0.003	-0.042, 0.047	0.12	0.907	0.993
lh_G_temporal_middle	-0.008	-0.038, 0.023	-0.49	0.628	0.993
lh_Lat_Fis.ant.Horizontal	-0.005	-0.039, 0.029	-0.29	0.769	0.993
lh_Lat_Fis.ant.Vertical	0.024	-0.011, 0.059	1.33	0.183	0.993
lh_Lat_Fis.post	-0.008	-0.035, 0.019	-0.59	0.553	0.993
lh_Pole_occipital	-0.005	-0.06, 0.05	-0.18	0.854	0.993
lh_Pole_temporal	-0.006	-0.044, 0.032	-0.32	0.751	0.993
lh_S_calcarine	0.011	-0.026, 0.047	0.58	0.562	0.993
lh_S_central	0.006	-0.021, 0.032	0.41	0.681	0.993
lh_S_cingul.Marginalis	-0.006	-0.026, 0.015	-0.52	0.603	0.993
lh_S_circular_insula_ant	-0.037	-0.067, -0.007	-2.46	0.015	0.875
lh_S_circular_insula_inf	-0.017	-0.04, 0.006	-1.44	0.151	0.993
lh_S_circular_insula_sup	-0.021	-0.046, 0.003	-1.75	0.081	0.875
lh_S_collat_transv_ant	-0.018	-0.063, 0.027	-0.79	0.428	0.993
lh_S_collat_transv_post	-0.033	-0.071, 0.005	-1.72	0.087	0.875
lh_S_front_inf	-0.003	-0.031, 0.024	-0.25	0.803	0.993
lh_S_front_middle	-0.011	-0.041, 0.018	-0.75	0.455	0.993
lh_S_front_sup	-0.019	-0.041, 0.003	-1.69	0.093	0.875
lh_S_interm_prim.Jensen	-0.004	-0.035, 0.026	-0.28	0.782	0.993
lh_S_intrapariet_and_P_trans	-0.011	-0.036, 0.014	-0.88	0.380	0.993
lh_S_oc_middle_and_Lunatus	-0.035	-0.075, 0.004	-1.76	0.080	0.875
lh_S_oc_sup_and_transversal	-0.016	-0.048, 0.016	-0.99	0.322	0.993
lh_S_occipital_ant	-0.014	-0.04, 0.013	-1.03	0.305	0.993
lh_S_oc.temp_lat	-0.038	-0.064, -0.012	-2.86	0.005	0.758

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_oc.temp_med_and_Lingual	-0.013	-0.04, 0.015	-0.91	0.364	0.993
lh_S_orbital_lateral	0.009	-0.035, 0.053	0.40	0.686	0.993
lh_S_orbital_med.olfact	-0.042	-0.118, 0.034	-1.08	0.281	0.993
lh_S_orbital.H_Shaped	-0.018	-0.052, 0.017	-1.00	0.316	0.993
lh_S_parieto_occipital	-0.003	-0.029, 0.022	-0.24	0.810	0.993
lh_S_pericallosal	-0.006	-0.035, 0.023	-0.42	0.673	0.993
lh_S_postcentral	-0.009	-0.032, 0.013	-0.80	0.425	0.993
lh_S_precentral.inf.part	-0.013	-0.041, 0.015	-0.90	0.368	0.993
lh_S_precentral.sup.part	-0.012	-0.036, 0.012	-0.98	0.326	0.993
lh_S_suborbital	-0.008	-0.066, 0.051	-0.25	0.799	0.993
lh_S_subparietal	0.007	-0.021, 0.034	0.48	0.630	0.993
lh_S_temporal_inf	0.007	-0.036, 0.049	0.32	0.747	0.993
lh_S_temporal_sup	-0.003	-0.027, 0.02	-0.27	0.786	0.993
lh_S_temporal_transverse	-0.004	-0.039, 0.031	-0.22	0.825	0.993
rh_G_and_S_frontomargin	-0.003	-0.06, 0.055	-0.09	0.927	0.993
rh_G_and_S_occipital_inf	-0.011	-0.055, 0.034	-0.48	0.633	0.993
rh_G_and_S_paracentral	-0.004	-0.031, 0.023	-0.27	0.784	0.993
rh_G_and_S_subcentral	0.003	-0.024, 0.03	0.21	0.837	0.993
rh_G_and_S_transv_frontopol	-0.005	-0.047, 0.036	-0.26	0.798	0.993
rh_G_and_S_cingul.Ant	-0.014	-0.044, 0.015	-0.97	0.332	0.993
rh_G_and_S_cingul.Mid.Ant	-0.020	-0.043, 0.004	-1.65	0.099	0.875
rh_G_and_S_cingul.Mid.Post	0.004	-0.017, 0.025	0.36	0.720	0.993
rh_G_cingul.Post.dorsal	-0.003	-0.027, 0.021	-0.24	0.810	0.993
rh_G_cingul.Post.ventral	-0.022	-0.065, 0.021	-1.00	0.318	0.993
rh_G_cuneus	-0.019	-0.055, 0.016	-1.06	0.290	0.993
rh_G_front_inf.Opercular	-0.008	-0.033, 0.018	-0.59	0.559	0.993
rh_G_front_inf.Orbital	-0.015	-0.055, 0.025	-0.74	0.460	0.993
rh_G_front_inf.Triangul	-0.013	-0.045, 0.019	-0.81	0.418	0.993
rh_G_front_middle	0.003	-0.025, 0.031	0.22	0.829	0.993
rh_G_front_sup	0.002	-0.017, 0.021	0.17	0.864	0.993
rh_G_Ins_lg_and_S_cent_ins	0.018	-0.009, 0.046	1.32	0.186	0.993
rh_G_insular_short	-0.007	-0.03, 0.016	-0.60	0.547	0.993
rh_G_occipital_middle	0.009	-0.021, 0.039	0.58	0.566	0.993
rh_G_occipital_sup	0.000	-0.032, 0.033	0.02	0.986	0.994
rh_G_oc.temp_lat.fusifor	0.002	-0.036, 0.04	0.10	0.919	0.993
rh_G_oc.temp_med.Lingual	-0.024	-0.067, 0.02	-1.07	0.287	0.993
rh_G_oc.temp_med.Parahip	-0.016	-0.055, 0.023	-0.83	0.408	0.993
rh_G_orbital	-0.021	-0.071, 0.029	-0.83	0.406	0.993
rh_G_pariet_inf.Angular	0.008	-0.017, 0.034	0.66	0.508	0.993
rh_G_pariet_inf.Supramar	-0.011	-0.032, 0.011	-0.99	0.324	0.993
rh_G_parietal_sup	-0.005	-0.03, 0.02	-0.39	0.694	0.993
rh_G_postcentral	-0.002	-0.028, 0.023	-0.19	0.847	0.993
rh_G_precentral	0.003	-0.017, 0.022	0.27	0.791	0.993

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_precuneus	0.004	-0.023, 0.03	0.27	0.785	0.993
rh_G_rectus	-0.105	-0.219, 0.01	-1.80	0.073	0.875
rh_G_subcallosal	0.009	-0.035, 0.054	0.41	0.680	0.993
rh_G_temp_sup.G_T_transv	-0.002	-0.031, 0.027	-0.14	0.892	0.993
rh_G_temp_sup.Lateral	-0.006	-0.027, 0.016	-0.52	0.603	0.993
rh_G_temp_sup.Plan_polar	0.007	-0.014, 0.028	0.66	0.508	0.993
rh_G_temp_sup.Plan_tempo	-0.015	-0.04, 0.01	-1.15	0.251	0.993
rh_G_temporal_inf	-0.040	-0.086, 0.007	-1.68	0.093	0.875
rh_G_temporal_middle	0.004	-0.023, 0.032	0.31	0.753	0.993
rh_Lat_Fis.ant.Horizontal	-0.005	-0.037, 0.027	-0.32	0.747	0.993
rh_Lat_Fis.ant.Vertical	-0.005	-0.042, 0.032	-0.26	0.797	0.993
rh_Lat_Fis.post	-0.003	-0.032, 0.025	-0.23	0.817	0.993
rh_Pole_occipital	-0.029	-0.084, 0.025	-1.05	0.293	0.993
rh_Pole_temporal	-0.011	-0.054, 0.032	-0.49	0.622	0.993
rh_S_calcarine	-0.012	-0.05, 0.026	-0.60	0.549	0.993
rh_S_central	0.010	-0.018, 0.037	0.67	0.501	0.993
rh_S_cingul.Marginalis	-0.008	-0.031, 0.014	-0.72	0.470	0.993
rh_S_circular_insula_ant	-0.032	-0.061, -0.002	-2.11	0.036	0.875
rh_S_circular_insula_inf	-0.004	-0.028, 0.02	-0.34	0.737	0.993
rh_S_circular_insula_sup	-0.015	-0.039, 0.01	-1.18	0.239	0.993
rh_S_collat_transv_ant	-0.039	-0.089, 0.011	-1.53	0.126	0.922
rh_S_collat_transv_post	-0.023	-0.059, 0.014	-1.21	0.228	0.993
rh_S_front_inf	0.000	-0.029, 0.029	-0.01	0.988	0.994
rh_S_front_middle	-0.027	-0.058, 0.003	-1.78	0.076	0.875
rh_S_front_sup	-0.009	-0.032, 0.013	-0.84	0.404	0.993
rh_S_interm_prim.Jensen	-0.017	-0.046, 0.012	-1.15	0.250	0.993
rh_S_intrapariet_and_P_trans	0.007	-0.019, 0.034	0.54	0.588	0.993
rh_S_oc_middle_and_Lunatus	-0.034	-0.07, 0.003	-1.83	0.068	0.875
rh_S_oc_sup_and_transversal	0.002	-0.027, 0.031	0.13	0.895	0.993
rh_S_occipital_ant	-0.002	-0.028, 0.024	-0.17	0.865	0.993
rh_S_oc.temp_lat	-0.027	-0.054, 0.001	-1.92	0.056	0.875
rh_S_oc.temp_med_and_Lingual	-0.010	-0.039, 0.018	-0.70	0.487	0.993
rh_S_orbital_lateral	0.000	-0.04, 0.04	0.00	1.000	1.000
rh_S_orbital_med.olfact	-0.051	-0.131, 0.029	-1.25	0.212	0.993
rh_S_orbital.H_Shaped	0.003	-0.035, 0.042	0.18	0.860	0.993
rh_S_parieto_occipital	0.000	-0.026, 0.026	0.03	0.978	0.994
rh_S_pericallosal	-0.011	-0.043, 0.022	-0.66	0.511	0.993
rh_S_postcentral	-0.013	-0.035, 0.01	-1.11	0.266	0.993
rh_S_precentral.inf.part	-0.003	-0.031, 0.025	-0.21	0.835	0.993
rh_S_precentral.sup.part	-0.015	-0.038, 0.008	-1.25	0.211	0.993
rh_S_suborbital	0.007	-0.085, 0.099	0.15	0.882	0.993
rh_S_subparietal	0.006	-0.022, 0.034	0.42	0.671	0.993
rh_S_temporal_inf	-0.005	-0.049, 0.039	-0.23	0.816	0.993

eTable 32. Differences in functional activity between the never-suicidal group and the child-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_temporal_sup	0.000	-0.023, 0.022	-0.02	0.987	0.994
rh_S_temporal_transverse	-0.012	-0.048, 0.024	-0.67	0.503	0.993

Bold denotes group differences at p<0.05 uncorrected

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.012	-0.052, 0.029	-0.56	0.573	0.940
Left.Thalamus.Proper	0.014	-0.005, 0.034	1.44	0.151	0.773
Left.Caudate	0.001	-0.027, 0.029	0.06	0.954	0.960
Left.Putamen	0.008	-0.017, 0.033	0.62	0.536	0.940
Left.Pallidum	0.016	-0.006, 0.038	1.47	0.144	0.773
Brain.Stem	-0.013	-0.037, 0.011	-1.05	0.293	0.843
Left.Hippocampus	-0.002	-0.023, 0.02	-0.14	0.888	0.940
Left.Amygdala	0.020	-0.008, 0.048	1.43	0.153	0.773
Left.Accumbens.area	0.014	-0.018, 0.046	0.87	0.384	0.916
Left.VentralDC	0.012	-0.014, 0.038	0.91	0.363	0.916
Right.Cerebellum.Cortex	0.004	-0.032, 0.039	0.19	0.846	0.940
Right.Thalamus.Proper	0.020	-0.001, 0.04	1.91	0.057	0.758
Right.Caudate	0.004	-0.025, 0.032	0.24	0.809	0.940
Right.Putamen	0.004	-0.021, 0.03	0.35	0.729	0.940
Right.Pallidum	0.002	-0.019, 0.024	0.22	0.825	0.940
Right.Hippocampus	0.019	-0.003, 0.042	1.68	0.094	0.773
Right.Amygdala	0.019	-0.011, 0.05	1.26	0.210	0.780
Right.Accumbens.area	0.009	-0.021, 0.039	0.60	0.548	0.940
Right.VentralDC	0.018	-0.006, 0.041	1.46	0.144	0.773
Cortical regions					
lh_G_and_S_frontomargin	-0.042	-0.098, 0.014	-1.49	0.137	0.773
lh_G_and_S_occipital_inf	-0.014	-0.059, 0.032	-0.60	0.551	0.940
lh_G_and_S_paracentral	0.003	-0.024, 0.031	0.25	0.801	0.940
lh_G_and_S_subcentral	0.001	-0.026, 0.029	0.10	0.918	0.940
lh_G_and_S_transv_frontopol	0.007	-0.047, 0.061	0.26	0.793	0.940
lh_G_and_S_cingul.Ant	0.011	-0.017, 0.039	0.78	0.439	0.916
lh_G_and_S_cingul.Mid.Ant	-0.002	-0.025, 0.021	-0.20	0.842	0.940
lh_G_and_S_cingul.Mid.Post	0.018	-0.004, 0.04	1.64	0.101	0.773
lh_G_cingul.Post.dorsal	0.028	0.001, 0.055	2.01	0.045	0.758
lh_G_cingul.Post.ventral	0.056	0, 0.111	1.98	0.048	0.758

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_cuneus	0.004	-0.032, 0.04	0.21	0.837	0.940
lh_G_front_inf.Opercular	0.000	-0.025, 0.025	0.01	0.990	0.990
lh_G_front_inf.Orbital	-0.007	-0.049, 0.034	-0.35	0.726	0.940
lh_G_front_inf.Triangul	0.003	-0.031, 0.038	0.20	0.845	0.940
lh_G_front_middle	0.016	-0.009, 0.042	1.25	0.212	0.780
lh_G_front_sup	0.013	-0.006, 0.033	1.32	0.186	0.780
lh_G_Ins_lg_and_S_cent_ins	0.011	-0.017, 0.038	0.75	0.455	0.926
lh_G_insular_short	0.003	-0.019, 0.025	0.24	0.813	0.940
lh_G_occipital_middle	-0.013	-0.043, 0.017	-0.84	0.402	0.916
lh_G_occipital_sup	0.004	-0.028, 0.036	0.26	0.794	0.940
lh_G_oc.temp_lat.fusifor	0.013	-0.023, 0.049	0.69	0.488	0.940
lh_G_oc.temp_med.Lingual	0.021	-0.021, 0.064	1.00	0.318	0.900
lh_G_oc.temp_med.Parahip	-0.003	-0.037, 0.031	-0.19	0.849	0.940
lh_G_orbital	0.016	-0.035, 0.066	0.62	0.539	0.940
lh_G_pariet_inf.Angular	0.016	-0.011, 0.044	1.17	0.242	0.809
lh_G_pariet_inf.Supramar	0.014	-0.008, 0.037	1.24	0.217	0.780
lh_G_parietal_sup	0.002	-0.024, 0.027	0.14	0.890	0.940
lh_G_postcentral	-0.003	-0.028, 0.021	-0.27	0.789	0.940
lh_G_precentral	0.016	-0.004, 0.036	1.57	0.117	0.773
lh_G_precuneus	0.011	-0.014, 0.035	0.87	0.387	0.916
lh_G_rectus	-0.050	-0.142, 0.043	-1.06	0.291	0.843
lh_G_subcallosal	0.030	-0.016, 0.075	1.28	0.200	0.780
lh_G_temp_sup.G_T_transv	0.028	-0.001, 0.056	1.91	0.057	0.758
lh_G_temp_sup.Lateral	0.005	-0.017, 0.028	0.47	0.642	0.940
lh_G_temp_sup.Plan_polar	-0.001	-0.024, 0.021	-0.11	0.911	0.940
lh_G_temp_sup.Plan_tempo	0.015	-0.012, 0.041	1.10	0.274	0.843
lh_G_temporal_inf	0.030	-0.018, 0.077	1.23	0.219	0.780
lh_G_temporal_middle	0.007	-0.022, 0.037	0.50	0.619	0.940
lh_Lat_Fis.ant.Horizontal	-0.010	-0.046, 0.026	-0.55	0.584	0.940
lh_Lat_Fis.ant.Vertical	0.038	0.002, 0.074	2.09	0.037	0.758
lh_Lat_Fis.post	0.011	-0.017, 0.039	0.79	0.428	0.916
lh_Pole_occipital	-0.021	-0.077, 0.036	-0.72	0.474	0.931
lh_Pole_temporal	-0.002	-0.041, 0.036	-0.11	0.913	0.940
lh_S_calcarine	0.026	-0.012, 0.063	1.36	0.176	0.780
lh_S_central	0.007	-0.02, 0.034	0.50	0.619	0.940
lh_S_cingul.Marginalis	0.006	-0.015, 0.028	0.58	0.563	0.940
lh_S_circular_insula_ant	-0.018	-0.05, 0.014	-1.11	0.268	0.843
lh_S_circular_insula_inf	-0.006	-0.03, 0.017	-0.52	0.605	0.940
lh_S_circular_insula_sup	-0.018	-0.042, 0.006	-1.46	0.146	0.773
lh_S_collat_transv_ant	0.033	-0.014, 0.08	1.37	0.171	0.780
lh_S_collat_transv_post	0.005	-0.033, 0.043	0.26	0.798	0.940
lh_S_front_inf	0.009	-0.021, 0.04	0.61	0.546	0.940
lh_S_front_middle	-0.001	-0.033, 0.03	-0.08	0.933	0.951

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_front_sup	0.006	-0.016, 0.027	0.49	0.623	0.940
lh_S_interm_prim.Jensen	0.027	-0.005, 0.059	1.65	0.100	0.773
lh_S_intrapariet_and_P_trans	0.014	-0.011, 0.038	1.08	0.283	0.843
lh_S_oc_middle_and_Lunatus	0.001	-0.04, 0.043	0.06	0.949	0.960
lh_S_oc_sup_and_transversal	0.006	-0.026, 0.038	0.35	0.726	0.940
lh_S_occipital_ant	0.010	-0.017, 0.037	0.72	0.474	0.931
lh_S_oc.temp_lat	0.010	-0.019, 0.039	0.66	0.508	0.940
lh_S_oc.temp_med_and_Lingual	0.023	-0.004, 0.05	1.69	0.091	0.773
lh_S_orbital_lateral	0.050	0.003, 0.097	2.11	0.036	0.758
lh_S_orbital_med.olfact	0.030	-0.051, 0.111	0.73	0.464	0.931
lh_S_orbital.H_Shaped	0.004	-0.034, 0.041	0.20	0.839	0.940
lh_S_parieto_occipital	0.021	-0.004, 0.047	1.63	0.103	0.773
lh_S_pericallosal	0.024	-0.004, 0.051	1.66	0.097	0.773
lh_S_postcentral	0.007	-0.015, 0.029	0.62	0.533	0.940
lh_S_precentral.inf.part	0.031	0.003, 0.06	2.16	0.032	0.758
lh_S_precentral.sup.part	0.016	-0.008, 0.041	1.32	0.188	0.780
lh_S_suborbital	0.027	-0.035, 0.09	0.86	0.393	0.916
lh_S_subparietal	0.030	0.002, 0.059	2.09	0.037	0.758
lh_S_temporal_inf	0.040	-0.002, 0.082	1.89	0.059	0.758
lh_S_temporal_sup	0.022	-0.003, 0.046	1.74	0.083	0.773
lh_S_temporal_transverse	0.050	0.016, 0.084	2.89	0.004	0.691
rh_G_and_S_frontomargin	-0.026	-0.083, 0.03	-0.92	0.360	0.916
rh_G_and_S_occipital_inf	0.017	-0.026, 0.061	0.78	0.439	0.916
rh_G_and_S_paracentral	-0.002	-0.03, 0.026	-0.12	0.905	0.940
rh_G_and_S_subcentral	0.015	-0.011, 0.041	1.11	0.269	0.843
rh_G_and_S_transv_frontopol	0.012	-0.034, 0.057	0.50	0.620	0.940
rh_G_and_S_cingul.Ant	-0.007	-0.037, 0.023	-0.48	0.632	0.940
rh_G_and_S_cingul.Mid.Ant	-0.005	-0.027, 0.018	-0.42	0.676	0.940
rh_G_and_S_cingul.Mid.Post	0.009	-0.013, 0.031	0.81	0.418	0.916
rh_G_cingul.Post.dorsal	0.030	0.003, 0.057	2.22	0.027	0.758
rh_G_cingul.Post.ventral	0.030	-0.014, 0.074	1.35	0.179	0.780
rh_G_cuneus	0.014	-0.022, 0.05	0.76	0.445	0.918
rh_G_front_inf.Opercular	0.005	-0.02, 0.031	0.42	0.677	0.940
rh_G_front_inf.Orbital	-0.002	-0.041, 0.037	-0.11	0.915	0.940
rh_G_front_inf.Triangul	0.008	-0.026, 0.042	0.44	0.659	0.940
rh_G_front_middle	0.005	-0.023, 0.034	0.37	0.708	0.940
rh_G_front_sup	-0.002	-0.022, 0.018	-0.19	0.848	0.940
rh_G_Ins_lg_and_S_cent_ins	0.026	-0.001, 0.052	1.89	0.059	0.758
rh_G_insular_short	0.003	-0.02, 0.026	0.23	0.818	0.940
rh_G_occipital_middle	0.018	-0.011, 0.047	1.21	0.227	0.788
rh_G_occipital_sup	0.008	-0.022, 0.039	0.54	0.587	0.940
rh_G_oc.temp_lat.fusifor	0.011	-0.028, 0.049	0.55	0.583	0.940
rh_G_oc.temp_med.Lingual	0.034	-0.008, 0.077	1.59	0.112	0.773

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_oc.temp_med.Parahip	-0.013	-0.049, 0.023	-0.69	0.493	0.940
rh_G_orbital	-0.003	-0.053, 0.047	-0.12	0.904	0.940
rh_G_pariet_inf.Angular	0.011	-0.016, 0.037	0.79	0.429	0.916
rh_G_pariet_inf.Supramar	0.007	-0.015, 0.029	0.63	0.526	0.940
rh_G_parietal_sup	0.006	-0.018, 0.03	0.50	0.620	0.940
rh_G_postcentral	-0.001	-0.025, 0.023	-0.12	0.908	0.940
rh_G_precentral	0.002	-0.016, 0.021	0.23	0.817	0.940
rh_G_precuneus	0.012	-0.013, 0.037	0.94	0.349	0.916
rh_G_rectus	-0.054	-0.178, 0.071	-0.85	0.395	0.916
rh_G_subcallosal	0.015	-0.03, 0.06	0.67	0.501	0.940
rh_G_temp_sup.G_T_transv	0.002	-0.028, 0.032	0.16	0.875	0.940
rh_G_temp_sup.Lateral	-0.005	-0.028, 0.018	-0.41	0.679	0.940
rh_G_temp_sup.Plan_polar	-0.008	-0.03, 0.015	-0.67	0.506	0.940
rh_G_temp_sup.Plan_tempo	0.034	0.009, 0.059	2.65	0.008	0.709
rh_G_temporal_inf	-0.007	-0.057, 0.043	-0.28	0.778	0.940
rh_G_temporal_middle	0.012	-0.019, 0.043	0.78	0.435	0.916
rh_Lat_Fis.ant.Horizontal	-0.026	-0.06, 0.007	-1.54	0.124	0.773
rh_Lat_Fis.ant.Vertical	-0.009	-0.048, 0.029	-0.48	0.631	0.940
rh_Lat_Fis.post	0.005	-0.023, 0.033	0.36	0.718	0.940
rh_Pole_occipital	0.024	-0.03, 0.077	0.88	0.382	0.916
rh_Pole_temporal	-0.028	-0.071, 0.015	-1.27	0.206	0.780
rh_S_calcarine	0.018	-0.02, 0.057	0.93	0.354	0.916
rh_S_central	-0.003	-0.03, 0.025	-0.18	0.855	0.940
rh_S_cingul.Marginalis	0.014	-0.009, 0.036	1.18	0.238	0.809
rh_S_circular_insula_ant	-0.013	-0.043, 0.017	-0.83	0.406	0.916
rh_S_circular_insula_inf	0.003	-0.021, 0.028	0.25	0.804	0.940
rh_S_circular_insula_sup	0.004	-0.02, 0.028	0.30	0.763	0.940
rh_S_collat_transv_ant	0.020	-0.03, 0.069	0.79	0.431	0.916
rh_S_collat_transv_post	0.016	-0.022, 0.055	0.82	0.410	0.916
rh_S_front_inf	0.006	-0.026, 0.038	0.37	0.712	0.940
rh_S_front_middle	-0.002	-0.031, 0.028	-0.11	0.911	0.940
rh_S_front_sup	0.007	-0.017, 0.031	0.59	0.557	0.940
rh_S_interm_prim.Jensen	-0.005	-0.032, 0.022	-0.39	0.699	0.940
rh_S_intrapariet_and_P_trans	0.002	-0.023, 0.026	0.13	0.898	0.940
rh_S_oc_middle_and_Lunatus	-0.007	-0.046, 0.032	-0.34	0.733	0.940
rh_S_oc_sup_and_transversal	0.016	-0.013, 0.045	1.07	0.283	0.843
rh_S_occipital_ant	0.024	-0.004, 0.051	1.70	0.090	0.773
rh_S_oc.temp_lat	0.002	-0.025, 0.03	0.18	0.860	0.940
rh_S_oc.temp_med_and_Lingual	0.020	-0.009, 0.048	1.37	0.171	0.780
rh_S_orbital_lateral	0.008	-0.034, 0.051	0.39	0.695	0.940
rh_S_orbital_med.olfact	0.021	-0.061, 0.103	0.50	0.617	0.940
rh_S_orbital.H_Shaped	0.008	-0.033, 0.048	0.38	0.704	0.940
rh_S_parieto_occipital	0.020	-0.006, 0.045	1.52	0.129	0.773

eTable 33. Differences in functional activity between the never-suicidal group and the caregiver-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_pericallosal	0.029	-0.005, 0.063	1.70	0.090	0.773
rh_S_postcentral	0.009	-0.013, 0.031	0.78	0.434	0.916
rh_S_precentral.inf.part	0.014	-0.016, 0.045	0.94	0.348	0.916
rh_S_precentral.sup.part	0.004	-0.019, 0.027	0.34	0.735	0.940
rh_S_suborbital	0.071	-0.02, 0.161	1.54	0.125	0.773
rh_S_subparietal	0.016	-0.013, 0.044	1.09	0.277	0.843
rh_S_temporal_inf	-0.029	-0.074, 0.017	-1.24	0.215	0.780
rh_S_temporal_sup	0.016	-0.008, 0.04	1.34	0.182	0.780
rh_S_temporal_transverse	0.008	-0.028, 0.043	0.42	0.671	0.940

Bold denotes group differences at p<0.05 uncorrected

eTable 34. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
Subcortical regions					
Left.Cerebellum.Cortex	-0.050	-0.122, 0.022	-1.38	0.170	0.754
Left.Thalamus.Proper	0.006	-0.026, 0.038	0.36	0.717	0.975
Left.Caudate	-0.036	-0.078, 0.006	-1.69	0.095	0.735
Left.Putamen	-0.005	-0.051, 0.041	-0.23	0.820	0.976
Left.Pallidum	0.009	-0.033, 0.052	0.43	0.670	0.975
Brain.Stem	-0.039	-0.082, 0.004	-1.80	0.074	0.730
Left.Hippocampus	-0.029	-0.065, 0.008	-1.56	0.122	0.735
Left.Amygdala	0.030	-0.018, 0.078	1.23	0.221	0.821
Left.Accumbens.area	0.009	-0.049, 0.067	0.30	0.766	0.975
Left.VentralDC	0.009	-0.035, 0.052	0.40	0.693	0.975
Right.Cerebellum.Cortex	-0.036	-0.105, 0.032	-1.05	0.295	0.898
Right.Thalamus.Proper	0.005	-0.028, 0.038	0.31	0.757	0.975
Right.Caudate	-0.029	-0.075, 0.016	-1.28	0.205	0.796
Right.Putamen	-0.039	-0.081, 0.004	-1.82	0.073	0.730
Right.Pallidum	-0.024	-0.058, 0.01	-1.40	0.165	0.754
Right.Hippocampus	0.015	-0.026, 0.055	0.71	0.477	0.960
Right.Amygdala	0.028	-0.02, 0.077	1.16	0.249	0.868
Right.Accumbens.area	-0.046	-0.089, -0.004	-2.17	0.032	0.492
Right.VentralDC	-0.009	-0.05, 0.032	-0.42	0.675	0.975
Cortical regions					
lh_G_and_S_frontomargin	-0.094	-0.169, -0.019	-2.48	0.015	0.492
lh_G_and_S_occipital_inf	-0.031	-0.115, 0.053	-0.74	0.462	0.960
lh_G_and_S_paracentral	-0.012	-0.053, 0.028	-0.60	0.548	0.975

eTable 34. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_G_and_S_subcentral	-0.020	-0.068, 0.029	-0.80	0.426	0.924
lh_G_and_S_transv_frontopol	-0.088	-0.179, 0.004	-1.90	0.060	0.669
lh_G_and_S_cingul.Ant	-0.004	-0.047, 0.04	-0.17	0.865	0.980
lh_G_and_S_cingul.Mid.Ant	-0.042	-0.081, -0.004	-2.20	0.030	0.492
lh_G_and_S_cingul.Mid.Post	0.001	-0.037, 0.038	0.05	0.962	0.992
lh_G_cingul.Post.dorsal	0.014	-0.028, 0.057	0.67	0.507	0.962
lh_G_cingul.Post.ventral	0.037	-0.055, 0.128	0.80	0.425	0.924
lh_G_cuneus	-0.041	-0.102, 0.019	-1.36	0.176	0.754
lh_G_front_inf.Opercular	-0.032	-0.072, 0.009	-1.55	0.124	0.735
lh_G_front_inf.Orbital	-0.066	-0.131, -0.001	-2.02	0.047	0.639
lh_G_front_inf.Triangul	-0.018	-0.078, 0.041	-0.60	0.548	0.975
lh_G_front_middle	-0.011	-0.053, 0.031	-0.53	0.600	0.975
lh_G_front_sup	0.003	-0.028, 0.035	0.21	0.835	0.976
lh_G_Ins_lg_and_S_cent_ins	0.010	-0.034, 0.054	0.44	0.658	0.975
lh_G_insular_short	-0.018	-0.054, 0.018	-1.00	0.318	0.898
lh_G_occipital_middle	-0.002	-0.057, 0.054	-0.05	0.956	0.992
lh_G_occipital_sup	-0.027	-0.084, 0.031	-0.93	0.357	0.898
lh_G_oc.temp_lat.fusifor	-0.042	-0.099, 0.015	-1.45	0.150	0.735
lh_G_oc.temp_med.Lingual	-0.036	-0.101, 0.028	-1.11	0.270	0.892
lh_G_oc.temp_med.Parahip	-0.036	-0.094, 0.022	-1.24	0.220	0.821
lh_G_orbital	-0.035	-0.109, 0.039	-0.93	0.353	0.898
lh_G_pariet_inf.Angular	0.000	-0.05, 0.05	0.01	0.992	0.992
lh_G_pariet_inf.Supramar	-0.013	-0.051, 0.025	-0.67	0.507	0.962
lh_G_parietal_sup	-0.025	-0.07, 0.02	-1.12	0.266	0.892
lh_G_postcentral	-0.010	-0.051, 0.032	-0.46	0.646	0.975
lh_G_precentral	0.000	-0.034, 0.034	-0.02	0.984	0.992
lh_G_precuneus	-0.001	-0.041, 0.038	-0.07	0.941	0.992
lh_G_rectus	-0.121	-0.273, 0.031	-1.57	0.119	0.735
lh_G_subcallosal	0.010	-0.066, 0.086	0.26	0.797	0.976
lh_G_temp_sup.G_T_transv	0.007	-0.047, 0.061	0.25	0.800	0.976
lh_G_temp_sup.Lateral	-0.001	-0.038, 0.037	-0.03	0.973	0.992
lh_G_temp_sup.Plan_polar	0.029	-0.007, 0.065	1.63	0.108	0.735
lh_G_temp_sup.Plan_tempo	-0.023	-0.066, 0.019	-1.08	0.284	0.898
lh_G_temporal_inf	0.004	-0.078, 0.087	0.10	0.922	0.992
lh_G_temporal_middle	-0.040	-0.093, 0.012	-1.53	0.129	0.735
lh_Lat_Fis.ant.Horizontal	-0.041	-0.099, 0.017	-1.41	0.161	0.754
lh_Lat_Fis.ant.Vertical	0.031	-0.03, 0.093	1.02	0.311	0.898
lh_Lat_Fis.post	0.005	-0.046, 0.056	0.20	0.841	0.976
lh_Pole_occipital	0.034	-0.059, 0.127	0.72	0.472	0.960
lh_Pole_temporal	-0.032	-0.1, 0.035	-0.94	0.347	0.898
lh_S_calcarine	0.024	-0.043, 0.09	0.70	0.483	0.960
lh_S_central	0.020	-0.029, 0.07	0.81	0.420	0.924
lh_S_cingul.Marginalis	-0.006	-0.04, 0.028	-0.36	0.718	0.975

eTable 34. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
lh_S_circular_insula_ant	-0.074	-0.128, -0.02	-2.71	0.008	0.449
lh_S_circular_insula_inf	-0.027	-0.069, 0.015	-1.28	0.203	0.796
lh_S_circular_insula_sup	-0.019	-0.061, 0.023	-0.89	0.378	0.898
lh_S_collat_transv_ant	-0.024	-0.106, 0.059	-0.57	0.569	0.975
lh_S_collat_transv_post	-0.001	-0.068, 0.066	-0.03	0.975	0.992
lh_S_front_inf	-0.019	-0.071, 0.033	-0.71	0.480	0.960
lh_S_front_middle	-0.007	-0.054, 0.041	-0.27	0.784	0.976
lh_S_front_sup	-0.001	-0.039, 0.037	-0.05	0.964	0.992
lh_S_interm_prim.Jensen	-0.003	-0.053, 0.048	-0.10	0.922	0.992
lh_S_intrapariet_and_P_trans	-0.012	-0.057, 0.033	-0.53	0.596	0.975
lh_S_oc_middle_and_Lunatus	-0.035	-0.107, 0.037	-0.97	0.335	0.898
lh_S_oc_sup_and_transversal	-0.028	-0.084, 0.029	-0.97	0.335	0.898
lh_S_occipital_ant	-0.006	-0.054, 0.041	-0.27	0.788	0.976
lh_S_oc.temp_lat	-0.076	-0.127, -0.025	-2.98	0.004	0.315
lh_S_oc.temp_med_and_Lingual	-0.004	-0.051, 0.044	-0.15	0.880	0.980
lh_S_orbital_lateral	0.001	-0.077, 0.079	0.02	0.982	0.992
lh_S_orbital_med.olfact	-0.070	-0.221, 0.081	-0.92	0.360	0.898
lh_S_orbital.H_Shaped	-0.046	-0.105, 0.014	-1.53	0.129	0.735
lh_S_parieto_occipital	0.008	-0.033, 0.05	0.40	0.690	0.975
lh_S_pericallosal	-0.015	-0.058, 0.028	-0.68	0.497	0.962
lh_S_postcentral	-0.022	-0.059, 0.014	-1.21	0.230	0.834
lh_S_precentral.inf.part	-0.005	-0.058, 0.049	-0.17	0.865	0.980
lh_S_precentral.sup.part	-0.009	-0.048, 0.031	-0.45	0.654	0.975
lh_S_suborbital	-0.029	-0.133, 0.074	-0.56	0.576	0.975
lh_S_subparietal	0.035	-0.012, 0.082	1.47	0.146	0.735
lh_S_temporal_inf	0.046	-0.033, 0.126	1.16	0.250	0.868
lh_S_temporal_sup	0.003	-0.04, 0.046	0.13	0.894	0.989
lh_S_temporal_transverse	0.011	-0.046, 0.069	0.39	0.696	0.975
rh_G_and_S_frontomargin	-0.100	-0.189, -0.01	-2.22	0.029	0.492
rh_G_and_S_occipital_inf	-0.014	-0.096, 0.068	-0.34	0.732	0.975
rh_G_and_S_paracentral	-0.017	-0.065, 0.032	-0.68	0.497	0.962
rh_G_and_S_subcentral	0.003	-0.038, 0.045	0.16	0.874	0.980
rh_G_and_S_transv_frontopol	-0.015	-0.084, 0.054	-0.44	0.662	0.975
rh_G_and_S_cingul.Ant	-0.023	-0.073, 0.027	-0.91	0.366	0.898
rh_G_and_S_cingul.Mid.Ant	-0.044	-0.083, -0.004	-2.18	0.032	0.492
rh_G_and_S_cingul.Mid.Post	-0.004	-0.042, 0.035	-0.19	0.848	0.976
rh_G_cingul.Post.dorsal	0.021	-0.023, 0.065	0.96	0.338	0.898
rh_G_cingul.Post.ventral	0.002	-0.068, 0.073	0.06	0.950	0.992
rh_G_cuneus	-0.008	-0.069, 0.054	-0.25	0.804	0.976
rh_G_front_inf.Opercular	-0.022	-0.063, 0.018	-1.10	0.272	0.892
rh_G_front_inf.Orbital	-0.058	-0.13, 0.013	-1.63	0.107	0.735
rh_G_front_inf.Triangul	-0.024	-0.08, 0.033	-0.83	0.406	0.924
rh_G_front_middle	-0.012	-0.056, 0.031	-0.56	0.574	0.975


eTable 34. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_G_front_sup	-0.007	-0.037, 0.023	-0.47	0.643	0.975
rh_G_Ins_lg_and_S_cent_ins	0.038	-0.013, 0.088	1.47	0.145	0.735
rh_G_insular_short	-0.029	-0.067, 0.009	-1.50	0.136	0.735
rh_G_occipital_middle	0.008	-0.045, 0.06	0.30	0.766	0.975
rh_G_occipital_sup	-0.007	-0.068, 0.054	-0.23	0.816	0.976
rh_G_oc.temp_lat.fusifor	-0.001	-0.062, 0.061	-0.02	0.984	0.992
rh_G_oc.temp_med.Lingual	0.001	-0.078, 0.079	0.01	0.988	0.992
rh_G_oc.temp_med.Parahip	-0.063	-0.127, 0.001	-1.97	0.052	0.639
rh_G_orbital	-0.101	-0.187, -0.015	-2.32	0.022	0.492
rh_G_pariet_inf.Angular	-0.016	-0.057, 0.025	-0.77	0.445	0.953
rh_G_pariet_inf.Supramar	-0.004	-0.039, 0.031	-0.21	0.837	0.976
rh_G_parietal_sup	-0.006	-0.047, 0.034	-0.31	0.756	0.975
rh_G_postcentral	0.004	-0.035, 0.043	0.20	0.842	0.976
rh_G_precentral	0.007	-0.027, 0.04	0.40	0.690	0.975
rh_G_precuneus	0.011	-0.035, 0.056	0.46	0.645	0.975
rh_G_rectus	-0.212	-0.428, 0.003	-1.96	0.054	0.639
rh_G_subcallosal	-0.032	-0.105, 0.041	-0.87	0.387	0.898
rh_G_temp_sup.G_T_transv	-0.006	-0.056, 0.044	-0.24	0.811	0.976
rh_G_temp_sup.Lateral	-0.017	-0.056, 0.022	-0.88	0.382	0.898
rh_G_temp_sup.Plan_polar	-0.010	-0.043, 0.022	-0.62	0.534	0.975
rh_G_temp_sup.Plan_tempo	0.014	-0.028, 0.056	0.66	0.513	0.963
rh_G_temporal_inf	-0.060	-0.146, 0.027	-1.37	0.173	0.754
rh_G_temporal_middle	0.012	-0.035, 0.06	0.52	0.605	0.975
rh_Lat_Fis.ant.Horizontal	-0.047	-0.105, 0.01	-1.64	0.104	0.735
rh_Lat_Fis.ant.Vertical	-0.052	-0.121, 0.017	-1.50	0.136	0.735
rh_Lat_Fis.post	0.021	-0.026, 0.069	0.89	0.376	0.898
rh_Pole_occipital	0.020	-0.084, 0.123	0.38	0.704	0.975
rh_Pole_temporal	-0.018	-0.096, 0.061	-0.45	0.656	0.975
rh_S_calcarine	0.016	-0.054, 0.087	0.46	0.645	0.975
rh_S_central	0.010	-0.042, 0.061	0.37	0.710	0.975
rh_S_cingul.Marginalis	0.006	-0.034, 0.045	0.29	0.770	0.975
rh_S_circular_insula_ant	-0.061	-0.111, -0.011	-2.44	0.017	0.492
rh_S_circular_insula_inf	0.003	-0.039, 0.046	0.16	0.875	0.980
rh_S_circular_insula_sup	-0.007	-0.05, 0.036	-0.33	0.746	0.975
rh_S_collat_transv_ant	-0.039	-0.124, 0.045	-0.93	0.357	0.898
rh_S_collat_transv_post	-0.032	-0.096, 0.033	-0.98	0.332	0.898
rh_S_front_inf	-0.033	-0.083, 0.017	-1.31	0.192	0.785
rh_S_front_middle	-0.047	-0.088, -0.006	-2.29	0.024	0.492
rh_S_front_sup	-0.014	-0.052, 0.023	-0.76	0.452	0.956
rh_S_interm_prim.Jensen	-0.070	-0.116, -0.025	-3.08	0.003	0.315
rh_S_intrapariet_and_P_trans	-0.010	-0.055, 0.035	-0.44	0.658	0.975
rh_S_oc_middle_and_Lunatus	-0.045	-0.113, 0.023	-1.31	0.193	0.785
rh_S_oc_sup_and_transversal	0.025	-0.029, 0.079	0.93	0.356	0.898


eTable 34. Differences in functional activity between the never-suicidal group and the concordantly-reported suicidality group for emotional faces: Welch's t-test

ROIs	Mean	95%CI	t	p (unc)	p FDR
rh_S_occipital_ant	0.019	-0.023, 0.061	0.89	0.374	0.898
rh_S_oc.temp_lat	-0.014	-0.061, 0.033	-0.60	0.552	0.975
rh_S_oc.temp_med_and_Lingual	0.019	-0.028, 0.067	0.81	0.418	0.924
rh_S_orbital_lateral	-0.003	-0.069, 0.064	-0.08	0.938	0.992
rh_S_orbital_med.olfact	-0.033	-0.19, 0.123	-0.42	0.674	0.975
rh_S_orbital.H_Shaped	-0.055	-0.121, 0.011	-1.66	0.101	0.735
rh_S_parieto_occipital	0.033	-0.01, 0.077	1.53	0.129	0.735
rh_S_pericallosal	-0.002	-0.056, 0.052	-0.08	0.935	0.992
rh_S_postcentral	-0.008	-0.042, 0.027	-0.45	0.655	0.975
rh_S_precentral.inf.part	-0.008	-0.059, 0.043	-0.30	0.767	0.975
rh_S_precentral.sup.part	-0.023	-0.066, 0.02	-1.06	0.292	0.898
rh_S_suborbital	0.044	-0.122, 0.21	0.53	0.596	0.975
rh_S_subparietal	0.015	-0.036, 0.067	0.59	0.554	0.975
rh_S_temporal_inf	-0.056	-0.134, 0.021	-1.45	0.150	0.735
rh_S_temporal_sup	0.006	-0.031, 0.043	0.34	0.738	0.975
rh_S_temporal_transverse	-0.012	-0.067, 0.043	-0.44	0.664	0.975


Bold denotes group differences at $p < 0.05$ uncorrected




eFigure 54. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 55 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 55 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 55 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the child-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 56. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 57 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 57 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 57 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the caregiver-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 58. Results of the Equivalence test for differences in functional activity in subcortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).




eFigure 59 (1 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 59 (2 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 59 (3 of 3). Results of the Equivalence test for differences in functional activity in cortical regions between the never-suicidal group and the concordantly-reported suicidality group for emotional faces. The figure shows the effect sizes and 95% confidence intervals of the difference against the interval of equivalence (-0.15 to 0.15).



eFigure 60. Equivalence testing for mean differences in functional activation during the MID task, the SST and EN-back task relating to suicidality. Distribution of absolute t-values from Equivalence tests comparing the regional means between the never-suicidal group and the child-reported suicidality group (Panel A) and the caregiver-reported suicidality group (Panel B); Higher t-values (i.e., darker green) suggest equivalence between groups. No statistical differences were found after applying FDR-correction for multiple comparisons.

eTable 35. Range of effect sizes (Cohen's d) and 90% confidence intervals bounds for each imaging modality and suicidality group comparison

Contrast		Child		Caregiver		Concordant	
		Min	Max	Min	Max	Min	Max
sMRI							
-	ES range	-0.11	0.12	-0.17	0.11	-0.23	0.23
	90%CI range	-0.20	0.20	-0.26	0.19	-0.39	0.39
rs-fMRI							
-	ES range	-0.17	0.14	-0.18	0.20	-0.34	0.28
	90%CI range	-0.29	0.25	-0.30	0.32	-0.55	-0.49
Task-based fMRI							
MID							
<i>Reward anticipation</i>	ES range	-0.17	0.08	-0.12	0.12	-0.34	0.20
	90%CI range	-0.28	0.18	-0.23	0.23	-0.54	0.40
<i>Loss anticipation</i>	ES range	-0.13	0.12	-0.15	0.09	-0.30	0.22
	90%CI range	-0.23	0.23	-0.26	0.21	-0.51	0.42
<i>Reward feedback</i>	ES range	-0.13	0.07	-0.17	0.10	-0.33	0.22
	90%CI range	-0.23	0.17	-0.28	0.21	-0.54	0.42
<i>Loss feedback</i>	ES range	-0.13	0.10	-0.08	0.20	-0.19	0.22
	90%CI range	-0.23	0.21	-0.19	0.31	-0.39	0.42
SST							
<i>Incorrect stop vs correct go</i>	ES range	-0.15	0.12	-0.15	0.12	-0.20	0.21
	90%CI range	-0.26	0.23	-0.26	0.21	-0.41	0.42
<i>Correct stop vs correct go</i>	ES range	-0.08	0.16	-0.08	0.16	-0.24	0.25
	90%CI range	-0.19	0.26	-0.25	0.26	-0.45	0.47
ENBack							
<i>2-back vs 0-back</i>	ES range	-0.15	0.06	-0.15	0.12	-0.32	0.17
	90%CI range	-0.26	0.17	-0.27	0.23	-0.53	0.38
<i>Emotional vs neutral faces</i>	ES range	-0.17	0.09	-0.09	0.17	-0.33	0.17
	90%CI range	-0.28	0.21	-0.21	0.29	-0.55	0.38

CI, confidence interval. ES, effect size. Min, minimum. Max, maximum. MID, Monetary incentive delay task. SST, Stop signal task. EN-back, Emotional n-back task.

eTable 36. Results, 95% confidence intervals and area under precision-recall curve with effect sizes (Cohen’s *d*) largest than |0.15| for each imaging modality in child- and caregiver-suicidality group comparison

	Child				Caregiver			
	Parcellation	ES	95%CI	PRC	Parcellation	ES	95%CI	PRC
sMRI								
	None	-	-		L bank superior temporal sulcus ^a	-0.17	-0.26, -0.08	0.07
rs-fMRI								
	Cingulo-parietal – Fronto-parietal	-0.17	-0.29, -0.06	0.07	Default - R hippocampus	-0.18	-0.3, -0.06	0.07
					Default – L amygdala	-0.17	-0.29, -0.05	0.07
					Dorsal attention – L cerebellum	0.15	0.03, 0.27	0.10
					Auditory – R ventral dc	0.15	0.03, 0.27	0.09
					Sensorimotor-mouth – L accumbens area	0.16	0.04, 0.28	0.09
					Cingulo-parietal -R accumbens area	0.18	0.06, 0.3	0.09
					Cingulo-parietal - L hippocampus	0.20	0.08, 0.33	0.09
Task-based fMRI								
MID								
<i>Reward anticipation</i>	R Marginal branch of the cingulate sulcus	-0.17	-0.28, -0.07	0.07	None	-	-	
	R parieto-occipital sulcus	-0.16	-0.27, -0.05	0.08				
	L calcarine sulcus	-0.16	-0.27, -0.05	0.07				
	R supramarginal gyrus	-0.16	-0.27, -0.05	0.08				
	L Marginal branch of the cingulate sulcus	-0.15	-0.26, -0.04	0.08				
<i>Loss anticipation</i>	None	-	-		L fronto-marginal gyrus and sulcus	-0.15	-0.27, -0.04	0.09
<i>Reward feedback</i>	None	-	-		R posterior segment of the lateral sulcus	-0.17	-0.28, -0.06	0.07
<i>Loss feedback</i>	None	-	-		R anterior segment of lateral sulcus that splits into vertical	0.18	0.07, 0.3	0.10
					L middle temporal gyrus	0.20	0.08, 0.31	0.10

eTable 36. Results, 95% confidence intervals and area under precision-recall curve with effect sizes (Cohen’s *d*) largest than |0.15| for each imaging modality in child- and caregiver-suicidality group comparison

	Child				Caregiver			
	Parcellation	ES	95%CI	PRC	Parcellation	ES	95%CI	PRC
SST								
<i>Incorrect stop vs correct go</i>	L superior occipital and transverse occipital sulci	-0.15	-0.27, -0.04	0.07	None	-	-	-
<i>Correct stop vs correct go</i>	L Superior segment of the circular sulcus of the insula	0.16	0.05, 0.28	0.10	None	-	-	-
ENBack								
<i>2-back vs 0-back</i>	None	-	-	-	None	-	-	-
<i>Emotional vs neutral faces</i>	L lateral occipito-temporal sulcus	-0.17	-0.29, -0.05	0.07	R planum temporale- superior temporal gyrus	0.16	0.04, 0.28	0.09
					L temporal transverse sulcus	0.17	0.05, 0.29	0.09

PRC, area under precision-recall curve. ES, effect size. CI, confidence interval. L, left. R, right.

^a Unique result that was statistically different ($p_{FDR} < 0.05$) after FDR-correction for multiple comparisons.

eReferences

1. Garavan H, Bartsch H, Conway K, et al. Recruiting the ABCD sample: Design considerations and procedures. *Developmental cognitive neuroscience*. 2018;32:16-22.
2. Kobak K, Kratochvil C, Stanger C, Kaufman J. Computerized screening of comorbidity in adolescents with substance or psychiatric disorders. Paper presented at: Poster presented at the 33rd annual anxiety disorders and depression conference, La Jolla, CA. Retrieved from http://www.telepsychology.net/resources/KSADS_ADAAPosterv4.pdf2013.
3. Luciana M, Bjork JM, Nagel BJ, et al. Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Developmental cognitive neuroscience*. 2018;32:67-79.
4. Barch DM, Albaugh MD, Avenevoli S, et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Developmental cognitive neuroscience*. 2018;32:55-66.
5. Zucker RA, Gonzalez R, Feldstein Ewing SW, et al. Assessment of culture and environment in the Adolescent Brain and Cognitive Development Study: Rationale, description of measures, and early data. *Developmental cognitive neuroscience*. 2018;32:107-120.
6. Knutson B, Westdorp A, Kaiser E, Hommer D. fMRI visualization of brain activity during a monetary incentive delay task. *NeuroImage*. 2000;12(1):20-27.
7. Logan GD. On the ability to inhibit thought and action: A users' guide to the stop signal paradigm. In: *Inhibitory processes in attention, memory, and language*. San Diego, CA, US: Academic Press; 1994:189-239.
8. Cohen A, Conley M, Dellarco D, Casey BJPotSfNSD, CA. November. The impact of emotional cues on short-term and long-term memory during adolescence. 2016.
9. Casey BJ, Cannonier T, Conley MI, et al. The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Developmental cognitive neuroscience*. 2018;32:43-54.
10. Hagler DJ, Jr., Hatton S, Cornejo MD, et al. Image processing and analysis methods for the Adolescent Brain Cognitive Development Study. *NeuroImage*. 2019:116091.
11. Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex*. 2016;26(1):288-303.
12. Van Dijk KR, Hedden T, Venkataraman A, Evans KC, Lazar SW, Buckner RL. Intrinsic functional connectivity as a tool for human connectomics: theory, properties, and optimization. *Journal of neurophysiology*. 2010;103(1):297-321.
13. Seibert TM, Brewer JB. Default network correlations analyzed on native surfaces. *Journal of neuroscience methods*. 2011;198(2):301-311.

14. Cohen AO, Breiner K, Steinberg L, et al. When Is an Adolescent an Adult? Assessing Cognitive Control in Emotional and Nonemotional Contexts. 2016;27(4):549-562.
15. Jollant F, Lawrence NS, Giampietro V, et al. Orbitofrontal cortex response to angry faces in men with histories of suicide attempts. *The American journal of psychiatry*. 2008;165(6):740-748.
16. Jollant F, Lawrence NS, Olie E, et al. Decreased activation of lateral orbitofrontal cortex during risky choices under uncertainty is associated with disadvantageous decision-making and suicidal behavior. *NeuroImage*. 2010;51(3):1275-1281.
17. Pan L, Segreti A, Almeida J, et al. Preserved hippocampal function during learning in the context of risk in adolescent suicide attempt. *Psychiatry research*. 2013;211(2):112-118.
18. Pan LA, Batezati-Alves SC, Almeida JR, et al. Dissociable patterns of neural activity during response inhibition in depressed adolescents with and without suicidal behavior. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2011;50(6):602-611.e603.
19. Pan LA, Hassel S, Segreti AM, Nau SA, Brent DA, Phillips ML. Differential patterns of activity and functional connectivity in emotion processing neural circuitry to angry and happy faces in adolescents with and without suicide attempt. *Psychological medicine*. 2013;43(10):2129-2142.
20. Vanyukov PM, Szanto K, Hallquist MN, et al. Paralimbic and lateral prefrontal encoding of reward value during intertemporal choice in attempted suicide. *Psychological medicine*. 2016;46(2):381-391.
21. Xie W, Li H, Luo X, et al. Anhedonia and pain avoidance in the suicidal mind: behavioral evidence for motivational manifestations of suicidal ideation in patients with major depressive disorder. *Journal of clinical psychology*. 2014;70(7):681-692.
22. McHugh CM, Chun Lee RS, Hermens DF, Corderoy A, Large M, Hickie IB. Impulsivity in the self-harm and suicidal behavior of young people: A systematic review and meta-analysis. *Journal of psychiatric research*. 2019;116:51-60.
23. Bauer BW, Gustafsson HC, Nigg J, Karalunas SL. Working memory mediates increased negative affect and suicidal ideation in childhood attention-deficit/hyperactivity disorder. *J Psychopathol Behav Assess*. 2018;40(2):180-193.
24. Bomyea J, Stout DM, Simmons AN. Attenuated prefrontal and temporal neural activity during working memory as a potential biomarker of suicidal ideation in veterans with PTSD. *Journal of affective disorders*. 2019;257:607-614.
25. Seymour KE, Jones RN, Cushman GK, et al. Emotional face recognition in adolescent suicide attempters and adolescents engaging in non-suicidal self-injury. *European child & adolescent psychiatry*. 2016;25(3):247-259.

26. Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and biomedical research, an international journal*. 1996;29(3):162-173.
27. Destrieux C, Fischl B, Dale A, Halgren E. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *NeuroImage*. 2010;53(1):1-15.
28. Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002;33(3):341-355.
29. Pornpattananangkul N, Leibenluft E, Pine DS, Stringaris A. Mapping Anhedonia in Youth: Large-Scale Resting-State Network, Task-Evoked Activation and Phenotypic Demarcation. *JAMA psychiatry*. 2019.
30. Fortin JP, Parker D, Tunc B, et al. Harmonization of multi-site diffusion tensor imaging data. *NeuroImage*. 2017;161:149-170.
31. Nielson DM, Pereira F, Zheng CY, et al. Detecting and harmonizing scanner differences in the ABCD study - annual release 1.0. In:2018.
32. Dick AS, Garcia NL, Pruden SM, et al. No evidence for a bilingual executive function advantage in the nationally representative ABCD study. *Nature human behaviour*. 2019;3(7):692-701.
33. Ialongo C. The logic of equivalence testing and its use in laboratory medicine. *Biochimica medica*. 2017;27(1):5-13.
34. Lakens D. Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. *Social psychological and personality science*. 2017;8(4):355-362.
35. Schuirmann DJ. A comparison of the two one-sided tests procedure and the power approach for assessing the equivalence of average bioavailability. *Journal of pharmacokinetics and biopharmaceutics*. 1987;15(6):657-680.
36. Rentería ME, Schmaal L, Hibar DP, et al. Subcortical brain structure and suicidal behaviour in major depressive disorder: a meta-analysis from the ENIGMA-MDD working group. *Translational psychiatry*. 2017;7(5):e1116.
37. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, NJ: Erlbaum; 1988.
38. Lakens D, McLatchie N, Isager PM, Scheel AM, Dienes Z. Improving Inferences about Null Effects with Bayes Factors and Equivalence Tests. *The journals of gerontology Series B, Psychological sciences and social sciences*. 2018.
39. Saito T, Rehmsmeier M. The precision-recall plot is more informative than the ROC plot when evaluating binary classifiers on imbalanced datasets. *PloS one*. 2015;10(3):e0118432.