TABLE 2
Normalized mean square errors (NMSE) in the normal cognition (NC) and mild cognitive impairment (MCI) groups.

Region	NMSE (SD)	
	NC	MCI
Whole Brain	0.68 (0.098)	0.7 (0.07)
Frontal	0.69 (0.092)	0.71(0.067)
Parietal	0.65 (0.11)	0.68 (0.079)
Occipital	0.64 (0.12)	0.67 (0.096)
Temporal	0.68 (0.10)	0.71 (0.078)



Figure 9. Average (95%-CI) whole-brain CA (top), CVR (middle), and PRR (bottom) and PDM gains in NC (blue) and MCI (red). Asterisk: $p < 0.05$.

TABLE 3
Average (SD) PDM derived CVR (left panel) compared to CVR during hypercapnic challenge (right panel) between NC and MCI in the whole brain and the frontal, parietal, occipital, and temporal lobes.
*: $p < 0.05$.

	PDM derived CVR		CVR during Hypercapnic challenge	
Region	NC	MCI	NC	MCI
Whole Brain	0.49(0.96)	0.21(0.85)	0.16(0.13)	0.17(0.1)
Frontal	0.24(1)	0.12(0.88)	0.13(0.13)	0.15(0.099)
Parietal	0.83(2)	0.32(1.8)	0.21(0.2)	0.26(0.15)
Occipital	1.1(2.1)	0.26(1.6)*	0.21(0.16)	0.22(0.16)
Temporal	0.58(1.2)	0.18(0.9)*	0.16(0.12)	0.17(0.1)



Figure 12. Association of whole-brain PDM-based CVR (left) and the 4th PDM gain of the PR input (right) using resting state BOLD with MOCA (top), delayed recall HVLT (middle), and Trail Making test B-A (model). β = slope.