

Supplementary Table 5. Primer sequences used for the RT-PCR of the in vitro model showing the effect of metformin on fibroblasts from women with POP (**Figure 2**).

Target gene		Oligonucleotide sequence	Annealing temperature (°C)	Product Size (bp)
COL1A1	Forward	5' TCCAACGAGATCGAGATCC 3'	56.05	191
	Reverse	5' AAGCCGAATTCCTGGTCT 3'	55.76	
COL3A1	Forward	5' GATCCGTTCTCTGCGATGAC 3'	58.52	279
	Reverse	5' AGTTCTGAGGACCAGTAGGG 3'	57.76	
ELN	Forward	5' TTTGGCCCGGGAGTAGTTGG 3'	62.13	136
	Reverse	5' CAGCTGCTTCTGGTGACACAAC 3'	62.2	
YWHAZ	Forward	5' GATGAAGCCATTGCTGAACCTG 3'	58.49	229
	Reverse	5' CTATTTGTGGGACAGCATGGA 3'	58.00	
HPRT1	Forward	5' GCTGACCTGCTGGATTACAT 3'	57.66	260
	Reverse	5' CTTGCGACCTTGACCATCT 3'	57.46	

RT-PCR, reverse-transcription polymerase chain reaction; COL1A1, α 1(I)procollagen; COL3A1, α 1(III)procollagen; ELN, elastin; YWHAZ, tyrosine 3-monooxygenase /tryptophan 5-monooxygenase activation protein, zeta polypeptide; HPRT1, hypoxanthine phosphoribosyltransferase 1.