



Supplementary Figure 1: Conversion distributions for repeat-tested patients with initial SARS-CoV-2 negative (**a**, **c**, **e**, and **g**) and positive (**b**, **d**, **f**, and **h**) results. For the analyses in **a** to **d** only the first conversion events (i.e. negative to positive - **a** and **c** - or positive to negative - **b** and **d**) were considered. N represents the total numbers of patients analyzed. The distributions in **a** to **d** are represented by density-adjusted histograms (in grey bars), kernel probability density lines (in blue), unadjusted cumulative distributions (red line) and Kaplan-Meier hazard rates (green); the timing of individual results are represented by the black marks along the x-axis. For the Kaplan Meier estimated hazard rates (**e** to **h**), in addition to the conversion events, patients were considered censored at the time of the last unchanged result were included and the number of patients at risk for conversion at each time point are shown in **g** and **h**.


a Conversion to Negative
In Patients Initially Positive




b Conversion to Positive
In Patients Initially Negative




c




d




e




f




g




h



Supplementary Figure 2: SARS-CoV-2 results from repeat-tested patients that converted from negative to positive and back to negative (**A**) or from positive to negative and back to positive (**B**). SARS-CoV-2 negative results are represented by blue dots and positive results are in red, plotted on a time scale from the date of the first test





Supplementary Tables

Supplementary Table 1: Numbers of unique patients in the Columbia Clinical COVID-19 dataset, NYP Laboratory SARS-CoV-2 Testing dataset, and the merged dataset.

Unique Patients	Merged dataset	Laboratory dataset only	Clinical dataset only
Total	1588	20750	36
SARS-CoV-2 Positive	990	9653	11
Repeat-tested	275	3157	0
With symptoms	529	0	8
Deceased	84	0	0
SARS-CoV-2 Tests			
Total	2023	25354	2
SARS-CoV-2 Positive	1266	11454	4
Repeat-tested	710	7761	0
With symptoms	707	0	1
Deceased	115	0	0

Supplementary Table 2: Patient characteristics, grouped by highest SARS-2-CoV2 result in each patient. **Negative** indicates a result of “Not Detected” and **Positive** includes results reported as “Detected” or “Indeterminate”. Time-dependent clinical factors, such as age group, symptom start to test interval, and encounter type at time of test order, are counted at the time of the first test.

	Negative (N=11352)	Positive (N=10920)	Total (N=22272)	p value
Gender at birth				< 0.001 ¹
Female	7045 (62.1%)	5061 (46.4%)	12106 (54.4%)	
Male	4291 (37.8%)	5851 (53.6%)	10142 (45.5%)	
Unknown	16 (0.1%)	7 (0.1%)	23 (0.1%)	
N	11352	10919	22271	
Age at order				< 0.001 ²
Median	42.9	61.5	54.4	
Mean (sd)	46.5 (23.0)	60.0 (19.3)	53.1 (22.3)	
Q1, Q3	31.4, 64.0	46.5, 74.5	35.5, 70.4	
Range	0.0 - 120.3	0.0 - 144.7	0.0 - 144.7	
N	11352	10920	22272	
Age group (y)				< 0.001 ³
0-19	1069 (9.4%)	155 (1.4%)	1224 (5.5%)	
20-44	4912 (43.3%)	2387 (21.9%)	7299 (32.8%)	
45-54	1255 (11.1%)	1533 (14.0%)	2788 (12.5%)	
55-64	1389 (12.2%)	2148 (19.7%)	3537 (15.9%)	
65-74	1236 (10.9%)	2065 (18.9%)	3301 (14.8%)	
75-84	868 (7.7%)	1588 (14.5%)	2456 (11.0%)	
>=85	616 (5.4%)	1044 (9.6%)	1660 (7.5%)	
N	11345	10920	22265	
Ethnicity				<(0.001 ¹)
Hispanic/Latino	154 (26.3%)	308 (31.0%)	462 (29.3%)	
Not Hispanic/Latino	311 (53.2%)	409 (41.2%)	720 (45.7%)	
Not Specified	120 (20.5%)	275 (27.7%)	395 (25.0%)	
N	585	992	1577	
Race				< 0.001 ¹
Asian	752 (6.6%)	531 (4.9%)	1283 (5.8%)	
Black or African American	1629 (14.3%)	1833 (16.8%)	3462 (15.5%)	
Caucasian	3586 (31.6%)	2356 (21.6%)	5942 (26.7%)	

	Negative (N=11352)	Positive (N=10920)	Total (N=22272)	p value
Declined	1593 (14.0%)	1308 (12.0%)	2901 (13.0%)	
Other	1769 (15.6%)	2287 (20.9%)	4056 (18.2%)	
Pacific Islander	20 (0.2%)	25 (0.2%)	45 (0.2%)	
Unavailable	2003 (17.6%)	2580 (23.6%)	4583 (20.6%)	
N	11352	10920	22272	
Highest level of care				0.025 ¹
Admitted	229 (41.4%)	305 (35.1%)	534 (37.6%)	
Discharged from ED	58 (10.5%)	71 (8.2%)	129 (9.1%)	
ICU-level Care	69 (12.5%)	119 (13.7%)	188 (13.2%)	
Outpatient	197 (35.6%)	373 (43.0%)	570 (40.1%)	
N	553	868	1421	
Encounter type at order				< 0.001 ¹
Emergency	2706 (23.8%)	4748 (43.5%)	7454 (33.5%)	
Inpatient	3827 (33.7%)	2555 (23.4%)	6382 (28.7%)	
Outpatient	4819 (42.5%)	3617 (33.1%)	8436 (37.9%)	
N	11352	10920	22272	
Location at order				< 0.001 ¹
Allen Hospital	448 (3.9%)	845 (7.7%)	1293 (5.8%)	
Columbia	4505 (39.7%)	3924 (35.9%)	8429 (37.8%)	
Columbia Outreach	509 (4.5%)	743 (6.8%)	1252 (5.6%)	
Lower Manhattan	633 (5.6%)	437 (4.0%)	1070 (4.8%)	
Queens	1588 (14.0%)	2555 (23.4%)	4143 (18.6%)	
Weill Cornell	3518 (31.0%)	2388 (21.9%)	5906 (26.5%)	
Westchester	151 (1.3%)	28 (0.3%)	179 (0.8%)	
N	11352	10920	22272	
Number of admissions				0.008 ²
Median	1.0	0.0	0.0	
Mean (sd)	0.5 (0.5)	0.5 (0.5)	0.5 (0.5)	
Q1, Q3	0.0, 1.0	0.0, 1.0	0.0, 1.0	
Range	0.0 - 2.0	0.0 - 2.0	0.0 - 2.0	
N	584	995	1579	
Admission duration (d)				0.046 ²

	Negative (N=11352)	Positive (N=10920)	Total (N=22272)	p value
Median	1.0	0.6	0.7	
Mean (sd)	5.7 (13.9)	4.5 (8.1)	4.9 (10.6)	
Q1, Q3	0.4, 4.0	0.3, 4.9	0.3, 4.5	
Range	0.0 - 139.0	0.0 - 82.2	0.0 - 139.0	
N	584	995	1579	
Admission 1 to test 1 (d)				< 0.001 ²
Median	0.4	0.2	0.3	
Mean (sd)	3.0 (11.4)	0.7 (2.6)	1.6 (7.4)	
Q1, Q3	0.2, 1.1	0.1, 0.6	0.2, 0.9	
Range	-13.9 - 133.1	0.0 - 55.4	-13.9 - 133.1	
N	555	924	1479	
Admission 2 to test 1 (d)				0.999 ²
Median	-7.9	-8.5	-8.3	
Mean (sd)	-9.4 (9.3)	-9.4 (7.5)	-9.4 (8.1)	
Q1, Q3	-16.9, -0.9	-14.3, -2.1	-15.4, -1.5	
Range	-29.0 - 1.2	-25.9 - 0.3	-29.0 - 1.2	
N	34	70	104	
Primary Outcome				< 0.001 ¹
Deceased/Discharged to Hospice	29 (5.0%)	72 (7.2%)	101 (6.4%)	
Discharged (not death)	542 (92.8%)	865 (86.9%)	1407 (89.1%)	
Intubated (still admitted)	1 (0.2%)	44 (4.4%)	45 (2.8%)	
Still admitted (not intubated)	12 (2.1%)	14 (1.4%)	26 (1.6%)	
N	584	995	1579	
Deceased				0.048 ¹
No	562 (96.2%)	933 (93.8%)	1495 (94.7%)	
Yes	22 (3.8%)	62 (6.2%)	84 (5.3%)	
N	584	995	1579	
Symptomatic				0.004 ¹
No	2 (6.9%)	10 (1.8%)	12 (2.0%)	
Unknown	6 (20.7%)	36 (6.4%)	42 (7.1%)	
Yes	21 (72.4%)	514 (91.8%)	535 (90.8%)	
N	29	560	589	
Symptoms start to death (d)				0.828 ²
Median	11.6	14.0	13.7	
Mean (sd)	11.6 (NA)	14.8 (9.3)	14.7 (9.2)	
Q1, Q3	11.6, 11.6	6.9, 19.2	7.0, 19.1	
Range	11.6 - 11.6	1.6 - 39.6	1.6 - 39.6	
N	1	46	47	

	Negative (N=11352)	Positive (N=10920)	Total (N=22272)	p value
Any fever				0.936 ¹
No	5 (23.8%)	133 (25.4%)	138 (25.4%)	
Yes	16 (76.2%)	390 (74.6%)	406 (74.6%)	
N	21	523	544	
Intubated				0.026 ¹
No	543 (93.0%)	889 (89.3%)	1432 (90.7%)	
Yes	41 (7.0%)	106 (10.7%)	147 (9.3%)	
N	584	995	1579	
Extubated				0.046 ¹
No	557 (95.4%)	969 (97.4%)	1526 (96.6%)	
Yes	27 (4.6%)	26 (2.6%)	53 (3.4%)	
N	584	995	1579	
Symptoms start to intubation (d)				0.152 ²
Median	7.0	8.0	8.0	
Mean (sd)	16.3 (19.7)	9.2 (7.3)	9.4 (7.9)	
Q1, Q3	5.0, 23.0	4.2, 11.0	4.0, 11.0	
Range	3.0 - 39.0	-3.0 - 39.0	-3.0 - 39.0	
N	3	90	93	
Intubation duration (d)				0.069 ²
Median	3.0	6.0	5.0	
Mean (sd)	4.2 (4.7)	6.9 (4.7)	5.4 (4.9)	
Q1, Q3	1.2, 5.0	4.5, 9.5	2.0, 8.0	
Range	-3.0 - 20.0	0.0 - 20.0	-3.0 - 20.0	
N	26	23	49	
Decompensated				0.005 ¹
No	535 (91.6%)	862 (86.6%)	1397 (88.5%)	
Yes	49 (8.4%)	133 (13.4%)	182 (11.5%)	
N	584	995	1579	
Symptoms start to decompensation (d)				
Median	9.0	8.0	8.0	
Mean (sd)	15.0 (16.3)	9.9 (8.2)	10.1 (8.5)	
Q1, Q3	6.0, 18.0	4.0, 13.0	4.0, 13.0	
Range	3.0 - 39.0	-3.0 - 39.0	-3.0 - 39.0	
N	4	108	112	
Symptoms start to first test (d)				
Median	4.4	4.7	4.7	
Mean (sd)	6.9 (13.7)	6.9 (17.0)	6.9 (16.9)	
Q1, Q3	2.5, 12.5	2.7, 7.9	2.7, 7.9	

	Negative (N=11352)	Positive (N=10920)	Total (N=22272)	p value
Range	-29.4 - 51.1	-4.3 - 366.8	-29.4 - 366.8	
N	21	508	529	
Number of tests				< 0.001 ²
Median	1.0	1.0	1.0	
Mean (sd)	1.2 (0.6)	1.2 (0.7)	1.2 (0.7)	
Q1, Q3	1.0, 1.0	1.0, 1.0	1.0, 1.0	
Range	1.0 - 8.0	1.0 - 41.0	1.0 - 41.0	
N	11352	10920	22272	
Number of positive results per patient				
Median	0.0	1.0	0.0	
Mean (sd)	0.0 (0.0)	1.0 (0.4)	0.5 (0.6)	
Q1, Q3	0.0, 0.0	1.0, 1.0	0.0, 1.0	
Range	0.0 - 0.0	0.0 - 20.0	0.0 - 20.0	
N	11352	10920	22272	
Number of positive results per patient (Indeterminate included)				
Median	0.0	1.0	0.0	
Mean (sd)	0.0 (0.0)	1.1 (0.4)	0.5 (0.6)	
Q1, Q3	0.0, 0.0	1.0, 1.0	0.0, 1.0	
Range	0.0 - 0.0	1.0 - 20.0	0.0 - 20.0	
N	11352	10920	22272	

1. Pearson's Chi-squared test (adjusted for multiple comparisons)
2. Linear Model ANOVA (adjusted for multiple comparisons)
3. Trend test for ordinal variables (adjusted for multiple comparisons)

Supplementary Table 3: Patient characteristics comparing repeat-tested with single-tested patients.

	Repeat (N=3432)	Single (N=18906)	Total (N=22338)	p value
Gender at birth				<0.001 ¹
Female	1644 (47.9%)	10503 (55.6%)	12147 (54.4%)	
Male	1787 (52.1%)	8380 (44.3%)	10167 (45.5%)	
Unknown	1 (0.0%)	22 (0.1%)	23 (0.1%)	
N	3432	18905	22337	
Age at order				<0.001 ²
Median	59.7	53.4	54.4	
Mean (sd)	55.5 (23.3)	52.7 (22.1)	53.1 (22.3)	
Q1, Q3	39.6, 72.3	35.0, 70.0	35.5, 70.4	
Range	0.0 - 120.3	0.0 - 144.7	0.0 - 144.7	
N	3432	18906	22338	
Age group (y)				<0.001 ³
0-19	278 (8.1%)	949 (5.0%)	1227 (5.5%)	
20-44	751 (21.9%)	6569 (34.8%)	7320 (32.8%)	
45-54	413 (12.0%)	2384 (12.6%)	2797 (12.5%)	
55-64	604 (17.6%)	2943 (15.6%)	3547 (15.9%)	
65-74	673 (19.6%)	2639 (14.0%)	3312 (14.8%)	
75-84	450 (13.1%)	2017 (10.7%)	2467 (11.0%)	
>=85	259 (7.6%)	1402 (7.4%)	1661 (7.4%)	
N	3428	18903	22331	
Ethnicity				0.231 ¹
Hispanic/Latino	92 (33.5%)	371 (28.3%)	463 (29.2%)	
Not Hispanic/Latino	121 (44.0%)	602 (46.0%)	723 (45.6%)	
Not Specified	62 (22.5%)	337 (25.7%)	399 (25.2%)	
N	275	1310	1585	
Race				< 0.001 ¹
Asian	220 (6.4%)	1065 (5.6%)	1285 (5.8%)	
Black or African American	620 (18.1%)	2856 (15.1%)	3476 (15.6%)	
Caucasian	963 (28.1%)	5006 (26.5%)	5969 (26.7%)	
Declined	476 (13.9%)	2432 (12.9%)	2908 (13.0%)	
Other	633 (18.4%)	3431 (18.1%)	4064 (18.2%)	
Pacific Islander	12 (0.3%)	34 (0.2%)	46 (0.2%)	
Unavailable	508 (14.8%)	4082 (21.6%)	4590 (20.5%)	
N	3432	18906	22338	
Highest level of care				<0.001 ¹
Admitted	120 (45.1%)	414 (35.6%)	534 (37.4%)	
Discharged from ED	16 (6.0%)	113 (9.7%)	129 (9.0%)	

ICU-level Care	79 (29.7%)	109 (9.4%)	188 (13.2%)	
Outpatient	51 (19.2%)	526 (45.3%)	577 (40.4%)	
N	266	1162	1428	
Encounter type at order				<0.001 ¹
Emergency	1273 (37.1%)	6197 (32.8%)	7470 (33.4%)	
Inpatient	1400 (40.8%)	4998 (26.4%)	6398 (28.6%)	
Outpatient	759 (22.1%)	7711 (40.8%)	8470 (37.9%)	
N	3432	18906	22338	
Location at order				<0.001 ¹
Allen Hospital	204 (5.9%)	1099 (5.8%)	1303 (5.8%)	
Columbia	1434 (41.8%)	7031 (37.2%)	8465 (37.9%)	
Columbia Outreach	116 (3.4%)	1144 (6.1%)	1260 (5.6%)	
Lower Manhattan	161 (4.7%)	911 (4.8%)	1072 (4.8%)	
NYHQ	495 (14.4%)	3654 (19.3%)	4149 (18.6%)	
Weill Cornell	973 (28.4%)	4936 (26.1%)	5909 (26.5%)	
WESTCHESTER	49 (1.4%)	131 (0.7%)	180 (0.8%)	
N	3432	18906	22338	
Number of admissions				<0.001 ²
Median	1.0	0.0	0.0	
Mean (sd)	0.9 (0.6)	0.4 (0.5)	0.5 (0.5)	
Q1, Q3	0.0, 1.0	0.0, 1.0	0.0, 1.0	
Range	0.0 - 2.0	0.0 - 1.0	0.0 - 2.0	
N	275	1312	1587	
Admission duration (d)				<0.001 ²
Median	5.8	0.6	0.7	
Mean (sd)	11.6 (13.5)	3.5 (9.3)	4.9 (10.6)	
Q1, Q3	1.0, 19.4	0.3, 2.9	0.3, 4.4	
Range	0.0 - 83.5	0.0 - 139.0	0.0 - 139.0	
N	275	1312	1587	
Admission 1 to test 1 (d)				0.112 ²
Median	0.4	0.3	0.3	
Mean (sd)	2.3 (6.7)	1.4 (7.5)	1.6 (7.3)	
Q1, Q3	0.2, 1.1	0.2, 0.9	0.2, 0.9	
Range	-13.9 - 56.1	0.0 - 133.1	-13.9 - 133.1	
N	268	1218	1486	
Admission 2 to test 1 (d)				<0.001 ²
Median	-9.0	0.1	-8.3	
Mean (sd)	-10.6 (7.8)	-2.6 (5.9)	-9.4 (8.1)	
Q1, Q3	-15.8, -4.0	-1.2, 0.2	-15.4, -1.5	
Range	-29.0 - 0.5	-19.1 - 1.2	-29.0 - 1.2	

N	88	16	104	
Primary Outcome	<0.001¹			
Deceased/Discharged to Hospice	27 (9.8%)	74 (5.6%)	101 (6.4%)	
Discharged (not death)	196 (71.3%)	1219 (92.9%)	1415 (89.2%)	
Intubated (still admitted)	31 (11.3%)	14 (1.1%)	45 (2.8%)	
Still admitted (not intubated)	21 (7.6%)	5 (0.4%)	26 (1.6%)	
N	275	1312	1587	
Deceased	0.038¹			
No	253 (92.0%)	1250 (95.3%)	1503 (94.7%)	
Yes	22 (8.0%)	62 (4.7%)	84 (5.3%)	
N	275	1312	1587	
Symptomatic	0.035¹			
No	4 (3.8%)	8 (1.6%)	12 (2.0%)	
Unknown	2 (1.9%)	42 (8.6%)	44 (7.4%)	
Yes	99 (94.3%)	436 (89.7%)	535 (90.5%)	
N	105	486	591	
Symptoms start to death (d)	0.320²			
Median	12.3	14.3	13.7	
Mean (sd)	17.4 (11.9)	14.0 (8.4)	14.7 (9.2)	
Q1, Q3	9.8, 25.6	6.9, 19.1	7.0, 19.1	
Range	3.3 - 39.6	1.6 - 38.3	1.6 - 39.6	
N	10	37	47	
Any fever	0.925¹			
No	25 (25.0%)	113 (25.5%)	138 (25.4%)	
Yes	75 (75.0%)	331 (74.5%)	406 (74.6%)	
N	100	444	544	
Intubated	<0.001¹			
No	208 (75.6%)	1232 (93.9%)	1440 (90.7%)	
Yes	67 (24.4%)	80 (6.1%)	147 (9.3%)	
N	275	1312	1587	
Extubated	<0.001¹			
No	251 (91.3%)	1283 (97.8%)	1534 (96.7%)	
Yes	24 (8.7%)	29 (2.2%)	53 (3.3%)	
N	275	1312	1587	
Symptoms start to intubation (d)	<0.001¹			
Median	9.0	7.0	8.0	
Mean (sd)	10.9 (7.0)	8.3 (8.4)	9.4 (7.9)	
Q1, Q3	7.0, 14.0	3.0, 10.2	4.0, 11.0	
Range	0.0 - 30.0	-3.0 - 39.0	-3.0 - 39.0	

N	41	52	93	
Intubation duration (d)				0.297 ²
Median	5.0	4.5	5.0	
Mean (sd)	6.3 (5.5)	4.7 (4.2)	5.4 (4.9)	
Q1, Q3	2.5, 8.5	1.0, 7.8	2.0, 8.0	
Range	1.0 - 20.0	-3.0 - 13.0	-3.0 - 20.0	
N	23	26	49	
Decompensated				<0.001 ¹
No	199 (72.4%)	1206 (91.9%)	1405 (88.5%)	
Yes	76 (27.6%)	106 (8.1%)	182 (11.5%)	
N	275	1312	1587	
Symptoms start to decompensation (d)				
Median	9.0	7.0	8.0	
Mean (sd)	11.7 (8.3)	9.0 (8.6)	10.1 (8.5)	
Q1, Q3	7.0, 15.0	3.5, 11.0	4.0, 13.0	
Range	0.0 - 39.0	-3.0 - 39.0	-3.0 - 39.0	
N	45	67	112	
Symptoms start to first test (d)				
Median	5.3	4.6	4.7	
Mean (sd)	9.8 (36.8)	6.3 (6.4)	6.9 (16.9)	
Q1, Q3	2.1, 8.3	2.8, 7.8	2.7, 7.9	
Range	-4.3 - 366.8	-29.4 - 60.8	-29.4 - 366.8	
N	98	431	529	
Number of tests				<0.001 ²
Median	2.0	1.0	1.0	
Mean (sd)	2.5 (1.1)	1.0 (0.0)	1.2 (0.7)	
Q1, Q3	2.0, 3.0	1.0, 1.0	1.0, 1.0	
Range	1.0 - 41.0	1.0 - 1.0	1.0 - 41.0	
N	3432	18906	22338	
Number of positive results per patient				
Median	0.0	0.0	0.0	
Mean (sd)	0.6 (0.9)	0.5 (0.5)	0.5 (0.6)	
Q1, Q3	0.0, 1.0	0.0, 1.0	0.0, 1.0	
Range	0.0 - 20.0	0.0 - 1.0	0.0 - 20.0	
N	3432	18906	22338	
Number of positive results per patient (indeterminate included)				
Median	0.0	0.0	0.0	
Mean (sd)	0.7 (1.0)	0.5 (0.5)	0.5 (0.6)	
Q1, Q3	0.0, 1.0	0.0, 1.0	0.0, 1.0	
Range	0.0 - 20.0	0.0 - 1.0	0.0 - 20.0	

N	3432	18906	22338	1.
Highest SARS-CoV-2 Result				<0.001 ¹
Invalid	1 (0.0%)	65 (0.3%)	66 (0.3%)	
Not Detected	1960 (57.1%)	9392 (49.7%)	11352 (50.8%)	
Indeterminate	100 (2.9%)	177 (0.9%)	277 (1.2%)	
Detected	1371 (39.9%)	9272 (49.0%)	10643 (47.6%)	
N	3432	18906	22338	
	Pearson's Chi-squared test (adjusted for multiple comparisons)			
2.	Linear Model ANOVA (adjusted for multiple comparisons)			
3.	Trend test for ordinal variables (adjusted for multiple comparisons)			

Supplementary Table 4: SARS-CoV-2 molecular tests in repeat-tested vs. single-tested patients. N represents the number of tests in each category.

	Repeat (N=8471)	Single (N=18906)	Total (N=27377)	p value
Method	<0.001 ¹			
cobas 6800	5841 (69.0%)	13354 (70.6%)	19195 (70.1%)	
Infiniti	1900 (22.4%)	4054 (21.4%)	5954 (21.7%)	
GeneXpert	91 (1.1%)	174 (0.9%)	265 (1.0%)	
ID Now	9 (0.1%)	44 (0.2%)	53 (0.2%)	
Panther Fusion	11 (0.1%)	15 (0.1%)	26 (0.1%)	
QuantStudio	571 (6.7%)	1224 (6.5%)	1795 (6.6%)	
7500 Fast	48 (0.6%)	41 (0.2%)	89 (0.3%)	
N	8471	18906	27377	
cobas	<0.001 ²			
No	8471 (100.0%)	18906 (100.0%)	27377 (100.0%)	
N	8471	18906	27377	
Initial result	<0.001 ¹			
Invalid	554 (6.5%)	65 (0.3%)	619 (2.3%)	
Not Detected	5613 (66.3%)	9392 (49.7%)	15005 (54.8%)	
Indeterminate	265 (3.1%)	177 (0.9%)	442 (1.6%)	
Detected	2039 (24.1%)	9272 (49.0%)	11311 (41.3%)	
N	8471	18906	27377	
Highest result on day 1	<0.001 ¹			
Invalid	367 (4.3%)	65 (0.3%)	432 (1.6%)	
Not Detected	5501 (64.9%)	9392 (49.7%)	14893 (54.4%)	
Indeterminate	267 (3.2%)	177 (0.9%)	444 (1.6%)	
Detected	2336 (27.6%)	9272 (49.0%)	11608 (42.4%)	
N	8471	18906	27377	
Highest result any day	<0.001 ¹			
Invalid	2 (0.0%)	65 (0.3%)	67 (0.2%)	
Not Detected	4775 (56.4%)	9392 (49.7%)	14167 (51.7%)	
Indeterminate	246 (2.9%)	177 (0.9%)	423 (1.5%)	
Detected	3448 (40.7%)	9272 (49.0%)	12720 (46.5%)	
N	8471	18906	27377	
Control Ct	<0.001 ³			
Median	34.0	33.9	33.9	
Mean (sd)	34.2 (1.1)	34.1 (1.0)	34.1 (1.0)	
Q1, Q3	33.5, 34.7	33.4, 34.5	33.5, 34.6	
Range	32.5 - 41.2	32.4 - 40.5	32.4 - 41.2	
N	2144	7262	9406	
Target 1 Ct	0.036 ³			

	Repeat (N=8471)	Single (N=18906)	Total (N=27377)	p value
Median	27.5	26.6	26.7	
Mean (sd)	26.7 (5.5)	26.2 (5.5)	26.3 (5.5)	
Q1, Q3	22.3, 31.4	21.8, 31.0	21.9, 31.0	
Range	13.3 - 36.8	12.4 - 38.0	12.4 - 38.0	
N	744	4533	5277	
Target 2 Ct				<0.001 ³
Median	29.1	27.3	27.6	
Mean (sd)	28.5 (6.5)	27.3 (6.1)	27.5 (6.2)	
Q1, Q3	23.2, 34.0	22.3, 32.5	22.5, 32.7	
Range	13.7 - 39.8	12.9 - 43.7	12.9 - 43.7	
N	795	4548	5343	
Target 2 Ct > 30				<0.001 ¹
FALSE	431 (54.2%)	2869 (63.1%)	3300 (61.8%)	
TRUE	364 (45.8%)	1679 (36.9%)	2043 (38.2%)	
N	795	4548	5343	

1. Pearson's Chi-squared test (adjusted for multiple comparisons)
2. Chi-squared test for given probabilities (adjusted for multiple comparisons)
3. Linear Model ANOVA (adjusted for multiple comparisons)

Supplementary Table 5: Number of SARS-CoV-2 molecular test results of repeat-tested patients on day 1 of testing. Rows represent the initial test result and columns the highest test result on day 1.

Highest First Day Result —>	Invalid (N=163)	Not Detected (N=2250)	Indeterminate (N=112)	Detected (N=907)	Total (N=3432)
Initial Result					p value < 0.001 ¹
Invalid	163 (100%)	32 (1.4%)	1 (0.9%)	45 (5.0%)	241 (7.0%)
Not Detected	0 (0.0%)	2218 (98.6%)	13 (11.6%)	49 (5.4%)	2280 (66.4%)
Indeterminate	0 (0.0%)	0 (0.0%)	98 (87.5%)	16 (1.8%)	114 (3.3%)
Detected	0 (0.0%)	0 (0.0%)	0 (0.0%)	797 (87.9%)	797 (23.2%)

1. Pearson's Chi-squared test (adjusted for multiple comparisons)

Supplementary Table 6: Number of SARS-CoV-2 molecular test results over the course of repeat testing, grouped by the initial test result on day 1.

Highest Result Any Day —>	Invalid (N=1)	Not Detected (N=1960)	Indeterminate (N=100)	Detected (N=1371)	Total (N=3432)
Initial Result					p-value < 0.001 ¹
Invalid	1 (100%)	110 (5.6%)	3 (3.0%)	127 (9.3%)	241 (7.0%)
Not Detected	0 (0.0%)	1850 (94.4%)	45 (45.0%)	385 (28.1%)	2280 (66.4%)
Indeterminate	0 (0.0%)	0 (0.0%)	52 (52.0%)	62 (4.5%)	114 (3.3%)
Detected	0 (0.0%)	0 (0.0%)	0 (0.0%)	797 (58.1%)	797 (23.2%)

1. Pearson's Chi-squared test (adjusted for multiple comparisons)

Supplementary Table 7: Characteristics of initially positive repeat-tested patients comparing those that converted from positive to negative (Pos2neg) with those that remained positive (Unchanged pos).

	Pos2neg (N=330)	Unchanged pos (N=413)	Total (N=743)	p value
Gender at birth				0.599 ¹
Female	138 (41.8%)	162 (39.2%)	300 (40.4%)	
Male	192 (58.2%)	251 (60.8%)	443 (59.6%)	
N	330	413	743	
Age at order				0.014 ²
Median	58.6	62.6	61.3	
Mean (sd)	56.4 (18.2)	60.3 (18.3)	58.6 (18.4)	
Q1, Q3	46.2, 68.2	49.7, 73.1	48.2, 70.9	
Range	0.0 - 100.4	0.0 - 96.5	0.0 - 100.4	
N	330	413	743	
Age group (y)				0.014 ³
0-19	11 (3.3%)	11 (2.7%)	22 (3.0%)	
20-44	64 (19.4%)	71 (17.2%)	135 (18.2%)	
45-54	62 (18.8%)	60 (14.5%)	122 (16.4%)	
55-64	80 (24.2%)	83 (20.1%)	163 (21.9%)	
65-74	72 (21.8%)	102 (24.7%)	174 (23.4%)	
75-84	30 (9.1%)	62 (15.0%)	92 (12.4%)	
>=85	11 (3.3%)	24 (5.8%)	35 (4.7%)	
N	330	413	743	
Ethnicity				0.721 ¹
Hispanic/Latino	15 (30.6%)	15 (39.5%)	30 (34.5%)	
Not Hispanic/Latino	22 (44.9%)	14 (36.8%)	36 (41.4%)	
Not Specified	12 (24.5%)	9 (23.7%)	21 (24.1%)	
N	49	38	87	
Race				0.314 ¹
Asian	24 (7.3%)	19 (4.6%)	43 (5.8%)	
Black or African American	57 (17.3%)	66 (16.0%)	123 (16.6%)	
Caucasian	81 (24.5%)	115 (27.8%)	196 (26.4%)	
Declined	55 (16.7%)	47 (11.4%)	102 (13.7%)	
Other	62 (18.8%)	91 (22.0%)	153 (20.6%)	
Pacific Islander	3 (0.9%)	3 (0.7%)	6 (0.8%)	
Unavailable	48 (14.5%)	72 (17.4%)	120 (16.2%)	
N	330	413	743	
Highest level of care				0.241 ¹
Admitted	12 (27.9%)	11 (29.7%)	23 (28.8%)	
Discharged from ED	5 (11.6%)	1 (2.7%)	6 (7.5%)	

	Pos2neg (N=330)	Unchanged pos (N=413)	Total (N=743)	p value
ICU-level Care	22 (51.2%)	15 (40.5%)	37 (46.2%)	
Outpatient	4 (9.3%)	10 (27.0%)	14 (17.5%)	
N	43	37	80	
Encounter type at order				< 0.001 ¹
Emergency	19 (5.8%)	56 (13.6%)	75 (10.1%)	
Inpatient	216 (65.5%)	278 (67.3%)	494 (66.5%)	
Outpatient	95 (28.8%)	79 (19.1%)	174 (23.4%)	
N	330	413	743	
Location at order				0.018 ¹
Allen Hospital	9 (2.7%)	18 (4.4%)	27 (3.6%)	
Columbia	162 (49.1%)	165 (40.0%)	327 (44.0%)	
Columbia Outreach	0 (0.0%)	8 (1.9%)	8 (1.1%)	
Lower Manhattan	7 (2.1%)	11 (2.7%)	18 (2.4%)	
Queens	35 (10.6%)	73 (17.7%)	108 (14.5%)	
Weill Cornell	114 (34.5%)	136 (32.9%)	250 (33.6%)	
Westchester	3 (0.9%)	2 (0.5%)	5 (0.7%)	
N	330	413	743	
Number of admissions				0.629 ²
Median	1.0	1.0	1.0	
Mean (sd)	0.7 (0.5)	0.8 (0.7)	0.8 (0.6)	
Q1, Q3	0.0, 1.0	0.0, 1.0	0.0, 1.0	
Range	0.0 - 2.0	0.0 - 2.0	0.0 - 2.0	
N	49	38	87	
Admission duration (d)				0.314 ²
Median	22.9	6.6	12.3	
Mean (sd)	17.6 (16.4)	13.1 (13.0)	15.7 (15.1)	
Q1, Q3	0.4, 28.3	1.0, 27.7	0.7, 28.2	
Range	0.0 - 82.2	0.2 - 38.6	0.0 - 82.2	
N	49	38	87	
Admission 1 to test 1 (d)				0.014 ²
Median	28.1	20.6	25.7	
Mean (sd)	28.3 (10.4)	20.7 (12.2)	24.7 (11.9)	
Q1, Q3	23.1, 33.4	12.2, 31.5	16.3, 32.9	
Range	0.8 - 69.0	0.4 - 42.6	0.4 - 69.0	
N	43	38	81	
Admission 2 to test 1 (d)				0.182 ²
Median	8.6	0.2	0.3	
Mean (sd)	11.8 (12.3)	3.8 (8.1)	6.4 (10.2)	

	Pos2neg (N=330)	Unchanged pos (N=413)	Total (N=743)	p value
Q1, Q3	0.7, 21.3	0.2, 1.5	0.2, 8.3	
Range	0.2 - 32.6	0.1 - 28.3	0.1 - 32.6	
N	8	16	24	
Primary Outcome				0.241 ¹
Deceased/Discharged to Hospice	1 (2.0%)	3 (7.9%)	4 (4.6%)	
Discharged (not death)	24 (49.0%)	24 (63.2%)	48 (55.2%)	
Intubated (still admitted)	17 (34.7%)	10 (26.3%)	27 (31.0%)	
Still admitted (not intubated)	7 (14.3%)	1 (2.6%)	8 (9.2%)	
N	49	38	87	
Deceased				0.314 ¹
No	48 (98.0%)	35 (92.1%)	83 (95.4%)	
Yes	1 (2.0%)	3 (7.9%)	4 (4.6%)	
N	49	38	87	
Symptomatic				
No	1 (2.5%)	2 (8.7%)	3 (4.8%)	
Unknown	0 (0.0%)	0 (0.0%)	0 (0.0%)	
Yes	39 (97.5%)	21 (91.3%)	60 (95.2%)	
N	40	23	63	
Symptoms start to death (d)				0.327 ²
Median	39.6	13.7	22.0	
Mean (sd)	39.6 (NA)	18.2 (10.6)	23.5 (13.8)	
Q1, Q3	39.6, 39.6	12.1, 22.0	12.9, 32.6	
Range	39.6 - 39.6	10.6 - 30.3	10.6 - 39.6	
N	1	3	4	
Any fever				0.744 ¹
No	11 (28.2%)	5 (23.8%)	16 (26.7%)	
Yes	28 (71.8%)	16 (76.2%)	44 (73.3%)	
N	39	21	60	
Intubated				0.327 ¹
No	26 (53.1%)	25 (65.8%)	51 (58.6%)	
Yes	23 (46.9%)	13 (34.2%)	36 (41.4%)	
N	49	38	87	
Extubated				0.128 ¹
No	44 (89.8%)	38 (100.0%)	82 (94.3%)	
Yes	5 (10.2%)	0 (0.0%)	5 (5.7%)	
N	49	38	87	

	Pos2neg (N=330)	Unchanged pos (N=413)	Total (N=743)	p value
Symptoms start to intubation (d)	0.721 ²			
Median	10.0	9.0	9.0	
Mean (sd)	10.5 (5.6)	11.6 (8.1)	10.9 (6.4)	
Q1, Q3	8.0, 13.0	6.5, 19.0	7.0, 14.2	
Range	0.0 - 23.0	2.0 - 25.0	0.0 - 25.0	
N	21	11	32	
Intubation duration (d)				
Median	11.0	NA	11.0	
Mean (sd)	11.8 (5.1)	NA	11.8 (5.1)	
Q1, Q3	11.0, 11.0	NA	11.0, 11.0	
Range	6.0 - 20.0	NA	6.0 - 20.0	
N	5	0	5	
Decompensated	0.343 ¹			
No	25 (51.0%)	24 (63.2%)	49 (56.3%)	
Yes	24 (49.0%)	14 (36.8%)	38 (43.7%)	
N	49	38	87	
Symptoms start to decompensation (d)	0.980 ²			
Median	10.0	9.0	9.5	
Mean (sd)	11.8 (8.1)	11.7 (7.7)	11.8 (7.9)	
Q1, Q3	8.0, 13.8	6.8, 18.0	7.0, 14.8	
Range	0.0 - 39.0	2.0 - 25.0	0.0 - 39.0	
N	22	12	34	
Symptoms start to first test (d)	0.314 ²			
Median	5.8	7.0	6.2	
Mean (sd)	6.7 (5.1)	24.0 (78.8)	12.9 (47.6)	
Q1, Q3	2.2, 9.1	2.5, 8.6	2.3, 9.0	
Range	-1.3 - 19.4	0.6 - 366.8	-1.3 - 366.8	
N	37	21	58	
Number of tests	< 0.001 ²			
Median	3.0	2.0	2.0	
Mean (sd)	3.2 (2.4)	2.3 (0.6)	2.7 (1.7)	
Q1, Q3	2.0, 4.0	2.0, 2.0	2.0, 3.0	
Range	2.0 - 41.0	2.0 - 7.0	2.0 - 41.0	
N	330	413	743	
Number of positive results per patient	< 0.001 ²			
Median	1.0	2.0	2.0	
Mean (sd)	1.4 (1.2)	2.0 (0.6)	1.8 (1.0)	
Q1, Q3	1.0, 2.0	2.0, 2.0	1.0, 2.0	

	Pos2neg (N=330)	Unchanged pos (N=413)	Total (N=743)	p value
Range	1.0 - 20.0	1.0 - 7.0	1.0 - 20.0	
N	330	413	743	

1. Pearson's Chi-squared test (adjusted for multiple comparisons)
2. Linear Model ANOVA (adjusted for multiple comparisons)
3. Trend test for ordinal variables (adjusted for multiple comparisons)

Supplementary Table 8: Characteristics of repeat-tested initially negative patients comparing those that converted from negative to positive (Neg2pos), with those that remained negative (Unchanged neg) or indeterminate (Unchanged ind) patients.

	Neg2pos (N=417)	Pos2neg (N=4)	Unchanged ind (N=43)	Unchanged neg (N=1705)	Total (N=2169)	p value
Gender at birth						0.062 ¹
Female	192 (46.0%)	1 (25.0%)	20 (46.5%)	900 (52.8%)	1113 (51.3%)	
Male	225 (54.0%)	3 (75.0%)	23 (53.5%)	805 (47.2%)	1056 (48.7%)	
N	417	4	43	1705	2169	
Age at order						< 0.001 ²
Median	63.6	68.1	63.4	57.1	58.9	
Mean (sd)	61.3 (18.8)	65.9 (6.7)	61.4 (14.4)	51.6 (26.2)	53.7 (25.1)	
Q1, Q3	49.1, 75.0	64.8, 69.2	52.1, 71.6	33.6, 72.0	36.3, 72.7	
Range	0.0 - 97.4	56.0 - 71.1	31.3 - 86.4	0.0 - 120.3	0.0 - 120.3	
N	417	4	43	1705	2169	
Age group (y)						< 0.001 ³
0-19	4 (1.0%)	0 (0.0%)	0 (0.0%)	238 (14.0%)	242 (11.2%)	
20-44	85 (20.4%)	0 (0.0%)	7 (16.3%)	401 (23.5%)	493 (22.7%)	
45-54	47 (11.3%)	0 (0.0%)	7 (16.3%)	172 (10.1%)	226 (10.4%)	
55-64	84 (20.1%)	1 (25.0%)	12 (27.9%)	248 (14.5%)	345 (15.9%)	
65-74	93 (22.3%)	3 (75.0%)	8 (18.6%)	301 (17.7%)	405 (18.7%)	
75-84	61 (14.6%)	0 (0.0%)	7 (16.3%)	221 (13.0%)	289 (13.3%)	
>=85	43 (10.3%)	0 (0.0%)	2 (4.7%)	124 (7.3%)	169 (7.8%)	
N	417	4	43	1705	2169	
Ethnicity						
Hispanic / Latino	23 (41.1%)	0	1 (100.0%)	25 (28.4%)	49 (33.8%)	
Not Hispanic/ Latino	22 (39.3%)	0	0 (0.0%)	44 (50.0%)	66 (45.5%)	
Not Specified	11 (19.6%)	0	0 (0.0%)	19 (21.6%)	30 (20.7%)	
N	56	0	1	88	145	
Race						0.085 ¹
Asian	30 (7.2%)	1 (25.0%)	1 (2.3%)	120 (7.0%)	152 (7.0%)	
Black or African American	88 (21.1%)	1 (25.0%)	9 (20.9%)	295 (17.3%)	393 (18.1%)	
Caucasian	94 (22.5%)	0 (0.0%)	10 (23.3%)	530 (31.1%)	634 (29.2%)	
Declined	49 (11.8%)	0 (0.0%)	7 (16.3%)	245 (14.4%)	301 (13.9%)	
Other	86 (20.6%)	1 (25.0%)	11 (25.6%)	279 (16.4%)	377 (17.4%)	
Pacific Islander	1 (0.2%)	0 (0.0%)	0 (0.0%)	4 (0.2%)	5 (0.2%)	
Unavailable	69 (16.5%)	1 (25.0%)	5 (11.6%)	232 (13.6%)	307 (14.2%)	
N	417	4	43	1705	2169	

	Neg2pos (N=417)	Pos2neg (N=4)	Unchanged ind (N=43)	Unchanged neg (N=1705)	Total (N=2169)	p value
Highest level of care						
Admitted	30 (53.6%)	0	1 (100.0%)	43 (50.0%)	74 (51.7%)	
Discharged from ED	2 (3.6%)	0	0 (0.0%)	3 (3.5%)	5 (3.5%)	
ICU-level Care	10 (17.9%)	0	0 (0.0%)	23 (26.7%)	33 (23.1%)	
Outpatient	14 (25.0%)	0	0 (0.0%)	17 (19.8%)	31 (21.7%)	
N	56	0	1	86	143	
Encounter type at order						
Emergency	84 (20.1%)	0 (0.0%)	3 (7.0%)	161 (9.4%)	248 (11.4%)	
Inpatient	228 (54.7%)	4 (100.0%)	26 (60.5%)	1130 (66.3%)	1388 (64.0%)	
Outpatient	105 (25.2%)	0 (0.0%)	14 (32.6%)	414 (24.3%)	533 (24.6%)	
N	417	4	43	1705	2169	
Location at order						
Allen Hospital	38 (9.1%)	0 (0.0%)	4 (9.3%)	67 (3.9%)	109 (5.0%)	
Columbia	178 (42.7%)	1 (25.0%)	21 (48.8%)	731 (42.9%)	931 (42.9%)	
Columbia Outreach	16 (3.8%)	0 (0.0%)	1 (2.3%)	39 (2.3%)	56 (2.6%)	
Lower Manhattan	11 (2.6%)	1 (25.0%)	3 (7.0%)	86 (5.0%)	101 (4.7%)	
Queens	100 (24.0%)	0 (0.0%)	1 (2.3%)	223 (13.1%)	324 (14.9%)	
Weill Cornell	69 (16.5%)	2 (50.0%)	12 (27.9%)	524 (30.7%)	607 (28.0%)	
Westchester	5 (1.2%)	0 (0.0%)	1 (2.3%)	35 (2.1%)	41 (1.9%)	
N	417	4	43	1705	2169	
Number of admissions						
Median	1.0	NA	1.0	1.0	1.0	
Mean (sd)	1.0 (0.8)	NA	1.0 (NA)	0.9 (0.6)	1.0 (0.7)	
Q1, Q3	0.0, 2.0	NA	1.0, 1.0	0.8, 1.0	0.0, 1.0	
Range	0.0 - 2.0	NA	1.0 - 1.0	0.0 - 2.0	0.0 - 2.0	
N	56	0	1	88	145	
Admission duration (d)						
Median	5.6	NA	2.9	7.4	7.2	
Mean (sd)	8.5 (8.1)	NA	2.9 (NA)	13.2 (15.6)	11.3 (13.3)	
Q1, Q3	1.4, 11.9	NA	2.9, 2.9	1.5, 18.4	1.5, 15.4	
Range	0.2 - 33.0	NA	2.9 - 2.9	0.0 - 83.5	0.0 - 83.5	
N	56	0	1	88	145	
Admission 1 to test 1 (d)						
Median	8.5	NA	1.6	13.2	9.5	
Mean (sd)	12.1 (11.7)	NA	1.6 (NA)	17.5 (16.0)	15.3 (14.7)	
Q1, Q3	3.5, 20.3	NA	1.6, 1.6	3.3, 29.0	3.4, 25.7	

	Neg2pos (N=417)	Pos2neg (N=4)	Unchanged ind (N=43)	Unchanged neg (N=1705)	Total (N=2169)	p value
Range	0.5 - 42.9	NA	1.6 - 1.6	0.1 - 74.1	0.1 - 74.1	
N	56	0	1	87	144	
Admission 2 to test 1 (d)						
Median	0.2	NA	NA	1.2	0.3	
Mean (sd)	0.4 (0.9)	NA	NA	4.7 (8.0)	2.4 (5.9)	
Q1, Q3	0.1, 0.4	NA	NA	0.2, 4.1	0.1, 1.8	
Range	-3.8 - 2.0	NA	NA	0.0 - 33.0	-3.8 - 33.0	
N	33	0	0	29	62	
Primary Outcome						
Deceased/ Discharged to Hospice	7 (12.5%)	0	0 (0.0%)	9 (10.2%)	16 (11.0%)	
Discharged (not death)	43 (76.8%)	0	1 (100.0%)	68 (77.3%)	112 (77.2%)	
Intubated (still admitted)	3 (5.4%)	0	0 (0.0%)	1 (1.1%)	4 (2.8%)	
Still admitted (not intubated)	3 (5.4%)	0	0 (0.0%)	10 (11.4%)	13 (9.0%)	
N	56	0	1	88	145	
Deceased						
No	51 (91.1%)	0	1 (100.0%)	81 (92.0%)	133 (91.7%)	
Yes	5 (8.9%)	0	0 (0.0%)	7 (8.0%)	12 (8.3%)	
N	56	0	1	88	145	
Symptomatic						
No	1 (3.8%)	0	0 (0.0%)	0 (0.0%)	1 (3.4%)	
Unknown	1 (3.8%)	0	0 (0.0%)	1 (50.0%)	2 (6.9%)	
Yes	24 (92.3%)	0	1 (100.0%)	1 (50.0%)	26 (89.7%)	
N	26	0	1	2	29	
Symptoms start to death (d)						
Median	18.5	NA	NA	NA	18.5	
Mean (sd)	18.5 (10.7)	NA	NA	NA	18.5 (10.7)	
Q1, Q3	14.7, 22.3	NA	NA	NA	14.7, 22.3	
Range	10.9 - 26.1	NA	NA	NA	10.9 - 26.1	
N	2	0	0	0	2	
Any fever						
No	6 (24.0%)	0	0 (0.0%)	0 (0.0%)	6 (22.2%)	
Yes	19 (76.0%)	0	1 (100.0%)	1 (100.0%)	21 (77.8%)	
N	25	0	1	1	27	
Intubated						
No	49 (87.5%)	0	1 (100.0%)	71 (80.7%)	121 (83.4%)	

	Neg2pos (N=417)	Pos2neg (N=4)	Unchanged ind (N=43)	Unchanged neg (N=1705)	Total (N=2169)	p value
Yes	7 (12.5%)	0	0 (0.0%)	17 (19.3%)	24 (16.6%)	
N	56	0	1	88	145	
Extubated						
No	54 (96.4%)	0	1 (100.0%)	75 (85.2%)	130 (89.7%)	
Yes	2 (3.6%)	0	0 (0.0%)	13 (14.8%)	15 (10.3%)	
N	56	0	1	88	145	
Symptoms start to intubation (d)						
Median	11.0	NA	NA	7.0	10.5	
Mean (sd)	14.4 (11.5)	NA	NA	7.0 (NA)	13.2 (10.7)	
Q1, Q3	10.0, 21.0	NA	NA	7.0, 7.0	7.8, 18.5	
Range	0.0 - 30.0	NA	NA	7.0 - 7.0	0.0 - 30.0	
N	5	0	0	1	6	
Intubation duration (d)						
Median	3.5	NA	NA	3.0	3.0	
Mean (sd)	3.5 (3.5)	NA	NA	5.0 (5.6)	4.8 (5.2)	
Q1, Q3	2.2, 4.8	NA	NA	2.0, 5.0	2.0, 5.0	
Range	1.0 - 6.0	NA	NA	1.0 - 20.0	1.0 - 20.0	
N	2	0	0	12	14	
Decompensated						
No	46 (82.1%)	0	1 (100.0%)	68 (77.3%)	115 (79.3%)	
Yes	10 (17.9%)	0	0 (0.0%)	20 (22.7%)	30 (20.7%)	
N	56	0	1	88	145	
Symptoms start to decompensation (d)						
Median	16.0	NA	NA	7.0	11.0	
Mean (sd)	16.3 (11.3)	NA	NA	7.0 (NA)	15.0 (10.9)	
Q1, Q3	10.2, 24.7	NA	NA	7.0, 7.0	8.5, 23.5	
Range	0.0 - 30.0	NA	NA	7.0 - 7.0	0.0 - 30.0	
N	6	0	0	1	7	
Symptoms start to first test (d)						
Median	3.6	NA	9.1	4.4	4.4	
Mean (sd)	5.0 (5.2)	NA	9.1 (NA)	4.4 (NA)	5.1 (5.1)	
Q1, Q3	1.3, 7.8	NA	9.1, 9.1	4.4, 4.4	1.6, 7.9	
Range	-4.3 - 15.9	NA	9.1 - 9.1	4.4 - 4.4	-4.3 - 15.9	
N	25	0	1	1	27	
Number of tests						
Median	2.0	3.5	2.0	2.0	2.0	
Mean (sd)	2.4 (0.8)	3.5 (0.6)	2.3 (0.6)	2.5 (0.9)	2.5 (0.9)	
Q1, Q3	2.0, 3.0	3.0, 4.0	2.0, 2.0	2.0, 3.0	2.0, 3.0	

0.009²

	Neg2pos (N=417)	Pos2neg (N=4)	Unchanged ind (N=43)	Unchanged neg (N=1705)	Total (N=2169)	p value
Range	2.0 - 9.0	3.0 - 4.0	2.0 - 5.0	2.0 - 8.0	2.0 - 9.0	
N	417	4	43	1705	2169	
Number of positive results per patient						
Median	1.0	1.0	1.0	0.0	0.0	
Mean (sd)	1.1 (0.3)	1.0 (0.0)	1.1 (0.6)	0.0 (0.0)	0.2 (0.5)	
Q1, Q3	1.0, 1.0	1.0, 1.0	1.0, 1.0	0.0, 0.0	0.0, 0.0	
Range	1.0 - 3.0	1.0 - 1.0	1.0 - 4.0	0.0 - 0.0	0.0 - 4.0	
N	417	4	43	1705	2169	

1. Pearson's Chi-squared test (adjusted for multiple comparisons)
2. Linear Model ANOVA (adjusted for multiple comparisons)
3. Trend test for ordinal variables (adjusted for multiple comparisons)

< 0.001²