Supplementary Figures



S1 Fig. Visualization of WM ROIs. The colored WM ROIs are overlaid on the T1 template.



S2 Fig. Mean of GM-GM functional correlation matrices ($mFCM_{GG}$) for each clinical group in AD progression. (**a-f**) $mFCM_{GG}$ for group of CN (**a**), SMC (**b**), EMCI (**c**), MCI (**d**), LMCI (**e**) and ADD (**f**). Each element in $mFCM_{GG}$ is the group mean of correlation coefficient of the averaged BOLD time courses between two GM regions. See **Table 1** for the list of GM ROIs.



S3 Fig. Raw overall-connectivity and neuropsychological scores for each clinical group in AD progression. (**a**) Raw group mean (gray/black solid square) and standard deviation of mean (gray/black solid bar) of the overall-connectivity, i.e., average over all elements in FCM, for each clinical group. The six clinical groups are CN, SMC, EMCI, MCI, LMCI and AD groups. (**b**-**h**) Raw mean (colored solid square) and standard deviation (colored solid bar) of neuropsychological scores: MMSE (**b**), CDR-Global (**c**), CDR-SOB (**d**), GDS (**e**), FAQ (**f**), WMS-LMII (**g**), and ADAS-Cog (**h**).