

Table S1: Key epidemiological parameters used in the SEIHURD model, with their mean estimates and respective estimates obtained according to best fit for Salvador, inland cities and all Bahia state. The parameters interval was consolidated by Supplementary Table 2 and File 3.

Parameter	Description	Interval	Fixed	Estimated (95% CI) Bahia	Estimated (95% CI) Salvador	Estimated (95% CI) Interior (95% CI) April 01 th
β_0	Pre-intervention transmission rate	[0, 2]	-	1.06 (1.04 - 1.07)	1.24 (1.23 - 1.25)	0.95 (0.93 - 0.96)
β_1	Post-intervention transmission rate	[0, 2]	-	0.63 (0.62 - 0.64)	0.59 (0.58 - 0.59)	0.63 (0.62 - 0.64)
t_1	Time of transmission rate change	[March 15 th , April 15 th]	-	March 28 th	March 26 th	April 01 th
δ	Factor that reduce the asymptomatic infectivity	[0, 0.7]	-	0.62 (0.60 - 0.64)	0.55 (0.54 - 0.56)	0.57 (0.56 - 0.58)
p	Proportion of latent (E) that proceed to symptomatic infective	[0.13, 0.5]	0.2	-	-	-
κ	Mean exposed period	[1/6, 1/3]	1/4	-	-	-
γ_a	Mean asymptomatic period	[1/3.70, 1/3.24]	1/3.5	-	-	-
γ_s	Mean symptomatic period	[1/5, 1/3]	1/4	-	-	-
h	Proportion of symptomatic needing hospitalization or ICU	[0.05, 0.25]	-	0.28 (0.28, 0.29)	-	-
$1 - \xi$	Proportion of symptomatic that proceed to ICU	[0.01, 0.5]	0.47	-	-	-
γ_H	Mean Hospital period	[1/12, 1/4]	-	0.14 (0.13, 0.14)	-	-
γ_U	Mean ICU period	[1/12, 1/3]	-	0.14 (0.14, 0.15)	-	-
μ_H	Death rate of hospitalized individuals	[0.1, 0.2]	0.15	-	-	-
μ_U	Death rate of ICU individuals	[0.4, 0.5]	0.4	-	-	-
ω_H	Proportion of hospitalized that goes to ICU	[0.1, 0.3]	0.14	-	-	-
ω_U	Proportion of ICU that goes to hospitalization	[0.1, 0.3]	0.29	-	-	-