

Traces of SARS-CoV-2 RNA in the Blood of COVID-19 Patients

Ahmed Moustafa^{1,*}, Ramy K. Aziz^{2,3,*}

Affiliations

¹ Department of Biology, American University in Cairo, New Cairo, Egypt

² Department of Microbiology and Immunology, Faculty of Pharmacy, Cairo University, Cairo, Egypt

³ Center for Genome and Microbiome Research, Cairo University, Cairo, Egypt

* Corresponding author

Emails

AM: amoustafa@aucegypt.edu

RKA: ramy.aziz@gmail.com

Short title

SARS-CoV-2 RNA in COVID-19 PBMC

Keywords

SARS-CoV-2, Coronavirus, COVID-19, Blood, PBMC, RNA-Seq

Supplementary Data

Table S1. Summary of the samples included in the study, raw hits raw and estimated abundance metrics

Nine RNA-Seq samples (Xiong et al. 2020) were analyzed, including 3 PBMC from healthy donors, 3 PBMC from patients, and 2 BALF from patients with an additional replicate each. The numbers are paired-end reads.

Sample	Description	Total Reads	Viral Reads	Viral read abundance (ppm)	Actin transcripts	Spike-to-actin RNA ratio
CRR119890	PBMC: Healthy Donor 1	78,077,390			48.52195	0
CRR125445	PBMC: Healthy Donor 2	47,983,296			27.3257	0
CRR125446	PBMC: Healthy Donor 3	47,804,999			49.644	0
CRR119891	PBMC: Patient 1	76,420,796	2	0.03	77.30287	0.000004696
CRR119892	PBMC: Patient 2	76,333,057		0.00	76.5192	0
CRR119893	PBMC: Patient 3	78,523,460		0.00	77.5553	0
CRR119894	BALF: Patient 1 rep1	8,254,725	272,184	32,973.12	24.0391	9.702268388
CRR119895	BALF: Patient 1 rep2	3,671,406	107,423	29,259.36	13.3584	8.889350521
CRR119896	BALF: Patient 2 rep1	4,660,406	64,058	13,745.15	20.762	2.450953665
CRR119897	BALF: Patient 2 rep2	2,251,299	27,334	12,141.43	13.0798	2.526200706

References

- Xiong, Yong, Yuan Liu, Liu Cao, Dehe Wang, Ming Guo, Ao Jiang, Dong Guo, et al. 2020. "Transcriptomic Characteristics of Bronchoalveolar Lavage Fluid and Peripheral Blood Mononuclear Cells in COVID-19 Patients." *Emerging Microbes & Infections* 9 (1): 761–70.