

Table 1: Characteristics of included studies

Serial number	Study	Sample size(F/M)	Mean age, years (SD)	Description of PCI	Experimental intervention	Control intervention	Outcome measurement
22	Abdelhalem 2018	Experimental: 20(2/18); Control: 20(4/16)	54.65(7.63) 51.95(8.07)	PCI with bare metal stent and drug-eluting stent implantation	HIIT program consisted of 5 min of warm-up exercises followed 2–5 min higher intensity at 85–95% of HRR with similar time of moderate-intensity exercise, and end by 5 min of cool down. This training started within 2 weeks after PCI, 2 times a week for 3 months.	MICT program consisted of 5 min of warm-up exercises followed 30–35 min continuous treadmill exercise at 40–60% of HRR, and end by 5 min of cooldown. This training started within 2 weeks after PCI, 2 times/week for 3 months.	METS, EF%, QoL, laboratory index (s. Creatinine, Hb, platelet count, WBCs, s. Cholesterol, LDL, HDL, s. TGs)
23	Gao ZZ 2015	Experimental: 22(4/18); Control: ①21(5/16) ②22(7/17)	59.4(7.9) ①61.2(8.0) ②60.4(8.6)	PCI with stent implantation	Treadmill training at the intensity of 80% of PP in CPET was exercised for 3 min interval with 1 min rest in all 10 groups for 40 min. Additionally, 60% PP can be preloaded for 1 week. The program lasted 12 weeks in 3 times per week.	① Treadmill training at the intensity of 60% PP in CPET was exercised for 40 min continuously. The program lasted 12 weeks in 3 times per week. ② Usual care	LVEF, PP, ED, VO <sub>2</sub> peak, AT
24	Kim 2015	Experimental: 14(2/12); Control: 14(4/10)	57(11.58) 60.2(13.64)	PCI with drug-eluting stent implantation	HIIT program consisted of a 10-min warm-up at 50%-70% of heart rate reserve(HRR), followed by four times of 4-min intervals of walking at 85%-95% of HRR with three active pauses of 3-min walking and 10-min cooldown at 50%-70% of HRR. This training started within 3 weeks after PCI, three times a week for 6 weeks.	MICT program contained a 10-min warm-up, followed by a 25-min walk on a treadmill continuously at 70%-85% of HRR, and a 10-min cooldown. This training exercised for a total of 45 mins.	HDL, LDL, hs-CRP
25	Munk 2009(1)	Experimental: 20(3/17); Control: 20(4/16)	57(14) 61(10)	PCI with metal stent or drug-eluting stent	HIIT program consisted of a warm-up period, followed by four 4-minute intervals at 80–90% of HRmax along with 3 minutes of active recovery at	Usual care	VO <sub>2</sub> peak; endothelial function; C-reactive protein

				implantation	60–70% of HRmax. There were 5 minutes cool-down, 10 minutes abdominal and spine-resistance exercises, and 5 minutes of stretching and relaxing. The program started $11 \pm 4$ days after PCI lasted for 6 months in 3 times a week.		
26	Munk 2010	Experimental: 20(3/17); Control: 18(3/15)	57.7(10.4) 59.7(8.5)	PCI	HIIT program consisted of a warm-up period, followed by four 4-minute intervals at 80–90% of HRmax along with 3 minutes of active recovery at 60–70% of HRmax. There were 5 minutes cool-down, 10 minutes abdominal- and spine-resistance exercises, and 5 minutes of stretching and relaxing. The program started $11 \pm 4$ days after PCI lasted for 6 months in 3 times a week.	Usual care	Heart rate variability (HRV)
27	Munk 2011	Experimental: 18(2/16); Control: 18(4/14)	59.5(10) 60.7(9)	PCI	HIIT program consisted of a warm-up period, followed by four 4-minute intervals at 80–90% of HRmax along with 3 minutes of active recovery at 60–70% of HRmax. There were 5 minutes cool-down, 10 minutes abdominal- and spine-resistance exercises, and 5 minutes of stretching and relaxing. The program started $11 \pm 4$ days after PCI lasted for 6 months in 3 times a week.	Usual care	Inflammatory and anti-inflammatory markers; Markers of endothelium activation; Makers of platelet-mediated inflammation; chemokines