

Using simulation to assess the potential effectiveness of implementing screening at national borders during international outbreaks of influenza, SARS, Ebola virus disease and COVID-19

Supplementary Material

Declan Bays¹, Emma Bennett¹, Thomas Finnie¹

Border Screening Model

For our model to run computationally, we break down the model description given in the papers main body into a form easily transferable into computer code. This takes the following form:

```
Non_fliers = 0  
Border_detected = 0  
Non_detected = 0  
For 0 ≤ i ≤ n do  
    Sample tinc, texp, tflight from Dexp, Dinc, Dflight  
    If tinc ≤ texp, do  
        No_fly_count += 1  
    Else if tinc ≤ texp + tflight, do  
        If weighted_coin_flip(α), do  
            If weighted_coin_flip(β), do  
                Border_detected += 1  
            Else do  
                Non_detected += 1  
            Fliers = n – Non_fliers  
            Detection_ratio = Border_detected / Fliers  
        Return Detection_ratio
```

Where we use the function **weighted_coin_flip(x)**, defined to return the value True with a probability x, to simulate the random process of a symptomatic case being screened and subsequently detected. Note that we do not need to consider whether non-symptomatic persons undergo screening, as they will not be detectable, and this will just add processing time.

¹ Emergency Response Department, Public Health England, Porton Down, Salisbury, SP4 0JG

Calculating incubation parameters for Ebola and influenza

In the referenced papers (Nishiura et al., 2009; Pettey et al., 2017), the authors state that a gamma distribution was used to fit incubation period in both cases. However, in the Pettey paper only the mean and the 95th percentile was given, while in the Nishiura paper only the mean and the variance was given.

To obtain shape and scale parameters from these values, we used two different methods for each of the different types of stated values. For both cases we applied the following formula:

$$\text{shape} = \left(\frac{\text{mean}}{\text{standard deviation}} \right)^2$$
$$\text{scale} = \frac{(\text{standard deviation})^2}{\text{mean}}$$

For the influenza parameters, where the mean and variance (hence standard deviation) was given in the text, this was a simple case of entering the values into the formulas. For the Ebola parameters however (where only the mean and the 95th percentile was given), this was done using Python to create a range of values for the standard deviation. From here, a range of values for the shape and scale parameters were then calculated using these formulas; one for each value in the range of standard deviations. A gamma distribution was then created from each pair of shape and scale parameters using pythons SciPy package. Each of these distributions was then assessed for the value of the 95th percentile, and the distribution which gave a value closest to the reported value in the paper was then selected, and the corresponding parameters recorded.

Calculating incubation parameters for COVID-19

In table 1 of the referenced paper (Linton et al., 2020), it is given that the mean and standard deviation for the incubation period of infected persons (obtained from 158 recordings of cases which include Wuhan resident) were 5.6 and 2.8 respectively. We can then transform these values to obtain the lognormal parameters for this distribution. Our parameterisation requires the values μ and σ , which are obtained through the formulas (see https://en.wikipedia.org/wiki/Log-normal_distribution):

$$\mu = \ln \left(\frac{(\text{mean})^2}{\sqrt{(\text{standard deviation})^2 + (\text{mean})^2}} \right)$$
$$\sigma^2 = \ln \left(1 + \frac{(\text{standard deviation})^2}{(\text{mean})^2} \right)$$