## **Supplementary Data**




Figure S 1: Meta-regression bubble plots of (a) IL-6 and (b) IL-10 SMD with percent difference of male patients across severe and non-severe groups




Figure S 2: Standardized Mean Difference of Neutrophil-to-Lymphocyte ratio across severe and non-severe patients

		Severe	Non-severe								
Author(s) and Year	n	mean	n	mean				Stan	dardized	d Mean	Difference [95% CI]
Chen, 2020	11	1.886	10	1.326			<u> </u>				0.86 [-0.04, 1.76]
Feng, 2020	20	2.326	94	1.934			<del></del>	<b>—</b>			0.33 [-0.22, 0.88]
He, 2020	69	1.559	135	1.693			-				-0.09 [-0.41, 0.22]
Nie, 2020	25	1.639	72	1.515				<b>-</b>			0.19 [-0.34, 0.72]
Shi, 2020	21	1.913	22	1.695		٠	<del></del>	-			0.07 [-0.53, 0.67]
Song, 2020	42	1.181	31	1.291		-	———				-0.08 [-0.55, 0.39]
Wan, 2020	21	1.47	102	1.564			<del></del>				-0.48 [-1.06, 0.11]
Wang, 2020	7	1.876	36	1.44				•	<b>⊸</b>		0.61 [-0.42, 1.64]
Xu, 2020	45	1.691	80	1.762			<b>⊢</b>				-0.06 [-0.43, 0.31]
Zhang, Zhou, 2020	21	1.803	44	1.697		٠	<del>-</del>	<b>⊣</b>			0.05 [-0.53, 0.63]
Zhang, Wang, 2020	14	1.206	29	1.344		-	<del></del>	ı			-0.14 [-0.77, 0.50]
Zheng, 2020	74	1.411	22	1.524		٠	<del>-</del>				-0.06 [-0.55, 0.42]
RE Model (Q = 10.17, df = 11, p = 0.515; I <sup>2</sup> = 0.0%)							+				0.00 [-0.15, 0.15] p = 0.9962
							<del>i</del>				
					-2	-1	0	1	2	3	
	CD4+/CD8+ T-cell Ratio SMD										

Figure S 3: Standardized Mean Difference of CD4+/CD8+ T-cell count ratio across severe and non-severe patient groups




Figure S 4: Meta-regression bubble plots of (a) IL-6 and (b) IL-10 SMD with neutrophil to lymphocyte ratio across severe and non-severe groups




Figure S 5: Meta-regression bubble plots of (a) IL-6 and (b) IL-10 SMD with CD4+/CD8+ T-cell count ratio across severe and non-severe groups




Figure S 6: Sensitivity analysis for (a) IL-6 and (b) IL-10 meta-analysis