

Supplementary Table 1. Characteristics of women tested positive for SARS-CoV-2 infection via RT-PCR

| Variable | Category | N (%) |
|--|--|--------------------|
| Study cohort | Sub-cohort 1 (Women delivering before 26/May/2020) | 6 |
| | Sub-cohort 2 (Women delivering from 26/May/2020 on) | 1 |
| | Sub-cohort 3 (Women in their 12 th week of pregnancy after 26/May/2020) | 1 |
| Partner RT-PCR | Negative | 5 |
| | Positive | 1 |
| | Missing | 2 |
| Age | <25 years | 0 |
| | 25-29 years | 0 |
| | 30-34 years | 3 (37.5) |
| | 35-39 years | 2 (25.0) |
| | ≥40 years | 3 (37.5) |
| Education level | Primary studies | 0 |
| | Secondary studies | 1 (12.5) |
| | FP | 2 (25.0) |
| | University studies | 5 (62.5) |
| Work status | Unemployed / no active worker | 1 (12.5) |
| | Employed | 7 (87.5) |
| | Student | 0 |
| Occupation | Health care worker | 3 (37.5) |
| | Working in restaurant or commerce | 3 (37.5) |
| | Others | 2 (25.0) |
| Fertilization type | Natural | 7 (87.5) |
| | Artificial insemination | 0 |
| | In vitro fertilization (own ovule) | 0 |
| | In vitro fertilization (ovodonation) | 0 |
| | Missing (still pregnant) | 1 (12.5) |
| Smoker in pregnancy | No | 8 (100.0) |
| | Yes | 0 |
| Alcohol consumption in pregnancy | No | 8 (100.0) |
| | Yes | 0 |
| Parity | 1 | 5 (62.5) |
| | 2 | 1 (12.5) |
| | ≥3 | 1 (12.5) |
| | Missing (still pregnant) | 1 (12.5) |
| Type of delivery | Eutocic | 3 (37.5) |
| | Caesarean rate | 2 (25.5) |
| | Instrumental | 2 (25.5) |
| | Still pregnant | 1 (12.5) |
| Age (years) | mean±sd (min, max) | 36.4±4.0 (30 – 40) |
| Gestational age low at work (weeks) | mean±sd (min, max) | 21.1±12.7 (0 – 32) |

| | | |
|--|--------------------|------------------------|
| Pregestational BMI (kg/m²) | mean±sd (min, max) | 22.3±2.9 (18.5 – 28.0) |
| Gestational weight gain (kg) | mean±sd (min, max) | 10.9±2.8 (8 – 16) |
| Symptoms associated with COVID-19 | Fever | 0 |
| | Chills | 0 |
| | Sore throat | 0 |
| | Cough | 0 |
| | Short of breathing | 1 |
| | Headache | 0 |
| | Nausea | 1 |
| | Loss of sense | 1 |

Supplementary Table 2. Characteristics of children born from women tested positive to SARS-CoV-2 infection using RT-PCR

| Variable | Category | N = 7 (%) |
|--|--------------------------------|-------------------------|
| Gestational age at birth | <34 weeks | 0 |
| | 34-36 weeks | 0 |
| | ≥37 weeks | 7 (100.0) |
| Birth weight | <2500 g | 0 |
| | 2500-4000 g | 7 (100.0) |
| | >4000 g | 0 |
| Gender | Women | 3 (42.9) |
| | Men | 4 (57.1) |
| Twin or singleton | Singleton | 7 (100.0) |
| | Twin | 0 |
| Breastfeeding at hospital discharge | Exclusive breastfeeding | 2 (28.6) |
| | Mixed | 3 (42.9) |
| | Infant formula | 2 (28.6) |
| Neonatology admission | No | 6 (85.7) |
| | Yes | 1 (14.3) |
| Birth size | mean±sd (min, max) | 49.1±2.4 (47 – 53) |
| Birth cephalic perimeter | mean±sd (min, max) | 34.6±2.0 (33 – 39) |
| Apgar 1' | mean±sd (min, max) | 8.3±1.9 (4 – 9) |
| Apgar 5' | mean±sd (min, max) | 9.4±0.8 (8 – 10) |
| pH at birth | mean±sd (min, max) | 7.27±0.04 (7.23 – 7.32) |
| SARS-CoV-2 via RT-PCR | | All negative |